

Subramanian, S.

Working Paper

A Re-scaled Version of the Foster-Greer-Thorbecke Poverty Indices based on an Association with the Minkowski Distance Function

WIDER Research Paper, No. 2004/10

Provided in Cooperation with:

United Nations University (UNU), World Institute for Development Economics Research (WIDER)

Suggested Citation: Subramanian, S. (2004) : A Re-scaled Version of the Foster-Greer-Thorbecke Poverty Indices based on an Association with the Minkowski Distance Function, WIDER Research Paper, No. 2004/10, The United Nations University World Institute for Development Economics Research (UNU-WIDER), Helsinki

This Version is available at:

<https://hdl.handle.net/10419/84649>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



United Nations
University

WIDER

World Institute for Development Economics Research

Research Paper No. 2004/10

**A Re-scaled Version
of the Foster-Greer-Thorbecke Poverty
Indices based on an Association
with the Minkowski Distance Function**

S. Subramanian

February 2004

Abstract

This note advances a family of poverty measures, Π_α , which are derived as simple, normalized Minkowski distance functions. The Π_α indices turn out to be the α th roots of the corresponding Foster, Greer and Thorbecke P_α indices. The re-calibration of P_α in terms of Π_α could have certain possible advantages, which are reviewed in the note. While the Π_α indices are not decomposable in the ordinarily understood sense of that term, they are amenable to the completely general decomposition procedure advanced by Shorrocks ('Decomposition Procedures for Distributional Analysis: A Unified Framework Based on the Shapley Value') and discussed, here, as an application in the poverty context.

Keywords: poverty measures, decomposition

JEL classification: I13

Copyright © UNU-WIDER 2004

* Madras Institute of Development Studies, Tamil Nadu, India; email: subbu@mids.ac.in

This study has been prepared within the UNU-WIDER project on 'New Directions in Development Economics', which is directed by Tony Addison

UNU-WIDER acknowledges the financial contributions to the research programme by the governments of Denmark (Royal Ministry of Foreign Affairs), Finland (Ministry for Foreign Affairs), Norway (Royal Ministry of Foreign Affairs), Sweden (Swedish International Development Cooperation Agency—Sida) and the United Kingdom (Department for International Development).

ISSN 1810-2611

ISBN 92-9190-589-5 (internet version)

Acknowledgements

A first draft of this paper was written in the course of a Visiting Fellowship at UNU-WIDER over April-June 2003. I would like to thank, without implicating, James Foster, Tony Shorrocks, and D. Jayaraj for helpful discussions of the subject of this note, and Adam Swallow for seeing it through.

The World Institute for Development Economics Research (WIDER) was established by the United Nations University (UNU) as its first research and training centre and started work in Helsinki, Finland in 1985. The Institute undertakes applied research and policy analysis on structural changes affecting the developing and transitional economies, provides a forum for the advocacy of policies leading to robust, equitable and environmentally sustainable growth, and promotes capacity strengthening and training in the field of economic and social policy making. Work is carried out by staff researchers and visiting scholars in Helsinki and through networks of collaborating scholars and institutions around the world.

www.wider.unu.edu

publications@wider.unu.edu

UNU World Institute for Development Economics Research (UNU-WIDER)
Katajanokanlaituri 6 B, 00160 Helsinki, Finland

Camera-ready typescript prepared by Liisa Roponen at UNU-WIDER
Printed at UNU-WIDER, Helsinki

The views expressed in this publication are those of the author(s). Publication does not imply endorsement by the Institute or the United Nations University, nor by the programme/project sponsors, of any of the views expressed.

1 Introduction

This is a simple note which addresses, by way of a possibly useful *curiosum*, a few simple issues in the measurement of poverty. First, a family of poverty indices, Π_α ($\alpha \geq 1$), is derived in a very elementary way as a set of normalized (Minkowski) distance functions. The indices in this family turn out, simply but interestingly, to be strictly concave transforms of the corresponding—and well-known—Foster, Greer and Thorbecke (hereafter FGT 1984) class of poverty measures P_α . Typically, the magnitudes of the P_α indices—especially for ‘larger’ values of α —are somewhat ‘inconveniently’ small; and the effect of the re-scaling, or ‘re-cardinalization’, of the P_α measures in terms of the Π_α measures is to provide a set of indices of more tractably larger magnitude. The Π_α indices inherit most of the important properties of the corresponding ‘parental’ P_α indices, except that the former are not *decomposable* in the FGT sense. However, the Π_α measures *are* amenable to a completely general decomposition procedure proposed by Shorrocks (1999), and this—a second issue—is also discussed.

2 Poverty measures and their properties

For every positive integer n , let \mathbf{X}_n be the set of all non-negative n -vectors of income $\mathbf{x} = (x_1, \dots, x_i, \dots, x_n)$, where x_i is the income of the i th individual ($i = 1, \dots, n$). \mathbf{X} will be taken to be the set $\cup_{n=1}^{\infty} \mathbf{X}_n$. The poverty line, which separates the poor from the non-poor, is a strictly positive level of income denoted by z . \mathbf{R} is the set of reals and \mathcal{S} the set of positive reals. A *poverty index* is a mapping $P: \mathbf{X} \times \mathcal{S} \rightarrow \mathbf{R}$ such that, for every combination of \mathbf{x} and z in its domain, P specifies a unique real number describing the extent of poverty in the regime $(\mathbf{x}; z)$. For every $(\mathbf{x}; z) \in \mathbf{X} \times \mathcal{S}$, $q(\mathbf{x}; z)$ will stand for the number of poor persons, that is, for the cardinality of the set $\{i \mid x_i < z\}$.

A number of axioms for poverty measurement have been advanced in the literature. The following is a quick and informal description of some salient axioms. *Focus* requires the poverty index to be invariant with respect to increases in non-poor incomes; *symmetry* requires that the personal identities of individuals should not matter in assessing the extent of poverty; *normalization* requires that the poverty index be bounded from below by zero (corresponding to the case of ‘no poverty’); *continuity* requires the poverty index to be continuous on the sub-vector of poor incomes; *scale-invariance* requires that for all $(\mathbf{x}; z) \in \mathbf{X} \times \mathcal{S}$, $P(\mathbf{x}; z) = P(\beta\mathbf{x}; \beta z)$ for any positive scalar β ; *replication-invariance* requires that for all $(\mathbf{x}; z) \in \mathbf{X} \times \mathcal{S}$ and any positive integer k , $P(\mathbf{x}; z) = P(k\mathbf{x}; z)$; *monotonicity* requires that, other things equal, an increase in any poor person’s income should cause poverty to decline; *transfer* requires that, *ceteris paribus*, any rank-preserving transfer of income from a poor person to a poorer person should reduce poverty; *transfer-sensitivity* requires that, everything else remaining the same, a progressive rank-preserving transfer between two poor persons a fixed income apart should cause poverty to decline by more the poorer the pair of persons involved in the transfer (Foster 1984); *subgroup consistency* (Foster and Shorrocks 1991) requires that, *ceteris paribus*, an increase in any subgroup’s poverty should increase aggregate poverty; and *decomposability*, which is a strengthened version of subgroup consistency, requires the poverty index to be capable of being expressed as a population-share weighted sum of subgroup poverty levels (FGT 1984).

3 Measuring poverty as a normalized distance function

We shall here consider a class of poverty indices called the Π_α class. Each member of the Π_α family can be straightforwardly derived as a specific type of normalized *distance function* (on the construction of distance functions between income distributions, the reader is referred to Shorrocks 1982, Ebert 1984, and Chakravarty and Dutta 1987). To this end, consider a non-decreasingly ordered non-negative n -vector of incomes $\mathbf{x} = (x_1, \dots, x_q, x_{q+1}, \dots, x_n)$, where—to recall— x_i is the income of the i th poorest person ($i = 1, \dots, n$), $x_i \leq x_{i+1}$ ($i = 1, \dots, n-1$), and q is the number of *poor* persons (so $x_q < z$ and $x_{q+1} \geq z$). Given \mathbf{x} and z , we define the following three n -vectors: $\mathbf{z} = (z, \dots, z)$, $\mathbf{x}^c = (x_1, \dots, x_q, z, \dots, z)$, and $\mathbf{0} = (0, \dots, 0)$. The vector \mathbf{z} is one in which every income level is just the poverty line income: it is the distribution with the smallest mean income that is compatible with a complete absence of poverty. The vector \mathbf{x}^c is what Takayama (1979) has called a ‘censored’ income vector, and is obtained from the distribution \mathbf{x} by replacing all the non-poor incomes in \mathbf{x} by the poverty line income z . The vector $\mathbf{0}$ is one in which every person receives zero income: it represents the case of ‘total poverty’. A fairly natural way of measuring the extent of poverty in the income vector \mathbf{x} , given that the poverty line is z , would be to identify it with the ‘shortfall’ or ‘distance’ of \mathbf{x}^c from \mathbf{z} , that is to say, with the distance of the ‘actual’ situation from one that corresponds to the most efficiently achieved ‘ideal’ of ‘no poverty’. (Implicit in this interpretation is respect for the *focus axiom* of poverty measurement which treats the magnitude of non-poor incomes as being irrelevant for the assessment of the extent of poverty associated with an income vector). To obtain a *normalized* picture of the extent of poverty, all one has to do is to divide the distance of \mathbf{x}^c from \mathbf{z} by the distance of $\mathbf{0}$ from \mathbf{z} : the latter distance represents the case of ‘maximal deficit’, that is to say, the distance of the ‘total poverty’ situation from the ‘no poverty’ situation. What remains to be done is to specify the form of the relevant distance function. A very natural set of candidates, in this context, is the well-worn *Minkowski class of α -distance functions*, written as D_α^M , of which the ‘city-block’ function and the *Euclidean norm* are special cases, realized for $\alpha = 1$ and 2 respectively (see Wilson and Martinez 1997). Given any two vectors \mathbf{a} and \mathbf{b} belonging to n -dimensional Euclidean space, the distance between the two, in terms of the Minkowski α -distance functions, is given by

$$(1) \quad D_\alpha^M(\mathbf{a}, \mathbf{b}) = [\sum_{i=1}^n |a_i - b_i|^\alpha]^{1/\alpha}, \quad \alpha \geq 1$$

In view of the preceding discussion, and given \mathbf{x} and z , a fairly straightforward expression for a class of normalized distance measures of poverty— Π_α —would be given by

$$(2) \quad \Pi_\alpha(\mathbf{x}; z) = D_\alpha^M(\mathbf{z}, \mathbf{x}^c) / D_\alpha^M(\mathbf{z}, \mathbf{0}), \quad \alpha \geq 1$$

Given (2), (3) can be written, with a little bit of manipulation, as

$$(3) \quad \Pi_\alpha(\mathbf{x}; z) = [(1/n) \sum_{i=1}^q \{(z - x_i)/z\}^\alpha]^{1/\alpha}, \quad \alpha \geq 1$$

The FGT (1984) family of poverty indices P_α , ($\alpha \geq 0$) is given by

$$(4) \quad P_\alpha(\mathbf{x}; z) = (1/n) \sum_{i=1}^q \{(z - x_i)/z\}^\alpha$$

Comparing (3) and (4), it is immediate that, for all $(\mathbf{x};z) \in \mathbf{X} \times \mathbf{S}$

$$(5) \quad \Pi_{\alpha}(\mathbf{x};z) = [P_{\alpha}(\mathbf{x};z)]^{1/\alpha}, \alpha \geq 1$$

It thus turns out that each member of the family of indices we have called Π_{α} is just the α th root of the corresponding member of the P_{α} family of indices. This is in itself a possibly small, but nevertheless, interesting result. Each member of the Π_{α} class is just a continuous, increasing and concave (strictly concave for $\alpha > 1$) transform of the counterpart member of the P_{α} class: it should not be surprising that the two classes of indices share their respective properties. (Indeed, Π_1 and P_1 are the same index). Specifically (see FGT 1984), the Π_{α} indices like the P_{α} indices satisfy focus, symmetry, normalization, continuity, scale- and replication-invariance, monotonicity, and subgroup consistency for all $\alpha \geq 1$; transfer for all $\alpha > 1$; and transfer-sensitivity for all $\alpha > 2$. Where the two sets of indices part company is in the matter of satisfying the decomposability property: every member of the P_{α} family is decomposable, while no member of the Π_{α} family is. We shall return to this issue later.

Setting aside for the moment the question of decomposability, are there any reasons for favouring the Π_{α} class over the P_{α} class? One can think of two reasons, each of which has something to commend it, though neither may be unqualifiedly compelling. The first has to do with a matter of interpretation: motivationally, the Π_{α} family of indices makes a direct appeal to intuition by presenting a poverty measure as a normalized distance measure, while the P_{α} family presents the poverty measure as a weighted sum of poverty-gaps, with the weights assuming the form of the poverty gaps themselves raised to a non-negative power, the basis for which is open to the (mild) criticism of a degree of arbitrariness. The second reason has to do with what, very informally, one may call ‘the psychology of numbers’. Take an index like P_3 , for instance. Inasmuch as it satisfies a number of desirable properties, including and in particular that of transfer-sensitivity, one would imagine that it would be widely employed in empirical work. The fact that it is not, one strongly suspects, is because, typically, P_3 yields ‘very small’ values when computed for actual empirical distributions of income. Consider, for instance, the vector $\mathbf{u} = (7,8,9,10,10,10,10,10,10)$ in a situation where $z = 10$, so that $q = 3$ and $n = 10$. It can be verified that $P_3(\mathbf{u};z) = 0.0036$: this is an ‘inconveniently small’ number, unlike $\Pi_3(\mathbf{u};z)$ which yields a ‘healthier-looking’ value of 0.1561. Indeed, this could be a matter of some practical (computational) import, as reflected in the following numerical example. Assuming that $z = 10$ and $n = 10$, let \mathbf{r} , \mathbf{s} and \mathbf{t} be three income vectors given, respectively, by $\mathbf{r} = (6,7,8,9,10,10,10,10,10,10)$, $\mathbf{s} = (6,7.1,7.9,9,10,10,10,10,10,10)$, and $\mathbf{t} = (6,7,8.1,8.9,10,10,10,10,10,10)$. It can be seen that each of \mathbf{s} and \mathbf{t} is derived from \mathbf{r} through a progressive transfer, and transfer sensitivity would require that $P(\mathbf{s};z) < P(\mathbf{t};z)$. Suppose we measure poverty correct to four decimal places. Then, it can be verified that $P_3(\mathbf{s};z) = P_3(\mathbf{t};z) = 0.0099$: P_3 is unable to differentiate between \mathbf{s} and \mathbf{t} , and one has to proceed to the fifth decimal place in order to secure the required discrimination ($P_3(\mathbf{s};z) = 0.00987 < P_3(\mathbf{t};z) = 0.00992$); on the other hand, Π_3 does the job when poverty is measured correct to four decimal places ($\Pi_3(\mathbf{s};z) = 0.2145 < \Pi_3(\mathbf{t};z) = 0.2149$).

The question remains whether the instrumental advantages discussed above in favour of Π_{α} outweigh the fact that no member of this class is decomposable, as that term is ordinarily understood. It may be as well here to formally define ‘decomposability’ after

the FGT (1984) fashion. Let $M = \{1, \dots, j, \dots, m\}$ be a set of m mutually exclusive and exhaustive subgroups into which a population of n individuals is partitioned. Let P be a measure of aggregate poverty for the population, and let P_j (respectively, t_j) be the poverty level (respectively, population share) of subgroup j , for every j belonging to M . The poverty index P may be written as a function f of the subgroup poverty levels

$$(6) \quad P = f(P_1, \dots, P_j, \dots, P_m)$$

Decomposability (or *Axiom D*), as FGT (1984) define it, requires the function f in (6) to take the form of a population share weighted sum of subgroup poverty levels, namely $P = \sum_{j \in M} t_j P_j$. Axiom D is obviously of considerable value in facilitating qualitative and quantitative assessments of the contributions of different subgroups to aggregate poverty, and, as such, holds a very useful key to policy analysis. For instance, it is common practice to interpret the decomposability of the poverty index P as suggesting that, for every $j \in M$, the contribution of subgroup j to total poverty is just $t_j P_j$. This is clearly useful information, and in particular not the sort of information one might wish to lose in the cause of a poverty index belonging to the Π_α family, despite the two reasons mentioned earlier in favour of this class of indices. This issue is investigated further in the next section.

4 Shorrocks-Shapley decomposition

The necessity of having to *choose* between indices which belong, respectively, to the Π_α and P_α families, would not arise if only there were available a workable and intuitively acceptable notion of ‘decomposability’ which is not simply *defined* to coincide with the demand made by Axiom D. Fortunately, such a notion of decomposability does exist, and can be found in Shorrocks’ regrettably unpublished paper (Shorrocks 1999). In this work Shorrocks presents a completely general technique of decomposition, based on Shapley’s (1953) formulation of the ‘Shapley value’ solution to the problem of allocation of output/costs among contributors/beneficiaries within the setting of an n -person cooperative game. Shorrocks’ adaptation of the Shapley value approach leads to a formulation of a general decomposition procedure which will here be called *Shorrocks-Shapley decomposition*, or SS-decomposition, for short. As applied specifically to the decomposition of a poverty index into subgroup poverty contributions, SS-decomposition can be described along the following lines (this description is heavily dependent on Shorrocks 1999).

We begin by recalling equation (6), in which aggregate poverty P is written as a function f of the set of m subgroup poverty levels $\{P_j\}_{j \in M}$. Let S be any subset of M , and define $F(S)$ to be the value which P assumes when all subgroup poverty levels P_j , $j \notin S$, have been eliminated from consideration, viz. $F(S) = f(\{P_j\}_{j \in S})$, $S \subseteq M$. Call the pair $\langle M, F \rangle$ a *model*, and given the model $\langle M, F \rangle$, a *decomposition* of the model is a set of real numbers $\{C_j(M, F)\}_{j \in M}$, such that, for every $j \in M$, C_j represents the contribution of subgroup j ’s poverty to total poverty. A *decomposition rule* C is a function which, for every model $\langle M, F \rangle$, assigns a set of subgroup poverty contributions $\{C_j(M, F)\}_{j \in M}$. It would be desirable for the decomposition rule to be (a) *exact* (which is the requirement that the subgroup poverty contributions add up to total poverty, namely $\sum_{j \in M} C_j(M, F) = F(M) (\equiv P)$); and (b) *symmetric* (which is the requirement that subgroup

contributions are invariant with respect to all possible permutations of subgroup labels). A decomposition rule which satisfies these properties is the Shorrocks-Shapley rule, the content of which can be explained as follows.

The basic idea here is to consider the marginal impact of each of the subgroups' poverty levels when the subgroups are eliminated in sequence. To do this, we first denote by Λ the set of all one-to-one mappings from M to itself. Every $\lambda \in \Lambda$ can then be interpreted as a particular sequence in which the subgroups in M are eliminated. For every $j \in M$ and $\lambda \in \Lambda$, let $S(j, \lambda)$ [respectively, $S'(j, \lambda)$] be the set of subgroups left after eliminating all the subgroups that *come before* subgroup j [respectively, the set of subgroups left after eliminating all the subgroups that *do not come after* subgroup j] according to the sequence represented by λ . That is, $S(j, \lambda) \equiv \{k \in \lambda(M) \mid k \geq j\}$ and $S'(j, \lambda) \equiv \{k \in \lambda(M) \mid k > j\}$. If we take away $F(S'(j, \lambda))$ from $F(S(j, \lambda))$, then it makes intuitive sense to interpret the remainder as the marginal impact of subgroup j 's poverty when λ represents the sequence of subgroup elimination; this marginal impact is denoted by C_j^λ

$$(7) \quad C_j^\lambda = F(S(j, \lambda)) - F(S'(j, \lambda))$$

It can be checked that $\sum_{j \in M} C_j^\lambda = F(M) (\equiv P)$, so (7) is an exact decomposition. To ensure *symmetry*, one can assign equal weight to the marginal impacts derived from different sequences of subgroup elimination, by taking the simple average of the marginal impacts corresponding to the various sequences of elimination. Specifically, with m subgroups, the cardinality of the set Λ will be $m!$; and a symmetric (in the sense of '[elimination] path independent') decomposition rule, C^S , is given by

$$(8) \quad C_j^S(M, F) = (1/m!) \sum_{\lambda \in \Lambda} C_j^\lambda$$

where C_j^λ is as defined in (7). From (8), it can be verified that C^S is an exact decomposition rule. Finally, since the decomposition rule is exact, it is meaningful to speak of the 'proportionate contribution' of each subgroup's poverty to total poverty: for every $j \in M$, this proportionate contribution will be given by $c_j^S \equiv C_j^S/P$. This is the substance of the SS-decomposition procedure, as applied to the present context.

Shorrocks (1999) has shown that the SS-decomposition of any poverty index belonging to the family P_α , will yield a contribution for the j th subgroup which is given by

$$(9) \quad C_j^S(M, F) = t_j P_{\alpha j}$$

This, precisely, is also the customary interpretation. What is the contribution of subgroup j 's poverty to poverty as measured by an index belonging to the Π_α family when the latter is subjected to SS-decomposition? Given (5), and noting that the FGT indices satisfy Axiom D, it is immediate that

$$(10) \quad \Pi_\alpha(\mathbf{x}; z) [= (P_\alpha(\mathbf{x}; z))^{1/\alpha} = (\sum_{j \in M} t_j P_{\alpha j})^{1/\alpha}] = (\sum_{j \in M} t_j \Pi_{\alpha j}^\alpha)^{1/\alpha}, \alpha \geq 1$$

For specificity, and by way of illustration, we consider the simple case of $m = 2$. There are two possible sequences of subgroup elimination available now: In the first sequence, subgroup 1 is first eliminated, followed by subgroup 2; in the second

sequence, subgroup 2 is first eliminated, followed by subgroup 1. For the first elimination sequence, the contribution of subgroup 1 is given by $F(M) - F(\{2\})$, and for the second elimination sequence, the contribution of subgroup 1 is given by $F(\{1\})$; averaging over the two sequences gives the SS-contribution of group 1: $C_1^S = (1/2)[F(M) - F(\{2\}) + F(\{1\})]$. Noting that $F(M) \equiv \Pi_\alpha$, $F(\{2\}) \equiv (t_2)^{1/\alpha}\Pi_{\alpha 2}$, and $F(\{1\}) \equiv (t_1)^{1/\alpha}\Pi_{\alpha 1}$, we have

$$(11) \quad C_1^S = (1/2)[\Pi_\alpha - (t_2)^{1/\alpha}\Pi_{\alpha 2} + (t_1)^{1/\alpha}\Pi_{\alpha 1}]$$

By virtue of the exactness property of the SS-decomposition rule, the contribution of subgroup 2 will just be total poverty *less* subgroup 1's contribution; or, in view of (11)

$$(12) \quad C_2^S = (1/2)[\Pi_\alpha - (t_1)^{1/\alpha}\Pi_{\alpha 1} + (t_2)^{1/\alpha}\Pi_{\alpha 2}]$$

Given (11) and (12), we are assured that the use of an index belonging to the Π_α family is in no way compromised by the possibility that we may not be able to infer subgroup contributions to total poverty: the Shorrocks-Shapley approach to decomposition has laid that problem to rest. Indeed, as Shorrocks (1999: 1) says:

[A] problem with conventional procedures is that they often place constraints on the kinds of poverty and inequality indices which can be used. Only certain forms of indices yield a set of contributions that sum up to the amount of poverty or inequality that requires explanation.

Finally, if one is persuaded that there is a case for favouring Π_α over P_α , then this could also have non-trivial implications for between-group allocations of anti-poverty budgets when such allocations are made, in line with a not unreasonable rule-of-thumb, according to the proportionate contributions of subgroups to aggregate poverty. By way of illustration, consider again the case in which $m = 2$ and, for specificity, $\alpha = 2$. Then, if $P_{21} = 0.05$, $P_{22} = 0.025$, and $t_1 = 0.1$, using (9) and (11) will enable confirmation of the fact that the proportionate contribution of subgroup 1 to total poverty is (a) 18.18 per cent when poverty is measured by P_2 , and (b) 26.08 per cent when poverty is measured by Π_2 . The difference between getting an 18 per cent and a 26 per cent share of the budgetary outlay is presumably non-negligible.

5 Summation

To summarize and conclude: a family of poverty indices Π_α which turn out to be just the α th roots of the corresponding members of the Foster-Greer-Thorbecke family P_α , has been derived as a set of plausible, normalized (Minkowski) distance functions. The re-scaled version Π_α of P_α , unlike the latter, is not 'decomposable' in the usually accepted sense of that term, that is, it cannot be written as a population share weighted sum of subgroup poverty levels. This, however, need not be a problem for Π_α : Shorrocks's (1999) work on what one may call Shorrocks-Shapley decomposition furnishes a completely general solution to the decomposition problem, in terms of which subgroup contributions to total poverty can be computed even for poverty indices (like Π_α) which cannot be expressed as population share weighted sums of subgroup poverty levels. This frees up poverty analysts to avail themselves of such advantages—

in terms of directness of interpretation and the convenience afforded by a re-calibration of the scale on which poverty is measured—as may be had from a ‘concavification’ of the P_α indices in terms of the Π_α indices. Finally, the switch from P_α to Π_α could also make for a non-trivial difference at the margin if subgroup contributions to poverty are invested with more than ordinal significance.

References

- Chakravarty, S. R., and B. Dutta (1987). ‘A Note on Measures of Distance between Income Distributions’. *Journal of Economic Theory*, 41.
- Ebert, U. (1984). ‘Measures of Distance between Income Distributions’. *Journal of Economic Theory*, 32.
- Foster, J. (1984). ‘On Economic Poverty: A Survey of Aggregate Measures’. *Advances in Econometrics*, 3.
- Foster, J., J. Greer, and E. Thorbecke (1984). ‘A Class of Decomposable Poverty Measures’. *Econometrica*, 42.
- Foster, J., and A. F. Shorrocks (1991). ‘Subgroup Consistent Poverty Indices’. *Econometrica*, 59.
- Sen, A. K. (1976). ‘Poverty: An Ordinal Approach to Measurement’. *Econometrica*, 44.
- Shapley, L. (1953). ‘A Value for n-person Games’, in H. W. Kuhn and A. W. Tucker (eds), *Contributions to the Theory of Games*, Vol. 2. Princeton: Princeton University Press.
- Shorrocks, A. F. (1982). ‘On the Distance Between Income Distributions’. *Econometrica*, 50.
- Shorrocks, A. F. (1999). ‘Decomposition Procedures for Distributional Analysis: A Unified Framework Based on the Shapley Value’. Wivenhoe Park: University of Essex and Institute for Fiscal Studies. Mimeo.
- Takayama, N. (1979). ‘Poverty, Income Inequality, and Their Measures: Professor Sen’s Axiomatic Approach Reconsidered’. *Econometrica*, 47.
- Wilson, D. R., and T. R. Martinez (1997). ‘Improved Heterogeneous Distance Functions’. *Journal of Artificial Intelligence Research*, 6.