

MacKenzie, C. R.; MacAleer, Michael

**Working Paper**

## Comparing tests of autoregressive versus moving average errors in regression models using Bahadur's asymptotic relative efficiency

ISER Discussion Paper, No. 537

**Provided in Cooperation with:**

The Institute of Social and Economic Research (ISER), Osaka University

*Suggested Citation:* MacKenzie, C. R.; MacAleer, Michael (2001) : Comparing tests of autoregressive versus moving average errors in regression models using Bahadur's asymptotic relative efficiency, ISER Discussion Paper, No. 537, Osaka University, Institute of Social and Economic Research (ISER), Osaka

This Version is available at:

<https://hdl.handle.net/10419/92622>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

**COMPARING TESTS OF AUTOREGRESSIVE  
VERSUS MOVING AVERAGE ERRORS  
IN REGRESSION MODELS USING BAHADUR'S  
ASYMPTOTIC RELATIVE EFFICIENCY**

C. R. McKenzie  
and  
Michael McAleer

May 2001

The Institute of Social and Economic Research  
Osaka University  
6-1 Mihogaoka, Ibaraki, Osaka 567-0047, Japan

# COMPARING TESTS OF AUTOREGRESSIVE VERSUS MOVING AVERAGE ERRORS IN REGRESSION MODELS USING BAHADUR'S ASYMPTOTIC RELATIVE EFFICIENCY

C. R. McKenzie

Osaka School of International Public Policy, Osaka University

Michael McAleer

Department of Economics, University of Western Australia

Revised: May 2001

## Abstract

The purpose of this paper is to use Bahadur's asymptotic relative efficiency measure to compare the performance of various tests of autoregressive (AR) versus moving average (MA) error processes in regression models. Tests to be examined include non-nested procedures of the models against each other, and classical procedures based upon testing both the AR and MA error processes against the more general autoregressive-moving average model.

**KEYWORDS:** Autoregressive model; Bahadur efficiency; inappropriate alternatives; Lagrange multiplier test; moving average model; separate (non-nested) tests.

## 1. INTRODUCTION

In contrast to the numerous empirical studies which have reported estimates of regression models with autoregressive errors, there have been relatively few studies reporting estimates of regression models with moving average errors. This situation prevails in spite of the computational ease with which regression models with moving average errors can be estimated by maximum likelihood (ML) methods (see, for example, Pesaran and Pesaran [1] and Quantitative Micro Software EViews 4.0 [2]), or by asymptotically efficient two step methods (see Reinsel [3] and Hannan et al. [4]). The Lagrange multiplier (LM) test for serial correlation can detect serial correlation but provides no indication of whether the serial correlation in a regression model arises from an autoregressive process or a moving average process (see, for example, Godfrey [5]). The major aim of this paper is to compare some tests that will enable a choice to be made between the two error processes in a regression model using Bahadur's asymptotic relative efficiency measure.

Moving average (MA) errors are a viable alternative to autoregressive (AR) errors in a regression model since there are strong a priori reasons to expect the errors in certain models to have an MA form. For example, MA errors in a regression context may arise when: (i) there are random measurement errors associated with the dependent variable and lagged dependent variables appear as explanatory variables (Walker [6] and Pesaran [7]); (ii) the forecasting period exceeds the sampling period in forecasting equations (Hansen and Hodrick [8]); (iii) the equation is a solution of a rational expectations model (Broze et al. [9] and Evans and Honkapohja [10]); (iv) the model is a discrete time approximation to a continuous time model (Bergstrom [11]); (v) Koyck lag distributions are employed (Chow [12, pp. 102-103]); (vi) overlapping data on the dependent variable are used (Rowley and Wilton [13] and Kenward [14]); (vii) structural time series models are used (Harvey and Todd [15]); (viii) the data have been adjusted using filters such as X-11 (Wallis [16]); or (ix) an error correction model is estimated for series that are cointegrated (Engle and Granger [17]). Nicholls et al. [18] and Schwert [19, pp. 77-78] indicate a number of other instances where moving average errors can also be expected.

On the other hand, AR errors in a regression context may arise from common factor restrictions in dynamic models (see Hendry and Mizon [20] and Sargan [21]) and from

stock adjustment models, or they may be indicative of general misspecification, especially in the form of the exclusion of important explanatory variables. If the excluded variables have the typical spectra of economic variables (Granger [22]), then the errors are likely to be AR. Alternatively, AR errors may be assumed because the consequences of doing so, even when the errors follow an MA error process, may not be too severe (see Griffiths and Beesley [23] for the case of a first-order MA, or MA(1), process).

Justifications for autoregressive-moving average (ARMA) errors rest on a combination of the reasons for AR and MA errors. The principle of parsimony, or Occam's razor (see, for example, Zellner et al. [24]), may lead to a simple AR or MA representation as an approximation to a higher-order ARMA process (see Box and Jenkins [25] and Hendry and Trivedi [26]).

The problem of testing between AR and MA models has been considered in the pure time series literature by Whittle [27], Walker [28], Pagan et al. [29], McAleer et al. [30], Hall and McAleer [31], Godfrey and Tremayne [32, 33], Franses [34], and Gouriéroux and Monfont [35]. There have been a few attempts to examine the problem of testing between two regression models with different error processes; for example, King and McAleer [36], Godfrey and Tremayne [32] and Burke et al. [37] test AR(1) errors against MA(1) errors, Silvapulle and King [38] test MA(1) errors against AR(1) errors, Silvapulle and King [39] test joint AR(1)-AR(4) disturbances against joint MA(1)-MA(4) disturbances, and McKenzie et al. [40] test AR(p) disturbances against MA(q) disturbances. The available Monte Carlo evidence for both the pure time series and regression cases suggests that non-nested tests of these two models can have high power against each other, as well as against inappropriate alternatives (see McAleer et al. [30], Hall and McAleer [31] and McKenzie et al. [40]).

As a complement to existing research which tests these non-nested models against each other, this paper uses Bahadur's asymptotic relative efficiency to compare various tests of AR versus MA errors in regression models against each other.

The plan of the paper is as follows. Section 2 contains details of the model being considered as well as definitions of a number of variables that are used in later sections.

Section 3 briefly reviews tests of the null hypothesis that the errors follow either an AR or an MA process against the non-nested alternative of an MA or an AR process, respectively, as well as diagnostic tests for the respective null hypotheses against higher-order ARMA processes. Section 4 provides a comparison of the Bahadur approximate slopes of the diagnostic and non-nested tests against appropriate and inappropriate fixed alternatives. The final section contains some concluding comments.

## 2. MODEL AND NOTATION

Consider the linear regression model

$$y_t = x_t' \beta + u_t, t = 1, \dots, T \quad (1)$$

where  $x_t$  is presumed to be a  $k$  by 1 vector of non-stochastic variables and  $\beta$  is a  $k$  by 1 vector of unknown parameters. Two alternative non-nested hypotheses concerning the nature of  $u_t$  are the AR( $p$ ) and MA( $q$ ) processes, namely:

$$H_0: u_t = \alpha_1 u_{t-1} + \dots + \alpha_p u_{t-p} + \varepsilon_t, \varepsilon_t \sim \text{NID}(0, \sigma^2) \quad (2)$$

$$H_1: u_t = \varepsilon_t - \beta_1 \varepsilon_{t-1} - \dots - \beta_q \varepsilon_{t-q}, \varepsilon_t \sim \text{NID}(0, \sigma^2). \quad (3)$$

The AR( $p$ ) process in (2) is assumed to be finite and stationary, and the MA( $q$ ) process in (3) is assumed to be finite and invertible. Provided  $\alpha_i \neq 0$  for at least one  $i$  ( $0 < i < p+1$ ) and  $\beta_j \neq 0$  for at least one  $j$  ( $0 < j < q+1$ ), the hypotheses  $H_0$  and  $H_1$  can be shown to be globally non-nested. Assuming fixed initial values for  $u_t$  ( $t = -p+1, \dots, 0$ ) in  $H_0$  and for  $\varepsilon_t$  ( $t = -q+1, \dots, 0$ ) in  $H_1$ , denote the ML estimates of (1) under  $H_0$  as  $(\hat{\beta}', \hat{\alpha}_1, \dots, \hat{\alpha}_p, \hat{\sigma}^2)$  with associated residuals  $\hat{u}_t = y_t - x_t' \hat{\beta}$ , and under  $H_1$  as  $(\tilde{\beta}', \tilde{\beta}_1, \dots, \tilde{\beta}_q, \tilde{\sigma}^2)$  with associated residuals  $\tilde{u}_t = y_t - x_t' \tilde{\beta}$ . The predictions and prediction errors from the two models are denoted by:

$$H_0: \hat{y}_t = x_t' \hat{\beta} + \hat{\alpha}_1 \hat{u}_{t-1} + \dots + \hat{\alpha}_p \hat{u}_{t-p} \quad \text{and} \quad \hat{\varepsilon}_t = y_t - \hat{y}_t,$$

and

$$H_1: \tilde{y}_t = x_t' \tilde{\alpha} - \tilde{\alpha}_1 \tilde{y}_{t-1} - \dots - \tilde{\alpha}_q \tilde{y}_{t-q} \text{ and } \tilde{\alpha}_t = y_t - \hat{y}_t.$$

For future reference, it will be useful to define the following polynomial lag functions:

$$\hat{L} = 1 - \hat{\alpha}_1 L - \dots - \hat{\alpha}_p L^p, \quad \tilde{L} = 1 - \tilde{\alpha}_1 L - \dots - \tilde{\alpha}_p L^p,$$

$$\hat{L} = 1 - \hat{\alpha}_1 L - \dots - \hat{\alpha}_q L^q, \quad \tilde{L} = 1 - \tilde{\alpha}_1 L - \dots - \tilde{\alpha}_q L^q,$$

and the following transformed variables:  $y_t^* = \hat{L} y_t$ ,  $x_t^* = \hat{L} x_t$ ,  $y_t^+ = \tilde{L}^{-1} y_t$ ,

$$x_t^+ = \tilde{L}^{-1} x_t, \quad \tilde{y}_t^+ = \tilde{L}^{-1} \tilde{y}_t, \text{ and } \tilde{u}_t^+ = \tilde{L}^{-1} \tilde{u}_t.$$

### 3. TESTS OF AR(p) VERSUS MA(q) ERRORS

#### 3.1 Testing $H_0: \text{AR}(p)$ by Variable Addition

The tests of the AR(p) null against the MA(q) alternative developed in McKenzie et al. [40] are based on the following auxiliary regression equation:

$$y_t^* = x_t^{*'} \alpha + \hat{u}_{t-1} + \dots + \hat{u}_{t-p} + Z_t' \gamma + w_t \quad (4)$$

where  $w_t = \tilde{y}_t + (\hat{L} - \tilde{L}) x_t' (\hat{\alpha} - \tilde{\alpha})$ ,  $\hat{u}_i = \hat{y}_i - \tilde{y}_i$ ,  $Z_t$  is an  $r$  by 1 vector of added variables that are asymptotically uncorrelated with  $\tilde{y}_t$ , and  $\gamma$  is an  $r$  by 1 vector of unknown parameters. Estimating (4) by OLS and denoting the F test of  $H_0: \gamma = 0$  by  $F$ , then  $rF \xrightarrow{d} \chi^2_{(r)}$  under  $H_0$  as  $T \rightarrow \infty$ .

The test associated with the hypothesis  $\gamma = 0$  in (4) can be computed as

$$rF = (ESS_R - ESS_U) / (ESS_U / T) \quad (5)$$

where  $ESS_R$  and  $ESS_U$  are the restricted and unrestricted error sums of squares from estimating (4) subject to  $\alpha = 0$  and  $\beta = 0$ , respectively. An identical procedure can be

obtained by subtracting  $\hat{x}_t^*$  from both sides of (4) to give

$$\hat{y}_t = \hat{x}_t^* (\hat{\alpha} - \hat{\beta}) + \hat{u}_{t-1} + \dots + \hat{u}_{t-p} + Z_t' \gamma + w_t. \quad (6)$$

Equation (6) is particularly useful for calculating the asymptotic relative efficiencies of the various tests. For testing the  $AR(p)$  null against the  $MA(q)$ ,  $AR(p+r)$  or  $ARMA(p, r)$  alternatives, the choices of  $Z_t$  are given in Table 1.

### 3.2 Testing $H_1: MA(q)$ by Variable Addition

The tests of the  $MA(q)$  null against the  $AR(p)$  alternative developed in McKenzie et al. [40] are based on the following auxiliary regression equation:

$$y_t^+ = x_t^+ \alpha + \tilde{u}_{t-1}^+ + \dots + \tilde{u}_{t-q}^+ + Z_t' \gamma + \epsilon_t \quad (7)$$

where  $\epsilon_t = y_t^+ - [ (L) - \tilde{(L)} ] \tilde{(L)}^{-1} (y_t^+ - \tilde{y}_t^+)$  and  $\tilde{u}_i = \tilde{y}_i - \tilde{y}_{i-1}$ . Estimating (7) by OLS, and

denoting the F test of  $H_0: \gamma = 0$  by  $F$ , then  $rF \xrightarrow{d} \chi^2_{(r)}$  under  $H_0$  as  $T \rightarrow \infty$ .

The test associated with the hypothesis  $\gamma = 0$  in (7) can be computed as  $rF$  in (5), where  $ESS_R$  and  $ESS_U$  are the restricted and unrestricted sums of squares from estimating (7) subject to  $\gamma = 0$  and  $\alpha = 0$ , respectively. An identical procedure can be obtained by

subtracting  $\tilde{x}_t^+$  from both sides of (7) to give

$$\tilde{y}_t = \tilde{x}_t^+ (\tilde{\alpha} - \tilde{\beta}) + \tilde{u}_{t-1}^+ + \dots + \tilde{u}_{t-q}^+ + Z_t' \gamma + \epsilon_t. \quad (8)$$

Equation (8) is particularly useful for calculating the asymptotic relative efficiencies of the various tests. For testing the  $MA(q)$  null against the  $AR(p)$ ,  $MA(q+r)$  or  $ARMA(p, q)$  alternatives, the choices of  $Z_t$  are given in Table 2.



#### 4. BAHADUR'S ASYMPTOTIC RELATIVE EFFICIENCY

Given the number of consistent tests in section 3, some criterion must be chosen to compare their performance. Since the AR(p) and MA(q) error processes are non-nested, non-local power comparisons are used to evaluate the tests. Specifically, Bahadur's [41, 42] asymptotic relative efficiency criterion is used for purposes of comparison (see Geweke [43, 44, 45], Pesaran [46], Wascher [47], Pesaran and Smith [48], and Zabel [49] for some econometric applications). In this context, since only the asymptotic properties of the estimators under both the null and alternative hypotheses are known, it is necessary to work with the approximate rather than the exact slopes of the tests, despite the difficulties noted in Geweke [43]. The difference between the tests labelled 1 and 2 in Tables 1 and 2 is whether the added variable is not modified before being included (method 1), or is modified by the AR transformation in Table 1 (method 2) or the MA transformation in Table 2 (method 2).

There are three important questions relating to the various tests: (i) For any given added variable, is any one of the testing methods clearly superior? (ii) For any given testing method, is any one of the added variables dominant? (iii) Is there an optimal test?

Bahadur's approach keeps the alternative hypothesis fixed and allows the probability of type I error (size) to tend to zero as the sample size increases. The slopes of the power functions can be approximated by evaluating the various statistics for testing the null hypothesis under the fixed alternative. In this section, the tests are compared using Bahadur's asymptotic relative efficiency for testing, in the context of the linear regression model, an AR(1) null hypothesis against an MA(1) alternative (section 4.1) and an MA(1) null hypothesis against an AR(1) alternative (section 4.2).

##### 4.1 Approximate Slopes of Tests of the AR(1) Model

A particularly useful property of the ML estimator  $\hat{\beta}$  is that  $\text{plim } \hat{\beta} = \beta$  regardless of whether the true disturbances follow an AR or an MA process. As  $x_t^*$  is asymptotically orthogonal to the remaining regressors in the equation for the LM, E and DOP tests, the

term  $x_t^*(\hat{\beta} - \beta)$  in (6) can be omitted without affecting the asymptotic properties of the test under either the null or the alternative hypothesis. Therefore, the LM, E and DOP statistics can be calculated from the auxiliary regression

$$\hat{u}_t = \hat{u}_{t-1} + \dots + \hat{u}_{t-p} + Z_t' \beta + w_{1t} \quad (9)$$

where  $w_{1t} = w_t + x_t^*(\hat{\beta} - \beta)$ . The Bahadur criterion requires an evaluation of the probability limit of the test statistic under the fixed alternative. When the linear regression model has been estimated assuming an AR(1) disturbance, the pseudo-true value of  $\beta$ ,  $\beta^*$ , is given, in general, as  $\beta^* = \text{plim}_1 \hat{\beta} = -\lambda/(1 + \lambda^2)$ , where  $\lambda$  is the true value of the MA(1) parameter and  $\text{plim}_1 \hat{\beta}$  denotes the probability limit of  $\hat{\beta}$  under  $H_1$ .

Using the test statistics calculated from (9), the Bahadur approximate slopes are computed as  $\text{plim}_1 rF/T$  for the various forms of the LM, E and DOP tests of the regression model with AR(1) errors, assuming a fixed alternative of a regression model with MA(1) errors. The probability limits of the approximate slopes of the tests will depend only on  $\beta^*$ , given the consistency of  $\hat{\beta}$  under both hypotheses.

Using the results in the Appendix, it can be shown that the approximate slope, AS, of each test defined using (9) can be computed as

$$AS = (ESS_R - ESS_U) / ESS_U, \quad (10)$$

where  $ESS_U$  is the unrestricted error sum of squares from one of the regressions defined in Table 3, and  $ESS_R = s_1' s_1$  (with  $s_1$  defined in Table 3). Let  $\tilde{s}_1' \tilde{s}_1$  be the unrestricted error sum of squares from a particular auxiliary regression arising from Table 3, so that the approximate slope of the test statistic is given by

$$AS = (s_1' s_1 - \tilde{s}_1' \tilde{s}_1) / \tilde{s}_1' \tilde{s}_1.$$

An interesting point to note from Table 3 is that the approximate slopes of E1, E2 and DOP1 are identical since the regressors used,  $a_2$ ,  $f_2$  and  $r_2$ , respectively, are linearly dependent.

Another test considered here is the adaptation by Burke et al. [37] of the test developed in Godfrey and Tremayne [32] for pure time series models for testing the null hypothesis of AR(1) errors against the alternative of MA(1) errors in a regression model. The test statistic is given by

$$= T^{1/2}(r_2 - r_1^2)/(1 - r_2) \quad (11)$$

where  $r_i \xrightarrow{d} N(0,1)$  under the null hypothesis of AR(1) errors, and  $r_i$  is an estimate of the  $i$ th-order autocorrelation coefficient ( $i=1,2$ ) calculated using OLS residuals. As the test has a negative mean under the MA(1) alternative, the AR(1) null is not rejected if  $> c$ , where  $c$  is such that  $\Pr(N(0,1) < c) = \alpha$  and  $\alpha$  is a given critical level. Since the theoretical autocorrelation coefficient for an MA(1) process cannot exceed 0.5, Burke et al. [37, p. 138] also suggest not rejecting the AR(1) model if  $r_1 > (0.5 + T^{-1/2})$ . However, the interpretation of this outcome should be that the MA(1) model is rejected, not that the AR(1) model is accepted. The reason for this interpretation is that there are many other stationary processes for which the theoretical autocorrelation coefficient at lag one can exceed 0.5, such as an MA(2) process. Under the MA(1) alternative,  $\text{plim}_1 T^{-1/2} = \theta^2/(1 + \theta^2)^2$ , where  $\theta$  is the true value of the MA(1) parameter (see Bera and Newbold [50]).

Figures 1 and 2 graph the approximate slopes of the tests for values of the true MA parameter between 0 and 0.9. Burke et al.'s [37] test is labelled TAU in Figure 1. In order that all these tests are distributed as  $\chi^2_{(1)}$  under the null hypothesis, the rule that the MA(1) model should be rejected when  $r_1 > (0.5 + T^{-1/2})$  is ignored in this analysis. One important result is that E1=E2=DOP1, so that no single testing method or added variable is

preferred. It is clear from these figures that the asymptotic relative efficiencies of the tests that use information from the alternative (E and DOP) are generally higher than the corresponding tests based on the LM principle (LMM and LMA). Moreover, in this simple case, of all the tests in the comparison, the E1 (=E2=DOP1) test would appear to be the best. The E1 test dominates Burke et al.'s [37] test.

#### 4.2 Approximate Slopes of Tests of the MA(1) Model

Since under  $H_0$  and  $H_1$ ,  $\text{plim } \tilde{\alpha} = \alpha$  and  $x_t^+$  is asymptotically orthogonal to all the other regressors in the equation for the LM, E and DOP tests,  $x_t^{+'}(\tilde{\alpha} - \alpha)$  in (8) can be omitted without affecting the asymptotic properties of the test under either the null or alternative hypotheses. Thus, the LM, E and DOP test statistics can be calculated from the auxiliary regression

$$\tilde{\epsilon}_t = \alpha_1 \tilde{\epsilon}_{t-1} + \dots + \alpha_q \tilde{\epsilon}_{t-q} + Z_t' \alpha + \epsilon_t \quad (12)$$

where  $\epsilon_t = \tilde{\epsilon}_t + x_t^{+'}(\tilde{\alpha} - \alpha)$ . When the linear regression model has been estimated assuming an MA(1) error, the pseudo-true values of  $\tilde{\alpha}$  and  $\tilde{\alpha}_0$  under  $H_0:AR(1)$  are  $\text{plim } \tilde{\alpha} = \alpha$  and  $\text{plim}_0 \tilde{\alpha} = \alpha^*$ , respectively, where  $\alpha^*$  is the solution to

$$\alpha^{*3} + 2\alpha^{*2} - \alpha^* - 1 = 0 \quad (13)$$

that lies in the interval  $(-1,1)$  (Whittle [27, p. 91]),  $\alpha$  is the true value of the AR(1) parameter, and  $\text{plim}_0 \tilde{\alpha}$  denotes the probability limit of  $\tilde{\alpha}$  under  $H_0$ . Thus, the probability limits for calculating the approximate slopes of the tests of  $H_1$  will depend only on  $\alpha$ .

Using the test statistics calculated from (12), the Bahadur approximate slopes are computed

as  $\text{plim}_0 rF/T$  for the various forms of the LM, E and DOP tests of the regression model with MA(1) errors, assuming a fixed alternative of a regression model with AR(1) errors. The probability limits of the approximate slopes of the tests will depend only on  $\rho$ , given the consistency of  $\hat{\rho}$  under both hypotheses.

Based on (12) and using the results in the Appendix, it can be shown that the approximate slope of each test defined using (12) can be computed as

$$AS = (\lim ESS_R - \lim ESS_U) / \lim ESS_U, \quad (14)$$

where  $ESS_U$  is the unrestricted error sum of squares from one of the regressions defined in Table 4,  $ESS_R$  is the error sum of squares from the regression of  $q$  on  $nq_1$  (with  $q$  and  $nq_1$  defined in the Appendix), and  $\lim$  denotes the limit as  $T \rightarrow \infty$ .

Figures 3-4 graph the approximate slopes of the tests for values of the true AR parameter between 0 and 0.9. It should be noted that the approximate slopes of LMA1, LMA2, LMM1, E2 and DOP2 are the same so that no single testing method or added variable is preferred. When they differ, the tests that use information from the alternative generally have higher approximate slopes than the corresponding tests based on the LM principle. In this case, the DOP1 test would appear to be the best of the tests in the comparison.

## 5. CONCLUSION

This paper has used Bahadur's asymptotic relative efficiency (ARE) to compare several simple procedures for the purpose of testing AR versus MA error processes in regression models. For the case of comparing a regression model with AR(1) errors as the null and MA(1) errors as the alternative, it was found on the basis of Bahadur's ARE criterion that the preferred test uses information from the alternative model through the prediction errors

or the difference in prediction errors of the two models. When the roles of the null and alternative hypotheses are reversed, so that the MA(1) model is the null and the AR(1) model is the alternative, the preferred test on the basis of Bahadur's ARE criterion uses information from the alternative model through the difference in the prediction errors.

## APPENDIX

Let A and B be T by T band matrices with elements  $a_{i,i} = 1$ ,  $a_{i+1,i} = 2, \dots, a_{i+k-1,i} = k$  and  $a_{i,j} = 0$  otherwise, and  $b_{i,i} = 1$ ,  $b_{i+1,i} = 2, \dots, b_{i+k-1,i} = k$  and  $b_{i,j} = 0$  otherwise, respectively. Denote the  $i$ th columns of A and B as  $\tilde{a}_i$  and  $\tilde{b}_i$ , respectively. In addition, the vectors  $\tilde{a}_i^j$  and  $\tilde{b}_i^j$  are  $j$  by 1 vectors with elements identical to the first  $j$  elements of  $\tilde{a}_i$  and  $\tilde{b}_i$ , respectively. Other vectors with a superscript are defined in a similar manner.

Lemma 1: (a) If  $k=T$  and  $\lim_{T \rightarrow \infty} \tilde{a}_1' \tilde{b}_1$  exists, then  $\lim_{T \rightarrow \infty} \text{tr}(A'B)/T = \lim_{T \rightarrow \infty} \tilde{a}_1' \tilde{b}_1$ .

(b) If  $k$  is fixed as  $T \rightarrow \infty$ ,  $\lim_{T \rightarrow \infty} \text{tr}(A'B)/T = \lim_{T \rightarrow \infty} \tilde{a}_1^k' \tilde{b}_1^k$ .

Proof: (a) If  $k=T$ , using the definitions of A and B,  $\text{tr}(A'B)$  is given by

$$\text{tr}(A'B) = \sum_{j=1}^T (T+1-j) \tilde{a}_j' \tilde{b}_j.$$

Taking limits as  $T \rightarrow \infty$  gives

$$\lim_{T \rightarrow \infty} \text{tr}(A'B)/T = \lim_{T \rightarrow \infty} \sum_{j=1}^T [1 - (j-1)/T] \tilde{a}_j' \tilde{b}_j.$$

The result follows from the properties of Cesaro sums (see Hatanaka [51]).

(b) Given (a), the case of fixed  $k$  is obvious.

QED

For the purpose of computing the Bahadur asymptotic relative efficiencies associated with tests of the AR(1) null under a fixed MA(1) alternative, and the MA(1) null under a fixed AR(1) alternative, it is useful to rewrite the models given in equations (1)-(3) in matrix

form for  $p=q=1$ . The regression model with AR(1) errors is rewritten as

$$y = X\beta + U, \quad R(\rho)U = e, \quad E(e) = 0, \quad V(e) = \sigma^2 I_T,$$

where  $y$ ,  $U$  and  $e$  are  $T$  by  $1$  vectors,  $X$  is a  $T$  by  $k$  matrix of non-stochastic variables,  $\beta$  is a  $k$  by  $1$  vector of unknown parameters, and  $R(\rho)$  is a  $T$  by  $T$  band matrix with  $r_{i,i} = 1$ ,  $r_{i,i-1} = \rho$  and  $r_{i,j} = 0$  otherwise. Denote the columns of  $R(\rho)$  by  $\tilde{r}_1(\rho)$ ,  $\tilde{r}_2(\rho)$ , ...,  $\tilde{r}_T(\rho)$ . The regression model with MA(1) errors is rewritten as

$$y = X\beta + U, \quad U = N(\theta)e, \quad E(e) = 0, \quad V(e) = \sigma^2 I_T,$$

where  $N(\theta)$  is a  $T \times T$  band matrix with  $a_{i,i} = 1$ ,  $a_{i,i-1} = \theta$  and  $a_{i,j} = 0$  otherwise. Denote the columns of  $N(\theta)$  by  $\tilde{a}_1(\theta)$ ,  $\tilde{a}_2(\theta)$ , ...,  $\tilde{a}_T(\theta)$ .

For the analysis that follows, define the  $T$  by  $1$  vectors  $U_j$ ,  $\hat{e}_j$ ,  $\hat{e}_j^+$ ,  $\hat{U}$ ,  $\hat{U}_j$ ,  $\tilde{U}$ ,  $\tilde{U}_j$ ,  $\tilde{e}_j$ ,  $\tilde{e}_j^+$ ,  $\tilde{e}_j^{++}$ ,  $\tilde{e}_j^{++}$  with typical elements  $u_{i-j}$ ,  $\hat{e}_i$ ,  $\hat{e}_{i-j}$ ,  $\hat{u}_i$ ,  $\hat{u}_{i-j}$ ,  $\tilde{u}_i$ ,  $\tilde{u}_{i-j}$ ,  $\tilde{e}_i$ ,  $\tilde{e}_{i-j}$ ,  $\tilde{e}_i^+$ ,  $\tilde{e}_{i-j}^+$  and  $\tilde{e}_{i-j}^{++}$  ( $j > 0$ ). Now  $\hat{e} = R(\rho)\hat{U}$  where  $\hat{U} = y - X\hat{\beta}$ , so that

$$\hat{e} = R(\rho)(y - X\hat{\beta}) = R(\rho)[U + X(\beta - \hat{\beta})] \quad (A1)$$

and  $\hat{U}_i$  can be written as

$$\hat{U}_i = y_i - X_i\hat{\beta} = U_i + X_i(\beta - \hat{\beta}) \quad (A2)$$

where  $y_i = L^i y$ ,  $X_i = L^i X$  and  $L$  is the lag operator. Similarly,  $\tilde{U} = N(\theta)\tilde{e}$  where  $\tilde{U} = y - X\tilde{\beta}$ , so that

$$\tilde{e} = N(\tilde{\cdot})^{-1}(y - X\tilde{\cdot}) = N(\tilde{\cdot})^{-1}[U + X(\tilde{\cdot} - \cdot)] \quad (A3)$$

and  $\tilde{e}_i^+$  can be written as

$$\tilde{e}_i^+ = N(\tilde{\cdot})^{-1}\tilde{e}_i = N(\tilde{\cdot})^{-2}[U_i + X_i(\tilde{\cdot} - \cdot)]. \quad (A4)$$

#### A. Tests of the AR(1) Null Under a Fixed MA(1) Alternative

It is useful to define the matrix  $S(\cdot, \cdot) = R(\cdot)N(\cdot)$ , which is a  $T$  by  $T$  band matrix with elements  $s_{i,i} = 1$ ,  $s_{i,i-1} = -(\cdot + \cdot)$ ,  $s_{i,i-2} = \cdot$  and  $s_{i,j} = 0$  otherwise. Denote the columns of  $S$  by  $s_1(\cdot, \cdot)$ ,  $s_2(\cdot, \cdot)$ , ...,  $s_T(\cdot, \cdot)$ . If the columns of the  $T$  by  $T$  identity matrix are

defined as  $f_1$ ,  $f_2$ ,  $f_3$ , ...,  $f_T$ , then define the  $T$  by  $T$  matrix  $L_1$  with the first  $(T-1)$

columns being  $f_2$ ,  $f_3$ , ...,  $f_T$  and the last column being a column of zeros. It should be

noted that  $s_i(\cdot, \cdot) = r_i(\cdot) - r_{i+1}(\cdot) = a_i(\cdot) - a_{i+1}(\cdot)$  and  $r_i(\cdot) = f_i - f_{i+1}$ . Define the  $T$  by

$T$  matrix  $N_1(\cdot)$  with columns  $a_2$ ,  $a_3$ , ...,  $a_T$ , with the last being a column of zeros.

Using (9), for the tests of the AR(1) model,  $ESS_R$  is given by

$$ESS_R = \hat{e}'\hat{e} - \hat{e}'\hat{U}_1(\hat{U}_1'\hat{U}_1)^{-1}\hat{U}_1'\hat{e}.$$

As indicated in section 4.1,  $\cdot^* = \text{plim}_1 \hat{\cdot} = -(1 + \cdot^2)$ , where  $\cdot$  is the true value of the MA(1) parameter. Let  $E_1 Z$  denote expectation of  $Z$  evaluated when  $H_1$ , the regression model with MA(1) errors, is true.

Proposition 1:  $\text{plim}_1 T^{-1}\hat{e}'\hat{e} = {}^2s_1^3(\cdot, \cdot^*)'s_1^3(\cdot, \cdot^*)$ .

Proof: Given the non-stochastic nature of  $x_t$ , and using (A1) and  $\text{plim}(\hat{\cdot} - \cdot) = 0$  under



both  $H_0$  and  $H_1$ , gives

$$p \lim T^{-1} \hat{e}' \hat{e} = p \lim T^{-1} [U' R'(\hat{\cdot}) R(\hat{\cdot}) U].$$

Evaluation under  $H_1$  gives

$$\begin{aligned} p \lim_1 T^{-1} \hat{e}' \hat{e} &= p \lim_1 T^{-1} [e' N'(\cdot) R'(\cdot^*) R(\cdot^*) N(\cdot) e] \\ &= p \lim_1 T^{-1} [e' S'(\cdot, \cdot^*) S(\cdot, \cdot^*) e] \\ &= {}^2 \lim T^{-1} \text{tr}[S'(\cdot, \cdot^*) S(\cdot, \cdot^*)] \\ &= {}^2 \underset{\sim}{s}_1^3(\cdot, \cdot^*)' \underset{\sim}{s}_1^3(\cdot, \cdot^*) \end{aligned}$$

where Lemma 1 is used to obtain the last result.

QED

Proposition 2:  $p \lim_1 T^{-1} \hat{e}' \hat{U}_1 = 0$ .

Proof: Using (A1) and (A2), and by similar reasoning to Proposition 1, gives

$$p \lim T^{-1} \hat{e}' \hat{U}_1 = p \lim T^{-1} [U' R'(\hat{\cdot}) U_1].$$

Evaluation under  $H_1$  yields

$$\begin{aligned} p \lim_1 T^{-1} \hat{e}' \hat{U}_1 &= p \lim_1 T^{-1} [e' N'(\cdot) R'(\cdot^*) L_1 N(\cdot) e] \\ &= p \lim_1 T^{-1} [e' S'(\cdot, \cdot^*) N_1(\cdot) e] \\ &= {}^2 \lim T^{-1} \text{tr}[S'(\cdot, \cdot^*) N_1(\cdot)] \\ &= {}^2 \underset{\sim}{s}_1^3(\cdot, \cdot^*)' \underset{\sim}{a}_2^3(\cdot) \\ &= - {}^2 [(\cdot + \cdot^*) + {}^2 \cdot^*]. \end{aligned}$$

Since  $\cdot^* = - \cdot / (1 + {}^2)$ ,  $p \lim_1 T^{-1} \hat{e}' \hat{U}_1 = 0$ .

QED

Proposition 3:  $p \lim_1 T^{-1} \hat{U}_1' \hat{U}_1 = {}^2 \underset{\sim}{a}_2^3(\cdot) \underset{\sim}{a}_2^3(\cdot)$ .

Proof: Using (A2), and by similar reasoning to Proposition 1, gives

$$p \lim T^{-1} \hat{U}_1' \hat{U}_1 = p \lim T^{-1} U_1' U_1.$$

Evaluation under  $H_1$  gives

$$\begin{aligned}
 \text{plim}_1 T^{-1} \hat{U}_1' \hat{U}_1 &= \text{plim}_1 T^{-1} [e' N'(\cdot) L_1' L_1 N(\cdot) e] \\
 &= \text{plim}_1 T^{-1} [e' N_1'(\cdot) N_1(\cdot) e] \\
 &= {}^2 \lim T^{-1} \text{tr}[N_1'(\cdot) N_1(\cdot)] \\
 &= {}^2 a_2^3(\cdot)' a_2^3(\cdot).
 \end{aligned}$$

QED

Combining the results in Propositions 1-3 implies  $\text{plim}_1 \text{ESS}_R / T = {}^2 s_1^3(\cdot, \cdot)' s_1^3(\cdot, \cdot)$ .

The other terms that appear in the various tests can be calculated in the same way using the following results for  $i, j > 0$ :

$$\text{plim}_1 \hat{e}' \hat{U}_j / T = {}^2 s_1^k(\cdot, \cdot)' a_{j+1}^k(\cdot), \text{ where } k=j+2$$

$$\text{plim}_1 \hat{U}_i' \hat{U}_j / T = {}^2 a_{i+1}^k(\cdot)' a_{j+1}^k(\cdot), \text{ where } k=2+\max(i, j)$$

$$\text{plim}_1 \hat{e}' \tilde{e}_j / T = {}^2 s_1^k(\cdot, \cdot)' f_{j+1}^k, \text{ where } k=1+\max(2, j)$$

$$\text{plim}_1 \hat{U}_i' \tilde{e}_j / T = {}^2 a_{i+1}^k(\cdot)' f_{j+1}^k, \text{ where } k=1+\max(i+1, j)$$

$$\text{plim}_1 \hat{e}_i' \hat{e}_j / T = {}^2 s_{i+1}^k(\cdot, \cdot)' s_{j+1}^k(\cdot, \cdot), \text{ where } k=3+\max(i, j)$$

$$\text{plim}_1 \hat{U}_i' \hat{e}_j / T = {}^2 a_{i+1}^k(\cdot)' s_{j+1}^k(\cdot, \cdot), \text{ where } k=2+\max(i, j+1)$$

$$\text{plim}_1 \hat{e}_i' \tilde{e}_j / T = {}^2 s_1^k(\cdot, \cdot)' s_{i+1}^k(\cdot, \cdot), \text{ where } k=i+3.$$

## B. Tests of the MA(1) Null Under a Fixed AR(1) Alternative

It is useful to define the matrix  $Q(\cdot, \cdot) = N(\cdot)^{-1} R(\cdot)^{-1} = S(\cdot, \cdot)^{-1}$ , which is a  $T$  by  $T$  band

matrix with elements  $q_{i,i} = 1$ ,  $q_{i,i-1} = +$ ,  $q_{i,i-j} = (+) q_{i,i-j+1} - q_{i,i-j+2}$  ( $j > 1$ ) and

$q_{i,j} = 0$  otherwise, and the matrix  $NQ(\tilde{\cdot}, \cdot) = N(\tilde{\cdot})^{-1}Q(\tilde{\cdot}, \cdot)$ , which is also a T by T band matrix with elements  $nq_{i,i} = 1$ ,  $nq_{i,i-1} = \alpha + 2$ ,  $nq_{i,i-2} = (\alpha + 2)nq_{i,i-1} - (2\alpha^2 + \alpha^2)$ ,  $nq_{i,i-j} = (\alpha + 2)nq_{i,i-j+1} - (2\alpha^2 + \alpha^2)nq_{i,i-j+2} + \alpha^2 nq_{i,i-j+3}$  ( $j > 2$ ) and  $nq_{i,j} = 0$  otherwise. Denote the columns of  $Q(\tilde{\cdot}, \cdot)$  by  $\tilde{q}_1(\tilde{\cdot}, \cdot), \tilde{q}_2(\tilde{\cdot}, \cdot), \dots, \tilde{q}_T(\tilde{\cdot}, \cdot)$  and the columns of  $NQ(\tilde{\cdot}, \cdot)$  by  $\tilde{nq}_1(\tilde{\cdot}, \cdot), \tilde{nq}_2(\tilde{\cdot}, \cdot), \dots, \tilde{nq}_T(\tilde{\cdot}, \cdot)$ . Define a T by T matrix  $NNQ(\tilde{\cdot}, \cdot)$  with columns  $\tilde{nq}_2(\tilde{\cdot}, \cdot), \tilde{nq}_3(\tilde{\cdot}, \cdot), \dots, \tilde{nq}_T(\tilde{\cdot}, \cdot)$ , and the last is a column of zeros. Denote the columns of  $R^{-1}(\tilde{\cdot})$  by  $\tilde{rr}_1(\tilde{\cdot}), \tilde{rr}_2(\tilde{\cdot}), \dots, \tilde{rr}_T(\tilde{\cdot})$ .  $R^{-1}(\tilde{\cdot})$  is a T by T band matrix with typical element  $rr_{i,i-j} = \alpha^j$  ( $j=0,1,2,3,\dots$ ) and  $rr_{i,j} = 0$  otherwise. Define the T by T matrix  $NNQ(\tilde{\cdot}, \cdot) = N(\tilde{\cdot})^{-2}Q(\tilde{\cdot}, \cdot)$  which is a band matrix with typical elements  $nnq_{i,i} = 1$ ,  $nnq_{i,i-1} = \alpha + 3$ ,  $nnq_{i,i-2} = (\alpha + 3)nnq_{i,i-1} - (3\alpha^2 + 3\alpha^2)$ ,  $nnq_{i,i-3} = (\alpha + 3)nnq_{i,i-2} - (3\alpha^2 + 3\alpha^2)nnq_{i,i-1} + (3\alpha^2 + \alpha^3)$ ,  $nnq_{i,i-j} = (\alpha + 3)nnq_{i,i-j+1} - (3\alpha^2 + 3\alpha^2)nnq_{i,i-j+2} + (3\alpha^2 + \alpha^3)nnq_{i,i-j+3} - \alpha^3 nnq_{i,i-j+4}$  ( $j > 3$ ) and  $nnq_{i,j} = 0$  otherwise. Denote the columns of  $NNQ(\tilde{\cdot}, \cdot)$  by  $\tilde{nnq}_1(\tilde{\cdot}, \cdot), \tilde{nnq}_2(\tilde{\cdot}, \cdot), \dots, \tilde{nnq}_T(\tilde{\cdot}, \cdot)$ .

Using (12) for the tests of the MA(1) model,  $ESS_R$  is given by

$$ESS_R = \tilde{e}'\tilde{e} - \tilde{e}'\tilde{e}_1^+(\tilde{e}_1^+\tilde{e}_1^+)^{-1}\tilde{e}_1^+\tilde{e}.$$

As indicated in section 4.2,  $\tilde{\cdot}^* = \text{plim}_0 \tilde{\cdot}$  and  $\tilde{\cdot}^*$  is given by the solution to (13) that lies

in the interval  $(-1, 1)$ . Let  $E_0 Z$  denote the expectation of  $Z$  evaluated when  $H_0$ , the regression model with AR(1) errors, is true.

Proposition 4:  $\text{plim}_0 T^{-1} \tilde{e}' \tilde{e} = {}^2 \lim_{\tilde{\alpha}} q_1(\tilde{\alpha}, \tilde{\alpha})' q_1(\tilde{\alpha}, \tilde{\alpha})$ .

Proof: Given the non-stochastic nature of  $x_t$ , and using (A3) and  $\text{plim}(\tilde{\alpha} - \alpha) = 0$  under both  $H_0$  and  $H_1$ , gives

$$\text{plim} T^{-1} \tilde{e}' \tilde{e} = \text{plim} T^{-1} [U' N'(\tilde{\alpha})^{-1} N(\tilde{\alpha})^{-1} U].$$

Evaluation under  $H_0$  gives

$$\begin{aligned} \text{plim}_0 T^{-1} \tilde{e}' \tilde{e} &= \text{plim}_0 T^{-1} [e' R'(\tilde{\alpha})^{-1} N'(\tilde{\alpha})^{-1} N(\tilde{\alpha})^{-1} R(\tilde{\alpha})^{-1} e] \\ &= \text{plim}_0 T^{-1} [e' Q'(\tilde{\alpha}, \tilde{\alpha}) Q(\tilde{\alpha}, \tilde{\alpha}) e] \\ &= {}^2 \lim_{\tilde{\alpha}} q_1(\tilde{\alpha}, \tilde{\alpha})' q_1(\tilde{\alpha}, \tilde{\alpha}). \end{aligned}$$

QED

Proposition 5:  $\text{plim}_0 \tilde{e}' e_i^+ = {}^2 \lim_{\tilde{\alpha}} q_1(\tilde{\alpha}, \tilde{\alpha})' n q_2(\tilde{\alpha}, \tilde{\alpha})$ .

Proof: Using (A3) and (A4), and by similar reasoning to Proposition 4, gives

$$\text{plim} T^{-1} \tilde{e}' e_i^+ = \text{plim} T^{-1} [U_1' N'(\tilde{\alpha})^{-1} N(\tilde{\alpha})^{-2} U_1].$$

Evaluation under  $H_0$  gives

$$\begin{aligned} \text{plim}_0 T^{-1} \tilde{e}' e_i^+ &= \text{plim}_0 T^{-1} [e' R'(\tilde{\alpha})^{-1} N'(\tilde{\alpha})^{-1} N(\tilde{\alpha})^{-2} L_1 R(\tilde{\alpha})^{-1} e] \\ &= \text{plim}_0 T^{-1} [e' Q'(\tilde{\alpha}, \tilde{\alpha}) N Q_1(\tilde{\alpha}, \tilde{\alpha}) e] \\ &= {}^2 \lim_{\tilde{\alpha}} q_1(\tilde{\alpha}, \tilde{\alpha})' n q_2(\tilde{\alpha}, \tilde{\alpha}). \end{aligned}$$

QED

Proposition 6:  $\text{plim}_0 T^{-1} \tilde{e}_i^+ \tilde{e}_i^+ = {}^2 \lim_{\tilde{\alpha}} n q_2(\tilde{\alpha}, \tilde{\alpha})' n q_2(\tilde{\alpha}, \tilde{\alpha})$ .

Proof: Using (A4), and by similar reasoning to Proposition 4, gives

$$p\lim T^{-1}\tilde{e}_i^+\tilde{e}_i^+ = p\lim T^{-1}[U_i'N'(\tilde{\cdot})^{-2}N(\tilde{\cdot})^{-2}U_i].$$

Evaluation under  $H_0$  gives

$$\begin{aligned} p\lim_0 T^{-1}\tilde{e}_i^+\tilde{e}_i^+ &= p\lim_0 T^{-1}[e'R'(\cdot)L_1'N'(\tilde{\cdot})^{-2}N(\tilde{\cdot})^{-2}L_1R'(\cdot)e] \\ &= p\lim_0 T^{-1}[e'NQ_1'(\cdot^*, \cdot)NQ_1(\cdot^*, \cdot)e] \\ &= {}^2\lim_{\tilde{\cdot}} nq_2(\cdot^*, \cdot)'nq_2(\cdot^*, \cdot). \end{aligned}$$

QED

The other terms that appear in the various tests can be calculated in the same way using the following results for  $i, j > 0$ :

$$\begin{aligned} p\lim_0 \tilde{e}_i^+\hat{U}_j/T &= {}^2\lim_{\tilde{\cdot}} nq_{i+1}(\cdot^*, \cdot)'rr_{j+1}(\cdot) \\ p\lim_0 \tilde{e}_i^+\tilde{e}_j^+/T &= {}^2\lim_{\tilde{\cdot}} nq_{i+1}(\cdot^*, \cdot)'nq_{j+1}(\cdot^*, \cdot) \\ p\lim_0 \hat{U}_i'\hat{U}_j/T &= {}^2\lim_{\tilde{\cdot}} rr_{i+1}(\cdot)'rr_{j+1}(\cdot) \\ p\lim_0 \hat{U}_i'\tilde{e}_j/T &= {}^2\lim_{\tilde{\cdot}} rr_{i+1}(\cdot)'q_{j+1}(\cdot^*, \cdot) \\ p\lim_0 \tilde{e}_i^+\tilde{e}_j^{++}/T &= {}^2\lim_{\tilde{\cdot}} nq_{i+1}(\cdot^*, \cdot)'nnq_{j+1}(\cdot^*, \cdot) \\ p\lim_0 \tilde{e}_i^{++}\tilde{e}_j^{++}/T &= {}^2\lim_{\tilde{\cdot}} nnq_{i+1}(\cdot^*, \cdot)'nnq_{j+1}(\cdot^*, \cdot) \\ p\lim_0 \tilde{e}_i\tilde{e}_j/T &= {}^2\lim_{\tilde{\cdot}} q_i(\cdot^*, \cdot)'q_{j+1}(\cdot^*, \cdot). \end{aligned}$$

## ACKNOWLEDGEMENTS

The authors would like to thank John Geweke, Soren Johansen, Kimio Morimune, and especially a referee, and seminar participants at the London School of Economics, University of Cambridge, and Institute of Mathematical Statistics – University of

Copenhagen, for helpful comments and suggestions. The first author wishes to acknowledge the financial support of the Asset Management Service Industry Fund, Osaka School of International Public Policy, Osaka University, and the second author wishes to acknowledge the financial support of the Australian Research Council, Japan Society for the Promotion of Science, Osaka School of International Public Policy at Osaka University, and the Institute of Social and Economic Research at Osaka University.

## REFERENCES

- [1] Pesaran, M. H.; Pesaran, B. Working with Microfit: Interactive Econometric Analysis, Oxford University Press: Oxford, 1997.
- [2] Quantitative Micro Software EViews 4.0 Command and Programming Reference, Quantitative Micro Software: Irvine, CA, 2000.
- [3] Reinsel, G. Maximum likelihood estimation of stochastic linear difference equations with autoregressive moving average errors. *Econometrica*, 1979, 47, 129-151.
- [4] Hannan, E. J.; Kavalieris, L.; Mackisack, M. Recursive estimation of linear systems. *Biometrika*, 1986, 73, 119-133.
- [5] Godfrey, L. G. On the invariance of the Lagrange multiplier test with respect to certain changes in the alternative hypothesis. *Econometrica*, 1981, 49, 1443-1456.
- [6] Walker, A. M. Some consequences of superimposed error in time series analysis. *Biometrika*, 1960, 47, 33-43.
- [7] Pesaran, M. H. Formation of inflation expectations in manufacturing industries. *Economic Journal*, 1985, 95, 948-975.
- [8] Hansen, L. P.; Hodrick, R. J. Forward exchange rates as optimal predictors of future spot rates: an econometric analysis. *Journal of Political Economy*, 1980, 88, 829-853.
- [9] Broze, L.; Gourieroux, C.; Szafarz, A. Solutions of linear rational expectations models. *Econometric Theory*, 1985, 1, 341-368.
- [10] Evans, G.; Honkapohja, S. A complete characterization of ARMA solutions to linear rational expectations models. *Review of Economic Studies*, 1986, 53, 227-239.
- [11] Bergstrom, A. R. Continuous time stochastic models and issues of aggregation over time. *Handbook of Econometrics, Volume II*; Z. Griliches and M.D. Intriligator, Eds, North-Holland: Amsterdam, 1984, 1145-1212.
- [12] Chow, G. C. *Econometrics*, McGraw-Hill: Tokyo, 1983.

- [13] Rowley, J. C. R.; Wilton, D. A. Quarterly models of wage determination: some new efficient estimates. *American Economic Review*, 1973, 63, 380-383.
- [14] Kenward, L. R. Autocorrelation and dynamic methodology with an application to wage determination models. *Journal of Econometrics*, 1975, 3, 179-187.
- [15] Harvey, A. C.; Todd, P. H. J. Forecasting economic time series with structural and Box-Jenkins models: a case study. *Journal of Business and Economic Statistics*, 1983, 1, 299-307.
- [16] Wallis, K. F. Seasonal adjustment and relations between variables. *Journal of the American Statistical Association*, 1974, 67, 18-31.
- [17] Engle, R. F.; Granger, C. W. J. Co-integration and error correction: representation, estimation, and testing. *Econometrica*, 1987, 55, 251-276.
- [18] Nicholls, D. F.; Pagan, A. R.; Terrell, R. D. The estimation and use of models with moving average disturbance terms: a survey. *International Economic Review*, 1975, 16, 113-134.
- [19] Schwert, G. W. Effects of model specification on tests for unit roots in macroeconomic data. *Journal of Monetary Economics*, 1987, 20, 73-103.
- [20] Hendry, D. F.; Mizon, G. E. Serial correlation as a convenient simplification not a nuisance: a comment on a study of the demand for money by the Bank of England. *Economic Journal*, 1978, 88, 549-553.
- [21] Sargan, J. D. Some tests of dynamic specification for a single equation. *Econometrica*, 1980, 48, 879-897.
- [22] Granger, C. W. J. The typical spectral shape of an economic variable. *Econometrica*, 1966, 34, 150-161.
- [23] Griffiths, W. E.; Beesley, P. A. A. The small-sample properties of some preliminary test estimators in a linear model with autocorrelated errors. *Journal of Econometrics*, 1984, 25, 49-61.
- [24] Zellner, A.; Keuzenkamp, H.; McAleer, M. (Eds.) *Simplicity, Inference and Modeling: Keeping It Sophisticatedly Simple*, Cambridge University Press: Cambridge and New York, 2001.
- [25] Box, G. E. P.; Jenkins, G. M. *Time Series Analysis: Forecasting and Control*, Holden Day: San Francisco, 1976.
- [26] Hendry, D. F.; Trivedi, P. K. Maximum likelihood estimation of difference equations with moving average errors: a simulation study. *Review of Economic Studies*, 1972, 39,

117-145.

[27] Whittle, P. Hypothesis Testing in Time Series Analysis, Almquist and Wiksell: Uppsala, 1951.

[28] Walker, A. M. Some tests of separate families of hypotheses in time series analysis. *Biometrika*, 1967, 54, 39-68.

[29] Pagan, A. R.; Hall, A. D.; Trivedi, P. K. Assessing the variability of inflation. *Review of Economic Studies*, 1983, 50, 585-596.

[30] McAleer, M.; McKenzie, C. R.; Hall, A. D. Testing separate time series models. *Journal of Time Series Analysis*, 1988, 9, 169-189.

[31] Hall, A. D.; McAleer, M. A Monte Carlo study of some tests of model adequacy in time series analysis. *Journal of Business and Economic Statistics*, 1989, 7, 95-106.

[32] Godfrey, L. G.; Tremayne, A. R. Checks of model adequacy for univariate time series models and their application to econometric relationships. *Econometric Reviews*, 1988, 7, 1-42.

[33] Godfrey, L. G.; Tremayne, A. R. Misspecification tests for univariate time series models and their application to econometric relationships. *The Implementation and Constructive Use of Misspecification Tests in Econometrics*; Godfrey, L.G., Ed.; Manchester University Press: Manchester, 1992, 1-66.

[34] Franses, P. H. A model selection test for an AR(1) versus an MA(1) model. *Statistics and Probability Letters*, 1992, 15, 281-284.

[35] Gouriéroux, C.; Monfort, A. Testing non-nested hypotheses. *Handbook of Econometrics*, Volume IV; R. F. Engle and D. L. McFadden, Eds; Elsevier: Amsterdam, 1994, 2585-2637.

[36] King, M. L.; McAleer, M. Further results on testing AR(1) against MA(1) disturbances in the linear regression model. *Review of Economic Studies*, 1987, 54, 649-663.

[37] Burke, S. P.; Godfrey, L. G.; Tremayne, A. R. Testing AR(1) against MA(1) disturbances in the linear regression model: an alternative procedure. *Review of Economic Studies*, 1990, 57, 135-145.

[38] Silvapulle, P.; King, M. L. Testing moving average against autoregressive disturbances in the linear regression model. *Journal of Business and Economic Statistics*, 1991, 9, 329-335.

[39] Silvapulle, P.; King, M. L. Non-nested testing for autocorrelation in the linear regression model. *Journal of Econometrics*, 1993, 58, 293-314.



- [40] McKenzie, C. R.; McAleer, M.; Gill, L. Simple procedures for testing autoregressive versus moving average errors in regression models. *Japanese Economic Review*, 1999, 50, 239-252.
- [41] Bahadur, R. R. Stochastic comparison of tests. *Annals of Mathematical Statistics*, 1960, 31, 276-295.
- [42] Bahadur, R. R. Rates of convergence of estimates and test statistics. *Annals of Mathematical Statistics*, 1967, 38, 303-324.
- [43] Geweke, J. The approximate slope of econometric tests. *Econometrica*, 1981a, 49, 1427-1442.
- [44] Geweke, J. A comparison of tests of independence of two covariance-stationary time series. *Journal of the American Statistical Association*, 1981b, 76, 363-373.
- [45] Geweke, J. Inference and causality in economic time series models. *Handbook of Econometrics*, Volume II; Z. Griliches and M. D. Intriligator, Eds; North-Holland: Amsterdam, 1984, 1101-1144.
- [46] Pesaran, M. H. Asymptotic power comparisons of tests of separate parametric families by Bahadur's approach. *Biometrika*, 1984, 71, 245-252.
- [47] Wascher, W. L. Assessing the relative efficiency of Hausman's test under Bahadur efficiency considerations. *Economics Letters*, 1986, 21, 343-347.
- [48] Pesaran, M. H.; Smith, R. J. A unified approach to estimation and orthogonality testing in linear single-equation econometric models. *Journal of Econometrics*, 1990, 44, 41-66.
- [49] Zabel, J. E. A comparison of nonnested tests for misspecified models using the method of approximate slopes. *Journal of Econometrics*, 1993, 57, 205-232.
- [50] Bera, A. K.; Newbold, P. Checks of model adequacy for univariate time series models and their application to econometric relationships: comment. *Econometric Reviews*, 1988, 7, 43-48.
- [51] Hatanaka, M. *Econometric Methods*, Sobunsha: Tokyo, 1991, in Japanese.

Table 1: Added Variables for Testing the AR(p) Null Against Three Alternatives

Alternative Hypothesis	Added Variables	Degrees of Freedom	Test
MA(q)	$\tilde{u}_{t-1}, \dots, \tilde{u}_{t-q}$	q	E1
MA(q)	$\tilde{u}_{t-1}^*, \dots, \tilde{u}_{t-q}^*$	q	E2
MA(q)	$\hat{u}_{t-1}^-, \tilde{u}_t$	1	DOP1
MA(q)	$\hat{u}_{t-1}^*, \tilde{u}_t^*$	1	DOP2
AR(p+r)	$\hat{u}_{t-p-1}, \dots, \hat{u}_{t-p-r}$	r	LMA1
AR(p+r)	$\hat{u}_{t-p-1}^*, \dots, \hat{u}_{t-p-r}^*$	r	LMA2
ARMA(p,r)	$\hat{u}_{t-1}, \dots, \hat{u}_{t-r}$	r	LMM1
ARMA(p,r)	$\hat{u}_{t-1}^*, \dots, \hat{u}_{t-r}^*$	r	LMM2

Notes:

1. An asterisk is used to denote the transformation of the relevant variable by  $\hat{\cdot}(L)$ ; for example,  $\tilde{u}_t^* = \hat{\cdot}(L) \tilde{u}_t$ .
2. LMA1=LMM1 (see Godfrey [5]).
3. E denotes the prediction error test, and DOP denotes the difference of prediction errors test.

Table 2: Added Variables for Testing the MA(q) Null Against Three Alternatives

Alternative Hypothesis	Added Variables	Degrees of Freedom	Test
AR(p)	$\hat{u}_{t-1}, \dots, \hat{u}_{t-p}$	p	E1
AR(p)	$\hat{u}_{t-1}^+, \dots, \hat{u}_{t-p}^+$	p	E2
AR(p)	$\hat{u}_t^-, \tilde{u}_t$	1	DOP1
AR(p)	$\hat{u}_t^+, \tilde{u}_t^+$	1	DOP2
MA(q+r)	$\tilde{u}_{t-q-1}^+, \dots, \tilde{u}_{t-q-r}^+$	r	LMM1
MA(q+r)	$\tilde{u}_{t-q-1}^{++}, \dots, \tilde{u}_{t-q-r}^{++}$	r	LMM2
ARMA(r,q)	$\tilde{u}_{t-1}^+, \dots, \tilde{u}_{t-r}^+$	r	LMA1
ARMA(r,q)	$\tilde{u}_{t-q-1}^{++}, \dots, \tilde{u}_{t-r}^{++}$	r-q [r>q]	LMA2

Notes:

1. A plus sign is used to denote the transformation of the relevant variable by  $\tilde{\cdot}(L)^{-1}$ ; for

example,  $\hat{u}_t^+ = \tilde{\cdot}(L)^{-1} \hat{u}_t$ .

2. LMM1=LMA1 (see Godfrey [5]).

3. E denotes the prediction error test, and DOP denotes the difference of prediction errors test.

4. Following the convention for the other tests, the added variables for LMA2 should be

$\tilde{u}_{t-1}^{++}, \dots, \tilde{u}_{t-r}^{++}$ . Since  $\tilde{u}_t^+ = \tilde{u}_t^{++}$ , it follows that  $\tilde{u}_{t-1}^{++}, \dots, \tilde{u}_{t-q}^{++}$  are perfectly correlated with the existing regressors in (7) and so are excluded from the above Table.

Table 3: Auxiliary Regressions to Calculate Bahadur Approximate Slopes for Tests of the AR(1) Null Evaluated Under the MA(1) Alternative

Test	Unrestricted Regression
E1	$s_1$ on $a_2$ and $f_2$
E2	$s_1$ on $a_2$ and $r_2$
DOP1	$s_1$ on $a_2$ and $-(f_2 + a_2)$
DOP2	$s_1$ on $a_2$ and $-(r_2 + (a_2 - a_3))$
LMA1	$s_1$ on $a_2$ and $a_3$
LMA2	$s_1$ on $a_2$ and $a_3 - a_4$
LMM1	$s_1$ on $a_2$ and $s_2$
LMM2	$s_1$ on $a_2$ and $s_2 - s_3$

where  $s_1 = s_1^5(\tilde{\cdot}, \cdot^*) = (1, -( \tilde{\cdot} + \cdot^*), \cdot^*, 0, 0)'$ ,  $s_2 = s_2^5(\tilde{\cdot}, \cdot^*) = (0, 1, -( \tilde{\cdot} + \cdot^*), \cdot^*, 0)'$ ,

$s_3 = s_3^5(\tilde{\cdot}, \cdot^*) = (0, 0, 1, -( \tilde{\cdot} + \cdot^*), \cdot^*)'$ ,  $a_2 = a_2^5(\tilde{\cdot}) = (0, 1, -\tilde{\cdot}, 0, 0)'$ ,  $a_3 = a_3^5(\tilde{\cdot}) = (0, 0, 1, -\tilde{\cdot}, 0)'$ ,

$a_4 = a_4^5(\tilde{\cdot}) = (0, 0, 0, 1, -\tilde{\cdot})'$ ,  $f_2 = f_2^5(\tilde{\cdot}) = (0, 1, 0, 0, 0)'$ , and  $r_2 = r_2^5(\tilde{\cdot}) = (0, 1, -\tilde{\cdot}, 0, 0)'$ . The vectors


$s_i^5(\tilde{\cdot}, \cdot^*)$  [ $i=1,2,3$ ],  $a_j^5(\tilde{\cdot})$  [ $j=2,3,4$ ],  $f_2^5$  and  $r_2^5(\tilde{\cdot})$  are defined in the Appendix.

Table 4: Auxiliary Regressions to Calculate Bahadur Approximate Slopes for Tests of the MA(1) Null Evaluated Under the AR(1) Alternative


Test	Unrestricted Regression
E1	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $rr_2(\cdot)_{\sim}$
E2	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $q_2(\cdot, \cdot)_{\sim}$
DOP1	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $-(\cdot)_{\sim}^* q_2(\cdot, \cdot)_{\sim} + rr_2(\cdot)_{\sim}$
DOP2	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $-(\cdot)_{\sim}^* nq_2(\cdot, \cdot)_{\sim} + q_2(\cdot, \cdot)_{\sim}$
LMM1	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $nq_3(\cdot, \cdot)_{\sim}$
LMM2	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $nnq_3(\cdot, \cdot)_{\sim}$
LMA1	$q_1(\cdot, \cdot)_{\sim}$ on $nq_2(\cdot, \cdot)_{\sim}$ and $q_2(\cdot, \cdot)_{\sim}$
LMA2	Not defined due to perfect collinearity among regressors.

where  $q_i(\cdot, \cdot)_{\sim}$ ,  $nq_i(\cdot, \cdot)_{\sim}$ ,  $nnq_i(\cdot, \cdot)_{\sim}$  and  $rr_i(\cdot)_{\sim}$  are defined in the Appendix.


**FIGURE 1: APPROXIMATE SLOPES (AS) OF METHOD 1 TESTS WITH AR(1) NULL**



**FIGURE 2: APPROXIMATE SLOPES (AS) OF METHOD 2 TESTS WITH AR(1) NULL**



**FIGURE 3: APPROXIMATE SLOPES (AS) OF METHOD 1 TESTS WITH  
MA(1) NULL**





**FIGURE 4: APPROXIMATE SLOPES (AS) OF METHOD 2 TESTS WITH  
MA(1) NULL**

