

Berti, Patrizia; Pratelli, Luca; Rigo, Pietro

Working Paper

Atomic Intersection of s-Fields and Some of Its Consequences

Quaderni di Dipartimento, No. 096

Provided in Cooperation with:

University of Pavia, Department of Economics and Quantitative Methods (EPMQ)

Suggested Citation: Berti, Patrizia; Pratelli, Luca; Rigo, Pietro (2009) : Atomic Intersection of s-Fields and Some of Its Consequences, Quaderni di Dipartimento, No. 096, Università degli Studi di Pavia, Dipartimento di Economia Politica e Metodi Quantitativi (EPMQ), Pavia

This Version is available at:

<https://hdl.handle.net/10419/95337>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.


Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



Quaderni di Dipartimento

**Atomic Intersection of σ -Fields
and Some of Its Consequences**

Patrizia Berti
(Università di Modena e Reggio Emilia)

Luca Pratelli
(Accademia Navale di Livorno)

Pietro Rigo
(Università di Pavia)

96 (05-09)

Dipartimento di economia politica
e metodi quantitativi
Università degli studi di Pavia
Via San Felice, 5
I-27100 Pavia

Maggio 2009

ATOMIC INTERSECTION OF σ -FIELDS AND SOME OF ITS CONSEQUENCES

PATRIZIA BERTI, LUCA PRATELLI, AND PIETRO RIGO

ABSTRACT. Let (Ω, \mathcal{F}, P) be a probability space. For each $\mathcal{G} \subset \mathcal{F}$, define $\overline{\mathcal{G}}$ as the σ -field generated by \mathcal{G} and those sets $F \in \mathcal{F}$ satisfying $P(F) \in \{0, 1\}$. Conditions for P to be atomic on $\cap_{i=1}^k \overline{\mathcal{A}_i}$, with $\mathcal{A}_1, \dots, \mathcal{A}_k \subset \mathcal{F}$ sub- σ -fields, are given. Conditions for P to be 0-1-valued on $\cap_{i=1}^k \overline{\mathcal{A}_i}$ are given as well. These conditions are useful in various fields, including Gibbs sampling, iterated conditional expectations and the intersection property.

1. INTRODUCTION

Throughout, (Ω, \mathcal{F}, P) is a probability space, $\mathcal{A}_1, \dots, \mathcal{A}_k \subset \mathcal{F}$ sub- σ -fields, $k \geq 2$, and we let

$$\mathcal{N} = \{F \in \mathcal{F} : P(F) \in \{0, 1\}\} \quad \text{and} \quad \overline{\mathcal{G}} = \sigma(\mathcal{G} \cup \mathcal{N}) \quad \text{for any } \mathcal{G} \subset \mathcal{F}.$$

As discussed in Section 3, the sub- σ -field

$$\mathcal{D} = \cap_{i=1}^k \overline{\mathcal{A}_i}$$

plays a role in various subjects, including Gibbs sampling, iterated conditional expectations and the intersection property. In a previous paper, in a Gibbs sampling framework, we investigated when $\cap_{i=1}^k \overline{\mathcal{A}_i} = \cap_{i=1}^k \mathcal{A}_i$; see [3].

In this paper, instead, we focus on atomicity of P on \mathcal{D} . In fact, atomicity of $P|_{\mathcal{D}}$ (i.e., the restriction of P to \mathcal{D}) has implications in each of the subjects mentioned above. It turns out that $P|_{\mathcal{D}}$ is actually atomic under mild conditions.

An extreme form of atomicity for $P|_{\mathcal{D}}$ is $\mathcal{D} = \mathcal{N}$, that is, P 0-1-valued on \mathcal{D} . Indeed, $\mathcal{D} = \mathcal{N}$ is fundamental for Gibbs sampling and very useful for the intersection property; see [3], [7] and [10].

Our main results are in Sections 4 and 5. Section 4 gives general results on atomicity of $P|_{\mathcal{D}}$. It includes a characterization (Theorem 2), a criterion for identifying the atoms (Theorem 3) and a sufficient condition (Theorem 4). Section 5, motivated by Gibbs sampling applications, concerns the particular case

$$\mathcal{A}_i = \sigma(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_k)$$

where X_1, \dots, X_k are any random variables on (Ω, \mathcal{F}, P) . It contains working sufficient conditions for $\mathcal{D} = \mathcal{N}$ (Theorem 8) and for $P|_{\mathcal{D}}$ to be atomic (Theorem 10). Indeed, $P|_{\mathcal{D}}$ is atomic whenever the probability distribution of (X_1, \dots, X_k) is absolutely continuous with respect to a σ -finite product measure.

2000 *Mathematics Subject Classification.* 60A10, 60J22, 62B99.

Key words and phrases. Atomic probability measure – Gibbs sampling – Graphical models – Intersection property – Iterated conditional expectations.

Finally, it is worth noting that $\mathcal{D} = \mathcal{N}$ whenever \mathcal{A}_r is independent of \mathcal{A}_s for some r, s . Given $D \in \mathcal{D}$, in fact, one has $P(A_i \Delta D) = 0$ for some $A_i \in \mathcal{A}_i$, $i = 1, \dots, k$. Hence, $P(D) = P(A_r \cap A_s) = P(A_r)P(A_s) = P(D)^2$.

2. PRELIMINARIES

Let $(\mathcal{X}, \mathcal{E}, Q)$ be a probability space. A Q -atom is any set $H \in \mathcal{E}$ such that $Q(H) > 0$ and $Q(\cdot | H)$ is 0-1-valued. In general, there are three possible situations: (i) Q is *nonatomic*, i.e., there are no Q -atoms; (ii) Q is *atomic*, i.e., the Q -atoms form a (countable) partition of \mathcal{X} ; (iii) there is $K \in \mathcal{E}$, $0 < Q(K) < 1$, such that $Q(\cdot | K)$ is nonatomic and K^c is a (countable) disjoint union of Q -atoms.

Thus, $D \subset \Omega$ is an atom of $P|\mathcal{D}$ if and only if $D \in \mathcal{D}$, $P(D) > 0$ and $P(\cdot | D)$ is 0-1-valued on \mathcal{D} . In the sequel, when $P|\mathcal{D}$ is atomic, we also say that \mathcal{D} is *atomic under P* .

For later purposes, we also note that Q is nonatomic if and only if $(\mathcal{X}, \mathcal{E}, Q)$ supports a real random variable with uniform distribution on $(0, 1)$. In fact, if U is a uniform random variable on $(\mathcal{X}, \mathcal{E}, Q)$, then Q is nonatomic since $\sigma(U) \subset \mathcal{E}$ and $Q|\sigma(U)$ is nonatomic. Conversely, by Lyapunov's convexity theorem, if Q is nonatomic the range of Q is $[0, 1]$; see e.g. [8] or Theorem 5.1.6 of [4] for a proof (based on transfinite induction or Zorn's lemma, respectively). Since the range of Q is $[0, 1]$, a uniform random variable on $(\mathcal{X}, \mathcal{E}, Q)$ can be obtained by arguing as in the proof of Lemma 2 of [1]; see also Theorem 3.1 of [2].

We finally recall that, for any sub- σ -field $\mathcal{G} \subset \mathcal{F}$,

$$\overline{\mathcal{G}} = \{F \in \mathcal{F} : P(F \Delta G) = 0 \text{ for some } G \in \mathcal{G}\}.$$

A straightforward consequence is that a real $\overline{\mathcal{G}}$ -measurable function on Ω coincides a.s. with some \mathcal{G} -measurable function. Thus, if $U : \Omega \rightarrow \mathbb{R}$ is \mathcal{D} -measurable, then $U = U_i$ a.s. for some \mathcal{A}_i -measurable function $U_i : \Omega \rightarrow \mathbb{R}$, $i = 1, \dots, k$.

3. FIELDS WHERE \mathcal{D} APPEARS

We list some fields involving \mathcal{D} , by paying particular attention to the case where $P|\mathcal{D}$ is atomic. We stress by now that, for atomicity of $P|\mathcal{D}$ to be a real advantage, the atoms of $P|\mathcal{D}$ and their probabilities should be known.

Throughout, X_i is a random variable on (Ω, \mathcal{F}, P) with values in the measurable space $(\mathcal{X}_i, \mathcal{B}_i)$, $i = 1, \dots, k$, and

$$X_i^* = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_k).$$

3.1. Intersection property. Let X be a random variable on (Ω, \mathcal{F}, P) with values in the measurable space $(\mathcal{X}, \mathcal{E})$. The intersection property (IP) is

$$X \perp X_i^* | X_i \text{ for } i = 1, \dots, k \implies X \perp (X_1, \dots, X_k)$$

where the notation " $U \perp V | W$ " stands for " U conditionally independent of V given W ". It is well known that IP may fail. As a trivial example, take X not independent of X_1 and $X_i = X_1$ for all i .

IP is involved in a number of arguments. It appears, for instance, in graphical models, zero entries in contingency tables, causal inference and estimation in Markov processes; see [10] and references therein.

The connections between IP and \mathcal{D} are made clear by part (b) of the next (obvious) result. Part (a) is already known for $k = 2$ (see Proposition 2.2 of [10] and references therein) but we give a proof to make the paper self-contained.

Theorem 1. *Let $\mathcal{A}_i = \sigma(X_i)$ for all i . Then:*

- (a) $X \perp X_i^* \mid X_i$ for $i = 1, \dots, k \iff E(f(X) \mid X_1, \dots, X_k) = E(f(X) \mid \mathcal{D})$ a.s. for each real bounded measurable function f on $(\mathcal{X}, \mathcal{E})$;
- (b) $X \perp X_i^* \mid X_i$ for $i = 1, \dots, k$ and $X \perp \mathcal{D} \iff X \perp (X_1, \dots, X_k)$;
- (c) $X \perp \mathcal{D}$ if and only if

$$P(X \in A, X_1 \in B_1) = P(X \in A)P(X_1 \in B_1) \quad \text{whenever}$$

$A \in \mathcal{E}, B_1 \in \mathcal{B}_1$ and $P(\{X_1 \in B_1\} \Delta \{X_i \in B_i\}) = 0$ for some $B_i \in \mathcal{B}_i, i = 2, \dots, k$.

Proof. (a) Suppose $E(f(X) \mid X_1, \dots, X_k) = E(f(X) \mid \mathcal{D})$ a.s. for all bounded measurable f on $(\mathcal{X}, \mathcal{E})$. Given i , since $\mathcal{D} \subset \overline{\sigma(X_i)} \subset \overline{\sigma(X_1, \dots, X_k)}$, then

$$E(f(X) \mid X_i) = E(f(X) \mid \mathcal{D}) = E(f(X) \mid X_1, \dots, X_k) \quad \text{a.s.}$$

for all bounded measurable f , that is, $X \perp X_i^* \mid X_i$. Conversely, suppose $X \perp X_i^* \mid X_i$ for all i . Given a bounded measurable f ,

$$E(f(X) \mid X_1) = E(f(X) \mid X_1, \dots, X_k) = E(f(X) \mid X_i) \quad \text{a.s. for all } i.$$

Thus, $E(f(X) \mid X_1)$ is $\overline{\sigma(X_i)}$ -measurable for all i , i.e., it is \mathcal{D} -measurable. Therefore, $E(f(X) \mid X_1, \dots, X_k) = E(f(X) \mid X_1) = E(f(X) \mid \mathcal{D})$ a.s..

(b) " \Leftarrow " is obvious. As to " \Rightarrow ", it suffices noting that $E(f(X) \mid X_1, \dots, X_k) = E(f(X) \mid \mathcal{D}) = E f(X)$ a.s. for all bounded measurable f , where the first equality is by part (a) and the second is because $X \perp \mathcal{D}$.

(c) Just note that

$$\mathcal{D} = \{F \in \mathcal{F} : P(F \Delta \{X_i \in B_i\}) = 0 \text{ for some } B_i \in \mathcal{B}_i, i = 1, \dots, k\}.$$

□

Thus, $X \perp \mathcal{D}$ is a (natural) sufficient condition for IP. In a sense, it is necessary as well, since it is a consequence of $X \perp (X_1, \dots, X_k)$. Heuristically, $X \perp \mathcal{D}$ means that X is not affected by that part of information which is common to X_1, \dots, X_k .

To test whether $X \perp \mathcal{D}$, atomicity of \mathcal{D} under P can help. If $P|_{\mathcal{D}}$ is atomic, in fact, $X \perp \mathcal{D}$ reduces to

$$P(X \in A, D) = P(X \in A)P(D) \quad \text{for all } A \in \mathcal{E} \text{ and atoms } D \text{ of } P|_{\mathcal{D}}.$$

As shown in Theorem 4, for $P|_{\mathcal{D}}$ to be atomic, it is enough that the distribution of (X_1, \dots, X_k) is absolutely continuous with respect to a σ -finite product measure; see also Lemma 6.

A last note is that $X \perp \mathcal{D}$ is trivially true whenever $\mathcal{D} = \mathcal{N}$. In [10], a paper which inspired the present Subsection, $\mathcal{D} = \mathcal{N}$ was firstly viewed as a sufficient condition for IP. In [3], in a Gibbs sampling framework, $\mathcal{D} = \mathcal{N}$ was given a characterization and various sufficient conditions.

3.2. Iterated conditional expectations. Let X be a real random variable on (Ω, \mathcal{F}, P) such that $EX^2 < \infty$ and

$$\mathcal{D}_{mk+i} = \mathcal{A}_i \quad \text{for all } m = 0, 1, \dots \text{ and } i = 1, \dots, k.$$

Define $Z_0 = X$ and $Z_n = E(Z_{n-1} \mid \mathcal{D}_n)$ for $n \geq 1$. Then,

$$(1) \quad Z_n \xrightarrow{\text{a.s.}} E(X \mid \mathcal{D}) \quad \text{as } n \rightarrow \infty.$$

This classical result was obtained by Burkholder-Chow [5] for $k = 2$ and Delyon-Delyon [6] for arbitrary k . See [7] for some historical notes.

Suppose now that \mathcal{D} is atomic under P and the goal is estimating EX . Then, $E(X | \mathcal{D}) = \sum_j I_{D_j} E(X | D_j)$ a.s. where D_1, D_2, \dots denote the (disjoint) atoms of $P|_{\mathcal{D}}$. Thus, one should apply relation (1) on each atom D_j , so as to obtain an estimate for $E(X | D_j)$, and then use the formula $EX = \sum_j P(D_j)E(X | D_j)$.

3.3. Gibbs sampling. As noted in [7], the limit theorem of Burkholder-Chow and Delyon-Delyon (Subsection 3.2) is intrinsically connected to Gibbs sampling.

Let X_1, \dots, X_k be the canonical projections on

$$(\Omega, \mathcal{F}) = \left(\prod_{i=1}^k \mathcal{X}_i, \prod_{i=1}^k \mathcal{B}_i \right).$$

Each X_i is assumed to admit a regular conditional distribution γ_i given X_i^* . In the notation $u = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k)$, where $x_j \in \mathcal{X}_j$ for all $j \neq i$, this means that: (i) $\gamma_i(u)$ is a probability measure on \mathcal{B}_i for every u ; (ii) $u \mapsto \gamma_i(u)(A)$ is measurable for $A \in \mathcal{B}_i$; (iii) $P(B) = \int \int I_B(x_1, \dots, x_k) \gamma_i(u) dx_i \gamma_i^*(du)$ for $B \in \prod_{i=1}^k \mathcal{B}_i$, where γ_i^* denotes the marginal distribution of X_i^* .

The Gibbs-chain

$$Y_n = (Y_{1,n}, \dots, Y_{k,n}), \quad n \geq 0,$$

can be informally described as follows. Starting from $\omega = (x_1, \dots, x_k)$, the next state $\omega^* = (a_1, \dots, a_k)$ is obtained by sequentially generating a_k, a_{k-1}, \dots, a_1 , each a_i being selected from the conditional distribution of X_i given $X_1 = x_1, \dots, X_{i-1} = x_{i-1}, X_{i+1} = a_{i+1}, \dots, X_k = a_k$. Formally, (Y_n) is the homogeneous Markov chain with state space (Ω, \mathcal{F}) and transition kernel

$$\begin{aligned} K(\omega, B) &= K((x_1, \dots, x_k), B) \\ &= \int \dots \int I_B(a_1, \dots, a_k) \prod_{i=1}^k \gamma_i(x_1, \dots, x_{i-1}, a_{i+1}, \dots, a_k) (da_i). \end{aligned}$$

Note that P is a stationary distribution for (Y_n) , i.e., $P(\cdot) = \int K(\omega, \cdot) P(d\omega)$. Let \mathbb{P} denote the law of (Y_n) such that $Y_0 \sim P$.

The Gibbs chain is constructed mainly for sampling from P . To this end, the following SLLN is fundamental

$$(2) \quad \frac{1}{n} \sum_{i=0}^{n-1} \phi(Y_i) \rightarrow \int \phi dP, \quad \mathbb{P}\text{-a.s.}, \quad \text{for all } \phi \in L_1(P).$$

Another basic requirement of (Y_n) , stronger than (2), is ergodicity on some set $S \in \mathcal{F}$, that is

$$P(S) = 1, \quad K(\omega, S) = 1 \quad \text{and} \quad \|K^n(\omega, \cdot) - P\| \rightarrow 0 \quad \text{for each } \omega \in S,$$

where $\|\cdot\|$ is total variation norm and K^n the n -th iterate of K .

Now, letting $\mathcal{A}_i = \sigma(X_i^*)$ for all i , the SLLN under (2) is equivalent to

$$\mathcal{D} = \mathcal{N}.$$

In addition, in case \mathcal{F} is countably generated and P absolutely continuous with respect to a σ -finite product measure, $\mathcal{D} = \mathcal{N}$ if and only if (Y_n) is ergodic on $S_0 = \{\omega \in \Omega : K(\omega, \cdot) \ll P\}$. See Theorems 4.2, 4.5 and Remark 4.7 of [3].

Conditions for $\mathcal{D} = \mathcal{N}$ (when $\mathcal{A}_i = \sigma(X_i^*)$ for all i) are given in Theorem 8; see also Lemma 6.

Strictly speaking, thus, Gibbs sampling is admissible only if $\mathcal{D} = \mathcal{N}$. At least in principle, however, it makes sense even if $\mathcal{D} \neq \mathcal{N}$, provided \mathcal{D} is atomic under P . In fact, if $P|\mathcal{D}$ is atomic (with disjoint atoms D_1, D_2, \dots), then (2) turns into

$$(2^*) \quad \frac{1}{n} \sum_{i=0}^{n-1} \phi(Y_i) \rightarrow \int \phi(\omega) P(d\omega | D_j),$$

\mathbb{P} -a.s. on $\{Y_0 \in D_j\}$, for all j and $\phi \in L_1(P)$.

The SLLN under (2*) can be proved by the same argument used for Theorem 4.2 of [3] plus the observation that $K(\cdot, B) = I_B(\cdot)$, P -a.s., for each $B \in \mathcal{D}$.

If the atoms D_j are only a finite number, and they are known together with their probabilities $P(D_j)$, then (2*) can be used to evaluate $\int \phi dP$. The chain (Y_n) should be started on each D_j , so as to obtain an estimate for $\int \phi(\omega) P(d\omega | D_j)$, and then the formula $\int \phi dP = \sum_j P(D_j) \int \phi(\omega) P(d\omega | D_j)$ should be applied.

As shown in Theorem 10, for $P|\mathcal{D}$ to be atomic, it is enough that the distribution of (X_1, \dots, X_k) is absolutely continuous with respect to a σ -finite product measure.

4. ATOMICITY OF \mathcal{D} UNDER P

We begin with a definition. Say that $H \subset \Omega$ has the *trivial intersection property*, or briefly that H is TIP, in case $H \in \mathcal{F}$, $P(H) > 0$, and

$$A_i \in \mathcal{A}_i \text{ and } P(A_i \Delta A_1 | H) = 0 \text{ for } i = 1, \dots, k \implies P(A_1 | H) \in \{0, 1\}.$$

Here are some obvious consequences of the definition.

(i) If H is TIP, $A_i \in \mathcal{A}_i$ and $P(A_i \Delta A_1 | H) = 0$ for all i , then either $P(A_i | H) = 0$ for all i or $P(A_i | H) = 1$ for all i .

(ii) Let $H \in \mathcal{F}$ with $P(H) > 0$ and write $P_H = P(\cdot | H)$. Then, H is TIP if and only if Ω is P_H -TIP (i.e., Ω is TIP under P_H). Moreover, Ω is TIP if and only if $\mathcal{D} = \mathcal{N}$. Therefore, the definition of TIP set may be rephrased as follows: H is TIP if and only if

$$\mathcal{D}_{P_H} = \mathcal{N}_{P_H}$$

where $\mathcal{N}_{P_H} = \{F \in \mathcal{F} : P_H(F) \in \{0, 1\}\}$ and $\mathcal{D}_{P_H} = \bigcap_{i=1}^k \sigma(\mathcal{A}_i \cup \mathcal{N}_{P_H})$.

(iii) Let Q be a probability on \mathcal{F} . If P and Q are equivalent (i.e., $P \ll Q$ and $Q \ll P$), then H is Q -TIP if and only if it is P -TIP. If $P \ll Q$ and $H \subset \{\frac{dP}{dQ} > 0\}$, for some given version of $\frac{dP}{dQ}$, then H is Q -TIP if and only if it is P -TIP.

The present notion of TIP set generalizes the one given in [3] for $k = 2$. Among other things, such a notion is basic for characterizing atomicity of $P|\mathcal{D}$.

Theorem 2. *Let $H \subset \Omega$. Then,*

- (a) *If H is TIP, there is an atom H^* of $P|\mathcal{D}$ satisfying $H^* \supset H$ and $P(D | H^*) = P(D | H)$ for all $D \in \mathcal{D}$;*
- (b) *For H to be an atom of $P|\mathcal{D}$ it is necessary and sufficient that $H \in \mathcal{D}$ and H is TIP;*
- (c) *\mathcal{D} is atomic under P if and only if $P(\cup_n H_n) = 1$ for some countable collection H_1, H_2, \dots of TIP sets.*

Proof. (a) Suppose H is TIP. We first prove that $P(\cdot | H)$ is 0-1 on \mathcal{D} . Given $D \in \mathcal{D}$, for each i there is $A_i \in \mathcal{A}_i$ such that $P(A_i \Delta D) = 0$. Hence, $P((A_i \Delta A_1) \cap H) \leq P(A_i \Delta A_1) = 0$ for all i , and H TIP implies $P(D | H) = P(A_1 | H) \in \{0, 1\}$. Next, by a standard argument, there is $H^* \in \mathcal{D}$ such that $H^* \supset H$ and

$$P(H^*) = \inf\{P(D) : H \subset D \in \mathcal{D}\}.$$

Let $D \in \mathcal{D}$. If $P(D | H) = 1$, then

$$H \subset (D \cap H^*) \cup (D^c \cap H) \in \mathcal{D},$$

so that $P(H^*) \leq P((D \cap H^*) \cup (D^c \cap H)) = P(D \cap H^*)$ by definition of H^* . Hence, $P(D | H^*) = 1$. Taking complements, if $P(D | H) = 0$ then $P(D | H^*) = 0$. Thus, H^* is an atom of $P|\mathcal{D}$ and $P(\cdot | H^*) = P(\cdot | H)$ on \mathcal{D} .

(b) If $H \in \mathcal{D}$ is TIP, then H is an atom of $P|\mathcal{D}$ since $P(H | H^*) = P(H | H) = 1$, where H^* is as in point (a). Conversely, suppose H is an atom of $P|\mathcal{D}$. To prove H TIP, we fix $A_i \in \mathcal{A}_i$ such that $P(A_i \Delta A_1 | H) = 0$ for $i = 1, \dots, k$. For each i , since $H \in \mathcal{D} \subset \overline{\mathcal{A}_i}$, some $H_i \in \mathcal{A}_i$ meets $P(H \Delta H_i) = 0$. Moreover,

$$P((A_1 \cap H) \Delta (A_i \cap H_i)) \leq P(H \Delta H_i) + P((A_i \Delta A_1) \cap H) = 0.$$

Hence, $A_1 \cap H \in \overline{\mathcal{A}_i}$ for all i , that is, $A_1 \cap H \in \mathcal{D}$. Since H is an atom of $P|\mathcal{D}$, it follows that $P(A_1 | H) = P(A_1 \cap H | H) \in \{0, 1\}$. Thus, H is TIP.

(c) If $P|\mathcal{D}$ is atomic, it suffices to take the H_n as the atoms of $P|\mathcal{D}$ and to apply point (b). Conversely, if $P(\cup_n H_n) = 1$ with the H_n TIP, for each n point (a) implies $H_n \subset H_n^*$ for some atom H_n^* of $P|\mathcal{D}$. Then, $P|\mathcal{D}$ is atomic since $P(\cup_n H_n^*) \geq P(\cup_n H_n) = 1$. \square

By Theorem 2, $P|\mathcal{D}$ is atomic provided Ω can be covered by countably many TIP sets H_1, H_2, \dots . In this case, every atom D admits the representation $D = \cup_{i \in I} H_i$ a.s. for some index set I (by point (a)). The next issue, thus, is identifying such atoms using the H_n as building blocks. Indeed, the atoms are maximal TIP sets, according to the following result.

Theorem 3. *Suppose $P(\cup_n H_n) = 1$, where H_1, H_2, \dots are TIP, and let $D \subset \Omega$. Then, D is an atom of $P|\mathcal{D}$ if and only if D is TIP and*

$$(3) \quad D \cup H_n \text{ fails to be TIP whenever } P(H_n \setminus D) > 0.$$

Proof. By Theorem 2, it can be assumed D TIP, and we have to prove that condition (3) is equivalent to $D \in \mathcal{D}$. Suppose (3) holds. Let $N = \{n : P(H_n \setminus D) > 0\}$. If $N = \emptyset$, then $P(D^c) \leq \sum_n P(H_n \setminus D) = 0$, so that $D \in \mathcal{N} \subset \mathcal{D}$. If $N \neq \emptyset$, by (3), for each $n \in N$ there are $A_{i,n} \in \mathcal{A}_i$, $i = 1, \dots, k$, satisfying

$$P(A_{i,n} \Delta A_{1,n} | D \cup H_n) = 0 \text{ and } P(A_{i,n} | D \cup H_n) \in (0, 1) \text{ for all } i.$$

Since D and H_n are TIP, one also has $P(A_{i,n} | D) \in \{0, 1\}$ and $P(A_{i,n} | H_n) \in \{0, 1\}$ for all i , and thus

$$P(A_{i,n} | D) = 1 - P(A_{i,n} | H_n) \text{ for all } i.$$

Define $F_{i,n} = A_{i,n}$ or $F_{i,n} = A_{i,n}^c$ as $P(A_{i,n} | D) = 1$ or $P(A_{i,n} | D) = 0$, and

$$A_i = \cap_{n \in N} F_{i,n}.$$

Then, $P(A_i | D) = 1$ and $P(A_i | H_n) = 0$ for all i and $n \in N$. Hence, given i ,

$$P(A_i \Delta D) = P(A_i \setminus D) \leq \sum_{n \in N} P(A_i \cap D^c \cap H_n) \leq \sum_{n \in N} P(A_i \cap H_n) = 0.$$

Since $A_i \in \mathcal{A}_i$, it follows that $D \in \overline{\mathcal{A}_i}$, that is, $D \in \mathcal{D}$. Conversely, suppose $D \in \mathcal{D}$ and $D \cup H_n$ is TIP for some n . Since $P(D \mid D \cup H_n) \in \{0, 1\}$ (by point (a) of Theorem 2) and $P(D) > 0$ (as D is TIP), then $P(H_n \setminus D) = 0$. \square

In real problems, it is not unusual that $P \ll Q$ for some probability Q on \mathcal{F} which makes $\mathcal{A}_1, \dots, \mathcal{A}_k$ independent. This does not imply $\mathcal{D} = \mathcal{N}$ (see Examples 3.16 and 3.17 of [3]) but it suffices for atomicity of $P \mid \mathcal{D}$. Actually, it is enough that a couple of the \mathcal{A}_i are independent under Q .

Theorem 4. *$P \mid \mathcal{D}$ is atomic provided $P \ll Q$ for some probability measure Q on \mathcal{F} which makes \mathcal{A}_r and \mathcal{A}_s independent for some r, s .*

Proof. Fix $H \in \mathcal{D}$ with $P(H) > 0$ and let $P_H = P(\cdot \mid H)$. If $P_H \mid \mathcal{D}$ is nonatomic, the probability space $(\Omega, \mathcal{D}, P_H \mid \mathcal{D})$ supports a real random variable with uniform distribution; see Section 2. Hence, it suffices to prove that each \mathcal{D} -measurable function $U : \Omega \rightarrow \mathbb{R}$ satisfies $P_H(U \in C) = 1$ for some countable set $C \subset \mathbb{R}$. Further, since $P_H \ll P$, it is enough to show that $P(U \in C) = 1$. Let $U : \Omega \rightarrow \mathbb{R}$ be \mathcal{D} -measurable. Then, $U = U_i$ a.s. for some $U_i : \Omega \rightarrow \mathbb{R}$ satisfying $\sigma(U_i) \subset \mathcal{A}_i$, $i = 1, \dots, k$. Define the countable set $C = \{c \in \mathbb{R} : Q(U_s = c) > 0\}$. Since U_r and U_s are independent under Q ,

$$Q(U_r \notin C, U_r = U_s) = \int_{\{U_r \notin C\}} Q\{x : U_s(x) = U_r(\omega)\} Q(d\omega) = 0.$$

Thus, $P \ll Q$ yields

$$P(U \in C) = 1 - P(U \notin C, U = U_r = U_s) = 1 - P(U_r \notin C, U_r = U_s) = 1. \quad \square$$

For $k = 2$, Theorem 4 reduces to Theorem 3.10 of [3].

Remark 5. Let $\mathcal{A}_i = \sigma(X_i)$ for all i , where $X_i : \Omega \rightarrow \mathcal{X}_i$ is a random variable and \mathcal{X}_i a separable metric space (equipped with its Borel σ -field \mathcal{B}_i). Then, $P \mid \mathcal{D}$ need not be atomic even though

$$(4) \quad P(X_i = f(X_j)) = 0$$

for all $i \neq j$ and all measurable functions $f : \mathcal{X}_j \rightarrow \mathcal{X}_i$. We mention this fact since, for some time, we conjectured $P \mid \mathcal{D}$ atomic under (4).

As an example, let $k = 2$, $X_1 = (U, W)$ and $X_2 = (V, W)$, where U, V, W are real independent random variables with nonatomic distributions. Take $\mathcal{X}_i = \mathbb{R}^2$ and $\mathcal{A}_i = \sigma(X_i)$ for $i = 1, 2$. Given a measurable function $f = (f_1, f_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, one obtains $P(X_2 = f(X_1)) \leq P(V = f_1(U, W)) = 0$ since V has nonatomic distribution and is independent of (U, W) . Likewise, $P(X_1 = f(X_2)) = 0$. However, $P \mid \mathcal{D}$ is nonatomic, as $\sigma(W) \subset \mathcal{D}$ and W has nonatomic distribution.

Finally, we state a simple but useful fact as a lemma. Let Q be a probability measure on \mathcal{F} . Say that P and Q are *equivalent on rectangles* in case

$$P(A) = 0 \Leftrightarrow Q(A) = 0 \text{ for each } A \in \mathcal{R},$$

$$\text{where } \mathcal{R} = \{\cap_{i=1}^k A_i : A_i \in \mathcal{A}_i, i = 1, \dots, k\}.$$

If $A \in \mathcal{R}$, then A^c is a finite union of elements of \mathcal{R} . Hence, $P(A) = 1 \Leftrightarrow Q(A) = 1$ and $P(A \Delta B) = 0 \Leftrightarrow Q(A \Delta B) = 0$ whenever $A, B \in \mathcal{R}$ and P, Q are equivalent on rectangles. Note that $\mathcal{A}_i \subset \mathcal{R}$ for all i . Note also that P and Q need not be equivalent on $\sigma(\mathcal{R})$ even though they are equivalent on rectangles.

Lemma 6. *Let P and Q be equivalent on rectangles. If D is an atom of $Q|\mathcal{D}_Q$, there is $A \in \mathcal{R}$ such that $Q(A\Delta D) = 0$ and A is an atom of $P|\mathcal{D}_P$. Moreover,*

$$\mathcal{D}_Q = \mathcal{N}_Q \Leftrightarrow \mathcal{D}_P = \mathcal{N}_P, \text{ and}$$

$$\mathcal{D}_Q \text{ is atomic under } Q \Leftrightarrow \mathcal{D}_P \text{ is atomic under } P.$$

(Here, $\mathcal{N}_Q = \{F \in \mathcal{F} : Q(F) \in \{0, 1\}\}$, $\mathcal{D}_Q = \cap_{i=1}^k \sigma(\mathcal{A}_i \cup \mathcal{N}_Q)$, $\mathcal{N}_P = \mathcal{N}$ and $\mathcal{D}_P = \mathcal{D}$).

Proof. We first prove that, for each $D \in \mathcal{D}_Q$ with $Q(D) > 0$, there is $A = A(D)$ satisfying $A \in \mathcal{A}_1 \cap \mathcal{D}_P$, $Q(A\Delta D) = 0$ and $P(A) > 0$. Take in fact $A_i \in \mathcal{A}_i$ with $Q(A_i\Delta D) = 0$, $i = 1, \dots, k$, and let $A = A_1$. Then $A \in \mathcal{A}_1$, $Q(A\Delta D) = 0$ and $P(A) > 0$. Since $Q(A_i\Delta A) = 0$ for all i , then $P(A_i\Delta A) = 0$ for all i , so that $A \in \mathcal{D}_P$. Next, let D be an atom of $Q|\mathcal{D}_Q$ and $A = A(D)$. Then, $A \in \mathcal{A}_1 \subset \mathcal{R}$. Given $G \in \mathcal{D}_P$, for each i there is $G_i \in \mathcal{A}_i$ such that $P(G\Delta G_i) = 0$. Again, $P(G_i\Delta G_1) = 0$ for all i implies $Q(G_i\Delta G_1) = 0$ for all i , so that $G_1 \in \mathcal{D}_Q$. Since A is an atom of $Q|\mathcal{D}_Q$ (as $Q(A\Delta D) = 0$), either $Q(A \cap G_1) = 0$ or $Q(A \cap G_1^c) = 0$. Accordingly, either $P(A \cap G) = P(A \cap G_1) = 0$ or $P(A \cap G^c) = P(A \cap G_1^c) = 0$, i.e., A is an atom of $P|\mathcal{D}_P$. Next, if $\mathcal{D}_Q = \mathcal{N}_Q$, then Ω is an atom of $Q|\mathcal{D}_Q$. Thus, $A = A(\Omega)$ is an atom of $P|\mathcal{D}_P$ and $P(A) = 1$, i.e., $\mathcal{D}_P = \mathcal{N}_P$. Finally, suppose $Q|\mathcal{D}_Q$ is atomic with (disjoint) atoms D_1, D_2, \dots . Let $A_j = A(D_j)$ and $A = \cup_j A_j$. Then, each A_j is an atom of $P|\mathcal{D}_P$, and $P(A) = 1$ since $Q(A) = 1$ and $A \in \mathcal{A}_1 \subset \mathcal{R}$. Therefore, $P|\mathcal{D}_P$ is atomic. \square

5. APPLICATIONS TO GIBBS SAMPLING

As remarked in Subsection 3.3, in a Gibbs sampling framework it is fundamental that $\mathcal{D} = \mathcal{N}$, or at least that \mathcal{D} is atomic under P , when the \mathcal{A}_i are given by

$$\mathcal{A}_i = \sigma(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_k).$$

In this section, we let $\mathcal{A}_i = \sigma(X_i^*)$ for all i , where $X_i^* = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_k)$ and the X_i are random variables on (Ω, \mathcal{F}, P) with values in the measurable spaces $(\mathcal{X}_i, \mathcal{B}_i)$, $i = 1, \dots, k$. We also let $\mathcal{D}_0 = \cap_i \overline{\sigma(X_i)}$. Since $\mathcal{D}_0 \subset \mathcal{D}$, P is 0-1-valued or atomic on \mathcal{D}_0 whenever it is so on \mathcal{D} .

Let $\mathcal{X} = \prod_{i=1}^k \mathcal{X}_i$ and let $\mathcal{B} = \prod_{i=1}^k \mathcal{B}_i$ denote the product σ -field on \mathcal{X} . Define two measures on \mathcal{B} as

$$\gamma(\cdot) = P((X_1, \dots, X_k) \in \cdot) \quad \text{and} \quad \mu = \mu_1 \times \dots \times \mu_k$$

where each μ_i is a σ -finite measure on \mathcal{B}_i . Thus, γ is the probability distribution of (X_1, \dots, X_k) and μ a σ -finite product measure.

By Theorem 4, it follows that $P|\mathcal{D}_0$ is atomic whenever $\gamma \ll \mu$. Whether or not $\gamma \ll \mu$ implies $P|\mathcal{D}$ atomic is a bit more delicate and is the main focus of this section. We start by noting that, in the independent case, things are as expected.

Lemma 7. *Let $\mathcal{A}_i = \sigma(X_i^*)$ for all i and*

$$\mathcal{D}_j = \cap_{i=1}^j \overline{\sigma(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_j)}, \quad j = 2, \dots, k.$$

If X_j is independent of (X_1, \dots, X_{j-1}) , then $\mathcal{D}_j = \mathcal{D}_{j-1}$. In particular, if X_1, \dots, X_k are independent, then $\mathcal{D} = \mathcal{N}$ and $H = \{X_1 \in B_1, \dots, X_k \in B_k\}$ is TIP as far as $B_i \in \mathcal{B}_i$ for all i and $P(H) > 0$.

Proof. Since $\mathcal{D}_{j-1} \subset \mathcal{D}_j$, it suffices to prove $\mathcal{D}_{j-1} \supset \mathcal{D}_j$. Let $A \in \mathcal{D}_j$. Then, $I_A = f_i(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_j)$ a.s. for some bounded measurable function f_i , $i = 1, \dots, j$. Let α_j denote the probability distribution of X_j . If X_j is independent of (X_1, \dots, X_{j-1}) , then

$$\begin{aligned} I_A &= E(I_A \mid X_1, \dots, X_{j-1}) \\ &= E(f_i(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_j) \mid X_1, \dots, X_{j-1}) \\ &= \int f_i(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_{j-1}, t) \alpha_j(dt) \quad \text{a.s. for each } i < j. \end{aligned}$$

Thus, $A \in \mathcal{D}_{j-1}$. Next, suppose X_1, \dots, X_k are independent. By what already proved,

$$\mathcal{D} = \mathcal{D}_k = \mathcal{D}_{k-1} = \dots = \mathcal{D}_2 = \overline{\sigma(X_1)} \cap \overline{\sigma(X_2)} = \mathcal{N},$$

or equivalently Ω is TIP. Since X_1, \dots, X_k are still independent under $P(\cdot \mid H)$, it follows that Ω is $P(\cdot \mid H)$ -TIP, that is, H is P -TIP. \square

The independence assumption can be considerably relaxed. Next result is inspired to Corollary 3.7 of [3].

Theorem 8. *Suppose $\mathcal{A}_i = \sigma(X_i^*)$ for all i , $\gamma \ll \mu$ and f is a version of $\frac{d\gamma}{d\mu}$. Let*

$$H = \{(X_1, \dots, X_k) \in B\} \quad \text{where } B \in \mathcal{B} \text{ and } B \subset \{f > 0\}.$$

Then, H is TIP provided

$$(5) \quad \cup_{i=1}^k \{X_i^* \in B_i^*\} \supset H \supset \cap_{i=1}^k \{X_i \in B_i\}, \quad \text{where } B_i^* = \times_{j \neq i} B_j, \\ \text{for some } B_i \in \mathcal{B}_i, i = 1, \dots, k, \text{ with } P(X_1 \in B_1, \dots, X_k \in B_k) > 0.$$

Proof. It can be assumed $(\Omega, \mathcal{F}, P) = (\mathcal{X}, \mathcal{B}, \gamma)$ and X_1, \dots, X_k the canonical projections (so that $H = B$). For each i , since μ_i is σ -finite, there is a probability Q_i on \mathcal{B}_i equivalent to μ_i . Let $Q = Q_1 \times \dots \times Q_k$ denote the corresponding product probability on \mathcal{B} . Since $P \ll Q$ and $P(H) > 0$, then $Q(H) > 0$. Since $f > 0$ on H , then $Q(\cdot \mid H)$ is equivalent to $P(\cdot \mid H)$. Thus, H is P -TIP if and only if it is Q -TIP. We next prove that H is Q -TIP. Let $K = \{X_1 \in B_1, \dots, X_k \in B_k\}$. Since $Q(K) > 0$ (due to $P(K) > 0$) and X_1, \dots, X_k are independent under Q , Lemma 7 implies that K is Q -TIP. Fix $A_i \in \mathcal{A}_i$ with $Q(A_i \Delta A_1 \mid H) = 0$, $i = 1, \dots, k$. Since K is Q -TIP and $K \subset H$, then $Q(A_1 \mid K) \in \{0, 1\}$, say $Q(A_1 \mid K) = 0$ (so that $Q(A_i \mid K) = 0$ for all i). Given j , since $Q(X_j \in B_j) > 0$ and

$$Q(A_j, X_j^* \in B_j^*) Q(X_j \in B_j) = Q(A_j, X_j^* \in B_j^*, X_j \in B_j) = Q(A_j \cap K) = 0,$$

then $Q(A_j, X_j^* \in B_j^*) = 0$. Also, $\{X_j^* \in B_j^*\} \cap \{X_r \notin B_r\} = \emptyset$ for $j \neq r$ and $H \subset \cup_{i=1}^k \{X_i^* \in B_i^*\}$. Thus, letting $A = \cap_i A_i$,

$$\begin{aligned} Q(A_1 \cap H) &= Q(A \cap H) = Q(A \cap H \cap K^c) \\ &\leq Q(A \cap (\cup_j \{X_j^* \in B_j^*\}) \cap (\cup_r \{X_r \notin B_r\})) \\ &\leq \sum_j \sum_r Q(A, X_j^* \in B_j^*, X_r \notin B_r) \\ &= \sum_j Q(A, X_j^* \in B_j^*, X_j \notin B_j) \\ &\leq \sum_j Q(A_j, X_j^* \in B_j^*) = 0. \end{aligned}$$

Thus, H is Q -TIP, and this concludes the proof. \square

By Theorem 8, $\mathcal{D} = \mathcal{N}$ in case $\gamma \ll \mu$ and condition (5) holds with $B = \{f > 0\}$.

Example 9. Let $X_3 = X_1 X_2$ where X_1 and X_2 are i.i.d. random variables with values in $\{-1, 1\}$ and $P(X_1 = -1) = P(X_1 = 1) = \frac{1}{2}$. Let $\mu_1 = \mu_2 = \mu_3$ be counting measure on $\{-1, 1\}$ and $\mathcal{D}_0 = \cap_i \overline{\sigma(X_i)}$. Since the X_i are pairwise independent (even if not independent), $\mathcal{D}_0 = \mathcal{N}$. Since $P(X_3 = 1) = \frac{1}{2}$ and $\mathcal{D} \supset \sigma(X_3)$, then $\mathcal{D} \neq \mathcal{N}$. Thus, $\mathcal{D}_0 = \mathcal{N}$ and $\gamma \ll \mu$ do not imply $\mathcal{D} = \mathcal{N}$. Note also that $\cup_{i=1}^3 \{X_i = 1\} = \Omega$, $P(X_1 = X_2 = X_3 = 1) > 0$ while $H = \Omega$ is not TIP. Thus, condition (5) cannot be weakened into

$$\cup_{i=1}^k \{X_i \in B_i\} \supset H \supset \cap_{i=1}^k \{X_i \in B_i\} \quad \text{with} \quad P(X_1 \in B_1, \dots, X_k \in B_k) > 0.$$

Our last and main result is that \mathcal{D} is atomic under P as far as $\gamma \ll \mu$.

Theorem 10. *Let $\mathcal{A}_i = \sigma(X_i^*)$ for all i . If $\gamma \ll \mu$, then $P|\mathcal{D}$ is atomic.*

Proof. Let Q be a probability measure on \mathcal{F} which makes X_1, \dots, X_k independent. Denote M_Q the class of those probabilities \mathbb{P} on \mathcal{F} such that $\mathbb{P} \ll Q$ and

$$\mathcal{N}_{\mathbb{P}} = \{F \in \mathcal{F} : \mathbb{P}(F) \in \{0, 1\}\}, \quad \mathcal{D}_{\mathbb{P}} = \cap_{i=1}^k \sigma(\sigma(X_i^*) \cup \mathcal{N}_{\mathbb{P}}), \quad \text{with } \mathbb{P} \in M_Q.$$

Arguing by induction on k , we now prove that each $\mathbb{P} \in M_Q$ is atomic on $\mathcal{D}_{\mathbb{P}}$.

Let $k = 2$ and $\mathbb{P} \in M_Q$. Since $X_1^* = X_2$ and $X_2^* = X_1$, then \mathbb{P} is atomic on $\mathcal{D}_{\mathbb{P}}$ by Theorem 4.

Given $k \geq 3$, define $V_i = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_{k-1})$. By induction, suppose that each $\mathbb{P} \in M_Q$ is atomic on

$$\mathcal{V}_{\mathbb{P}} = \cap_{i=1}^{k-1} \sigma(\sigma(V_i) \cup \mathcal{N}_{\mathbb{P}}).$$

We have to prove that each $\mathbb{P} \in M_Q$ is atomic on $\mathcal{D}_{\mathbb{P}}$. Accordingly, we fix $\mathbb{P} \in M_Q$ and a $\mathcal{D}_{\mathbb{P}}$ -measurable function $U : \Omega \rightarrow \mathbb{R}$. Arguing as in the proof of Theorem 4, it suffices to show that $\mathbb{P}(U \in C) = 1$ for some countable set $C \subset \mathbb{R}$.

Since $\sigma(U) \subset \mathcal{D}_{\mathbb{P}}$ and $\mathcal{A}_i = \sigma(X_i^*)$, one obtains $\mathbb{P}(U = f_i(X_i^*)) = 1$, $i = 1, \dots, k$, for some real measurable function f_i on $(\prod_{j \neq i} \mathcal{X}_j, \prod_{j \neq i} \mathcal{B}_j)$. Let

$$A_x = \{f_1(V_1, x) = \dots = f_{k-1}(V_{k-1}, x)\} \quad \text{for } x \in \mathcal{X}_k, \\ F(t, x) = Q(A_x \cap \{f_1(V_1, x) \leq t\}) \quad \text{for } t \in \mathbb{R} \text{ and } x \in \mathcal{X}_k.$$

Since $F(t, \cdot)$ is \mathcal{B}_k -measurable for fixed t , F is a real cadlag process on the measurable space $(\mathcal{X}_k, \mathcal{B}_k)$. Let $J = \{(t, x) : F(t, x) > F(t-, x)\}$ be the jump set of F . By a well known result (see e.g. [9], Proposition 2.26), J is contained in a countable union of graphs, that is,

$$J \subset \cup_n \{(g_n(x), x) : x \in \mathcal{X}_k\}$$

for suitable \mathcal{B}_k -measurable functions $g_n : \mathcal{X}_k \rightarrow \mathbb{R}$, $n = 1, 2, \dots$

Fix $x \in \mathcal{X}_k$ with $Q(A_x) > 0$ and define $Q_x(\cdot) = Q(\cdot | A_x)$. Then, Q_x is atomic on \mathcal{V}_{Q_x} (since $Q_x \in M_Q$) and $f_1(V_1, x)$ is \mathcal{V}_{Q_x} -measurable (since $Q_x(A_x) = 1$). Thus, $F(\cdot, x)$ is a purely jump function, that is, $Q(A_x \cap \{f_1(V_1, x) \notin J_x\}) = 0$ where $J_x = \{t : F(t, x) > F(t-, x)\}$. Integrating over x yields

$$Q(f_1(X_1^*) = \dots = f_{k-1}(X_{k-1}^*)) \\ = Q(f_1(X_1^*) = \dots = f_{k-1}(X_{k-1}^*) = g_n(X_k) \text{ for some } n).$$

Let $C = \{c \in \mathbb{R} : Q(f_k(X_k^*) = c) > 0\}$. Since $f_k(X_k^*)$ and $g_n(X_k)$ are independent under Q , then $Q(f_k(X_k^*) \notin C, f_k(X_k^*) = g_n(X_k)) = 0$ for all n . Hence,

$$\begin{aligned} & Q(f_k(X_k^*) \notin C \text{ and } f_1(X_1^*) = \dots = f_k(X_k^*)) \\ & \leq Q(f_k(X_k^*) \notin C \text{ and } f_k(X_k^*) = g_n(X_k) \text{ for some } n) = 0. \end{aligned}$$

Therefore, $\mathbb{P} \ll Q$ and $\mathbb{P}(U = f_i(X_i^*)) = 1$ for all $i = 1, \dots, k$ imply

$$\mathbb{P}(U \in C) = 1 - \mathbb{P}(f_k(X_k^*) \notin C \text{ and } f_1(X_1^*) = \dots = f_k(X_k^*)) = 1.$$

Since C is countable, \mathbb{P} is atomic on $\mathcal{D}_{\mathbb{P}}$. This concludes the induction argument and proves that each $\mathbb{P} \in M_Q$ is atomic on $\mathcal{D}_{\mathbb{P}}$.

Finally, to prove P atomic on $\mathcal{D} = \mathcal{D}_P$, it can be assumed $(\Omega, \mathcal{F}, P) = (\mathcal{X}, \mathcal{B}, \gamma)$ and X_1, \dots, X_k the canonical projections. Also, since μ is a σ -finite product measure, μ is equivalent to some probability Q on $\mathcal{B} = \mathcal{F}$ which makes X_1, \dots, X_k independent. Hence, $\gamma \ll \mu$ implies $P = \gamma \ll Q$. This concludes the proof. \square

Note that Theorem 4 could be obtained as a corollary of previous Theorem 10. However, Theorem 4 has been stated as an autonomous result, since it is a useful preliminary step toward Theorem 10.

Finally, the scope of Theorems 8 and 10 can be enlarged via Lemma 6. Following this route, sometimes, the assumption $\gamma \ll \mu$ can be circumvented. Let $Z_i : \mathcal{X} \rightarrow \mathcal{X}_i$ denote the i -th canonical projection and $Z_i^* = (Z_1, \dots, Z_{i-1}, Z_{i+1}, \dots, Z_k)$. Moreover, suppose γ is equivalent on rectangles to some probability ν on \mathcal{B} , i.e., $\gamma(A) = 0 \Leftrightarrow \nu(A) = 0$ for each set A of the form $A = \{Z_1^* \in C_1, \dots, Z_k^* \in C_k\}$ with $C_i \in \prod_{j \neq i} \mathcal{B}_j$ for all i . Then, \mathcal{D} is atomic under P provided $\nu \ll \mu$; cfr. Lemma 6 and Theorem 10. Or else, $\mathcal{D} = \mathcal{N}$ whenever $\nu \ll \mu$ and

$$\cup_{i=1}^k \{Z_i^* \in B_i^*\} \supset \left\{ \frac{d\nu}{d\mu} > 0 \right\} \supset \cap_{i=1}^k \{Z_i \in B_i\}$$

for some B_1, \dots, B_k such that $\nu(Z_1 \in B_1, \dots, Z_k \in B_k) > 0$; cfr. Lemma 6 and Theorem 8. Note also that, for $k = 2$, equivalence on rectangles reduces to

$$\gamma(B_1 \times B_2) = 0 \Leftrightarrow \nu(B_1 \times B_2) = 0 \quad \text{whenever } B_1 \in \mathcal{B}_1, B_2 \in \mathcal{B}_2.$$

As an example (suggested by an anonymous referee) suppose $(X_n : n \geq 1)$ is an exchangeable sequence of real random variables with Ferguson-Dirichlet mixing measure. For $k = 2$, (X_1, X_2) is distributed as

$$\gamma(B_1 \times B_2) = a \beta(B_1) \beta(B_2) + (1 - a) \beta(B_1 \cap B_2)$$

where $0 < a < 1$ and β is a probability on the real Borel sets. Then $\mathcal{D} = \mathcal{N}$, as γ is equivalent on rectangles to $\beta \times \beta$. However, if β is nonatomic, γ fails to be absolutely continuous with respect to any σ -finite product measure. It can be shown that, for every $k \geq 2$, one obtains $\mathcal{D} = \mathcal{N}$ for (X_1, \dots, X_k) as well.

Acknowledgment: This paper benefited from the helpful suggestions of an anonymous referee.

REFERENCES

- [1] Berti P., Pratelli L., Rigo P. (2006) Asymptotic behaviour of the empirical process for exchangeable data, *Stoch. Proc. Appl.*, 116, 337-344.
- [2] Berti P., Pratelli L., Rigo P. (2007) Skorohod representation on a given probability space, *Prob. Theo. Rel. Fields*, 137, 277-288.
- [3] Berti P., Pratelli L., Rigo P. (2008) Trivial intersection of σ -fields and Gibbs sampling, *Ann. Probab.*, 36, 2215-2234.
- [4] Bhaskara Rao K.P.S., Bhaskara Rao M. (1983) *Theory of charges*, Academic Press.
- [5] Burkholder D.L., Chow Y.S. (1961) Iterates of conditional expectation operators, *Proc. Amer. Math. Soc.*, 12, 490-495.
- [6] Delyon B., Delyon F. (1999) Generalization of von Neumann's spectral sets and integral representation of operators, *Bull. Soc. Math. Fr.*, 127, 25-41.
- [7] Diaconis P., Khare K., Saloff-Coste L. (2007) Stochastic alternating projections, *Preprint*, Dept. of Statistics, Stanford University, currently available at: <http://www-stat.stanford.edu/~cgates/PERSI/papers/altproject-2.pdf>
- [8] Halmos P.R. (1947) On the set of values of a finite measure, *Bull. Amer. Math. Soc.*, 53, 138-141.
- [9] Karatzas I., Shreve S.E. (1991) *Brownian motion and stochastic calculus* (second edition), Springer.
- [10] San Martin E., Mouchart M., Rolin J.M. (2005) Ignorable common information, null sets and Basu's first theorem, *Sankhya*, 67, 674-698.

PATRIZIA BERTI, DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA "G. VITALI", UNIVERSITA' DI MODENA E REGGIO-EMILIA, VIA CAMPI 213/B, 41100 MODENA, ITALY

E-mail address: `patrizia.berti@unimore.it`

LUCA PRATELLI, ACCADEMIA NAVALE, VIALE ITALIA 72, 57100 LIVORNO, ITALY

E-mail address: `pratel@mail.dm.unipi.it`

PIETRO RIGO (CORRESPONDING AUTHOR), DIPARTIMENTO DI ECONOMIA POLITICA E METODI QUANTITATIVI, UNIVERSITA' DI PAVIA, VIA S. FELICE 5, 27100 PAVIA, ITALY

E-mail address: `prigo@eco.unipv.it`