

Packham, Natalie; Kalkbrener, Michael; Overbeck, Ludger

**Working Paper**

## Default probabilities and default correlations under stress

Frankfurt School - Working Paper Series, No. 211

**Provided in Cooperation with:**

Frankfurt School of Finance and Management

*Suggested Citation:* Packham, Natalie; Kalkbrener, Michael; Overbeck, Ludger (2014) : Default probabilities and default correlations under stress, Frankfurt School - Working Paper Series, No. 211, Frankfurt School of Finance & Management, Frankfurt a. M.

This Version is available at:

<https://hdl.handle.net/10419/97176>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

**Frankfurt School – Working Paper Series**

**No. 211**

**Default probabilities and default  
correlations under stress**

by Natalie Packham, Michael Kalkbrener, Ludger Overbeck

April 2014



**Frankfurt School of  
Finance & Management**  
Bankakademie | HfB

Sonnemannstr. 9–11 60314 Frankfurt am Main, Germany  
Phone: +49 (0) 69 154 008 0 Fax: +49 (0) 69 154 008 728  
Internet: [www.frankfurt-school.de](http://www.frankfurt-school.de)

## Abstract

We investigate default probabilities and default correlations of Merton-type credit portfolio models in stress scenarios where a common risk factor is truncated. The analysis is performed in the class of elliptical distributions, a family of light-tailed to heavy-tailed distributions encompassing many distributions commonly found in financial modelling. It turns out that the asymptotic limit of default probabilities and default correlations depend on the max-domain of the elliptical distribution's mixing variable. In case the mixing variable is regularly varying, default probabilities are strictly smaller than 1 and default correlations are in  $(0; 1)$ . Both can be expressed in terms of the Student t-distribution function. In the rapidly varying case, default probabilities are 1 and default correlations are 0. We compare our results to the tail dependence function and discuss implications for credit portfolio modelling.

Keywords: financial risk management, credit portfolio modelling, stress testing, elliptic distribution, max-domain

MSC classification: 60G70, 91G40

ISSN: 1436-9753

### Contact:

Natalie Packham  
Juniorprofessorin für Quantitative Finance  
Frankfurt School of Finance & Management  
Sonnemannstraße 9-11  
60314 Frankfurt am Main  
Phone: +49-69-154008-723  
n.packham@fs.de

Michael Kalkbrener  
Deutsche Bank AG  
Taunusanlage 12  
60325 Frankfurt am Main

Ludger Overbeck  
Institute of Mathematics  
University of Giessen  
Arndtstrasse 2  
35392 Giessen

## Contents

<b>1</b>	<b>Introduction</b>	<b>4</b>
<b>2</b>	<b>Preliminaries</b>	<b>6</b>
2.1	Structural credit portfolio models . . . . .	6
2.2	Distribution of model variables . . . . .	7
2.3	Stress testing in credit portfolio models . . . . .	8
<b>3</b>	<b>Default probabilities under stress</b>	<b>8</b>
<b>4</b>	<b>Default correlations under stress</b>	<b>17</b>
<b>5</b>	<b>Implications for credit portfolio modelling</b>	<b>22</b>

# 1 Introduction

In the aftermath of the subprime crisis and the European sovereign debt crisis, stress testing of bank portfolios has become an integral part of financial risk management and banking supervision (Turner, 2009; Larosi re and others, 2009; Brunnermeier *et al.*, 2009; BIS, 2009). Stress tests for credit portfolios are of particular importance, since in a typical bank risk capital for credit risk far outweighs capital requirements for any other risk class.

In this paper, we analyse the behaviour of credit portfolio models under stress depending on the joint distribution of the stochastic variables of the model. Although widely questioned, the industry standard is still to employ multivariate normally distributed random variables. In order to cover a wide range of light-tailed to heavy-tailed distributions we use the family of elliptical distributions, which contains the normal distribution as a special case. More formally, let  $Z = (Z_0, \dots, Z_d)^T$  be a random vector on the probability space  $(\Omega, \mathcal{A}, \mathbf{P})$ . We assume that  $Z$  follows an *elliptical distribution* with representation

$$Z \stackrel{\mathcal{L}}{=} GAU, \tag{1}$$

where  $G > 0$  is a scalar random variable, the so-called *mixing variable*,  $A$  is a deterministic  $(d+1) \times (d+1)$  matrix with  $AA^T := \Sigma$ , which in turn is a  $(d+1) \times (d+1)$  nonnegative definite symmetric matrix of rank  $d+1$ , and  $U$  is a  $(d+1)$ -dimensional random vector uniformly distributed on the unit sphere  $\mathcal{S}_{d+1} := \{z \in \mathbb{R}^{d+1} : z^T z = 1\}$ , and  $U$  is independent of  $G$ . Recent papers study the asymptotic properties of value-at-risk in a similar setup, e.g. Embrechts *et al.* (2009); Mainik and Embrechts (2013).

In the next section, we provide a short survey of structural credit portfolio models. In this setting,  $Z_0$  will be interpreted as a risk factor of the model and  $Z_1, \dots, Z_d$  as asset return variables of  $d$  firms. The default of the  $i$ -th firm is represented by  $\{Z_i \leq D_i\}$  for a given default threshold  $D_i \in \mathbb{R}$  and the corresponding default probability (PD) is defined by  $p_i := \mathbf{P}(Z_i \leq D_i) = \mathbb{E}(\mathbf{1}_{\{Z_i \leq D_i\}})$ . The default correlations are defined as the correlations of the default indicators  $\mathbf{1}_{\{Z_i \leq D_i\}}$  and  $\mathbf{1}_{\{Z_j \leq D_j\}}$ . To simplify the exposition, we assume throughout that

the correlations of  $Z_0, Z_1, \dots, Z_d$  are in  $(0, 1)$ .

The objective of this paper is to analyse the impact of stress on default probabilities and default correlations. Stress scenarios are specified by truncating the risk factor  $Z_0$ , i.e., by conditioning on  $\{Z_0 \leq C\}$  with stress level  $C \in \mathbb{R}$ . Using techniques from Extreme Value Theory (EVT), we derive the limit of conditional default probabilities and default correlations as  $C \rightarrow -\infty$ . The limit depends on whether the mixing variable  $G$  is in the max-domain of the Fréchet or the Gumbel distribution, or more generally, on whether the tail distribution function  $\mathbf{P}(G > \cdot)$  is regularly varying or rapidly varying. For stressed default probabilities, we show that for any  $D_i \in \mathbb{R}$

$$\lim_{C \rightarrow -\infty} \mathbf{P}(Z_i \leq D_i | Z_0 \leq C) = 1,$$

if  $\mathbf{P}(G > \cdot)$  is rapidly varying. In contrast, if  $\mathbf{P}(G > \cdot)$  is regularly varying with tail index  $-\alpha$ , then

$$\lim_{C \rightarrow -\infty} \mathbf{P}(Z_i \leq D_i | Z_0 \leq C) = t_{\alpha+1} \left( \frac{\sqrt{\alpha+1} \rho}{\sqrt{1-\rho^2}} \right) \in [1/2, 1),$$

where  $t_v$  denotes the Student  $t$  distribution function with parameter  $v$  and  $\rho$  denotes the correlation of  $Z_0$  and  $Z_i$ . These results imply that the limiting default probability under stress is strictly smaller in the heavy-tailed case than in the light-tailed case. Essentially, in the heavy-tailed case, extreme outcomes are driven by the joint mixing variable, implying a strictly positive probability for a conditional extreme positive outcome of  $Z_i$ .

It is interesting to note that this behaviour of limiting default probabilities is fundamentally different to tail dependence, which is positive for heavy-tailed  $G$ , and converges to 0 as the tail index of  $G$  tends to infinity, that is, to the light-tailed case, see (Schmidt, 2002; Klüppelberg *et al.*, 2008; Hult and Lindskog, 2002). Limiting default correlations, on the other hand, behave like tail dependence: we show that  $\text{Corr}(\mathbf{1}_{\{Z_i \leq D_i\}}, \mathbf{1}_{\{Z_j \leq D_j\}} | Z_0 \leq C)$  converges to 0 in the light-tailed case and to a positive number in the heavy-tailed case.

The paper is structured as follows: in Section 2, we define stress tests in structural credit portfolio models. The results on asymptotic stressed default probabilities are derived in section 3. Section 4 focuses on stressed default correlations. In section 5, implications for credit

portfolio modelling are discussed.

## 2 Preliminaries

### 2.1 Structural credit portfolio models

Depending on their formulation, credit portfolio models can be divided into reduced-form models and structural (or firm-value) models. The progenitor of all structural models is the model of Merton (Merton, 1974), which links the default of a firm to the relationship between its assets and the liabilities at the end of a given time period  $[0, T]$ . More precisely, in a structural credit portfolio model the  $i$ -th counterparty defaults if its asset return (or ability-to-pay) variable  $Z_i$  falls below a default threshold  $D_i$ : the default event at time  $T$  is defined as  $\{Z_i \leq D_i\} \subseteq \Omega$ , where  $Z_i$  is a real-valued random variable on the probability space  $(\Omega, \mathcal{A}, \mathbf{P})$  and  $D_i \in \mathbb{R}$ . The portfolio loss variable is defined by

$$L := \sum_{i=1}^d l_i \cdot \mathbf{1}_{\{Z_i \leq D_i\}}, \quad (2)$$

where  $d$  denotes the number of counterparties and  $l_i$  is the loss-at-default of the  $i$ -th counterparty. In order to reflect risk concentrations, each  $Z_i$  is decomposed into a sum of systematic factors  $X_1, \dots, X_m$ , which are often identified with geographic regions or industries, and a firm-specific factor  $\varepsilon_i$ , that is,

$$Z_i = \sqrt{R_i^2} \sum_{j=1}^m w_{ij} X_j + \sqrt{1 - R_i^2} \varepsilon_i. \quad (3)$$

The impact of the risk factors on  $Z_i$  is determined by  $R_i^2 \in [0, 1]$  and the factor weights  $w_{ij} \in \mathbb{R}$ .

In order to quantify portfolio risk, measures of risk are applied to the portfolio loss distribution (2). The expected loss of the credit portfolio is used for specifying credit reserves. It is defined as the mean of  $L$ :

$$\mathbb{E}(L) = \sum_{i=1}^d l_i \cdot p_i,$$

where  $p_i = \mathbf{P}(Z_i \leq D_i) = \mathbb{E}(\mathbf{1}_{\{Z_i \leq D_i\}})$  denotes the default probability of the  $i$ -th counterparty.

Capital requirements for covering unexpected losses are typically derived from the value-at-risk  $\text{VaR}_\alpha(L)$  for a predefined probability  $\alpha \in (0, 1)$ , where  $\text{VaR}_\alpha(L)$  is simply defined as the  $\alpha$ -quantile of  $L$ . Obviously, the default probabilities and risk concentrations specified by the dependence structure of the default variables  $\mathbf{1}_{\{Z_i \leq D_i\}}$  determine the value-at-risk of the credit portfolio. Default correlations

$$\text{Corr}(\mathbf{1}_{\{Z_i \leq D_i\}}, \mathbf{1}_{\{Z_j \leq D_j\}}) = \frac{\mathbf{P}(Z_i \leq D_i, Z_j \leq D_j) - p_i p_j}{\sqrt{p_i(1-p_i)p_j(1-p_j)}}$$

are used as a measure of dependence by portfolio management to identify risk concentrations on counterparty level.

## 2.2 Distribution of model variables

The standard approach in credit risk management is to model the risk factors and ability-to-pay variables through a joint multi-variate Gaussian distribution. Since the purpose of this paper is to analyze the impact of stress scenarios under different distribution assumptions we use a more general framework and consider elliptical distributions instead.

Elliptical distributions cover a variety of light-tailed to heavy-tailed distributions depending on the tail behaviour of the mixing variable  $G$ , i.e., whether  $G$  is rapidly varying or regularly varying. A special role is played by *normal variance mixture (NVM) distributions* (see, for instance, McNeil *et al.* (2005); Bingham and Kiesel (2002)). First, NVM distributions encompass a number of distributions commonly used in financial modelling, most prominently normal distributions,  $t$ -distributions and symmetric generalised hyperbolic distributions. Second, all elliptical distributions of interest in the credit portfolio context can be represented as NVM distributions: any elliptical distribution whose so-called characteristic generator does not depend on the dimension  $d$  can be represented as an NVM distribution, see Theorem 2.21 of Fang *et al.* (1990), or Theorem 3.25 of McNeil *et al.* (2005). For details on elliptical distributions, we refer to (Fang *et al.*, 1990; Cambanis *et al.*, 1981) and for their application in finance and risk management we refer to (McNeil *et al.*, 2005).



## 2.3 Stress testing in credit portfolio models

In a stress test, credit portfolios are typically evaluated under the assumption of adverse economic conditions. A natural way for implementing stress tests in portfolio models is to translate the stress scenario into constraints on risk factors. In our setup, the constraints are formalised by truncating risk factor variables  $X_1, \dots, X_m$ , that is, by conditioning on the range of values that a risk factor may attain. This is a commonly used stress testing technique for credit risk management and capital management of financial institutions, see e.g. (Bonti *et al.*, 2006; Duellmann and Erdelmeier, 2009; Kalkbrener and Packham, 2013). More precisely, let us consider the situation when the risk factor  $Z_0 \in \{X_1, \dots, X_m\}$  is truncated by  $C \in \mathbb{R}$ , that is,  $Z_0 \leq C$  and write

$$\mathbf{P}^C(A) = \mathbf{P}(A|Z_0 \leq C), \quad A \in \mathcal{A},$$

for the corresponding conditional distribution. In this setting,  $C$  is interpreted as the level of stress applied to the risk factor  $Z_0$ . The objective of this paper is to calculate the limit of default probabilities, joint default probabilities and corresponding default correlations under  $\mathbf{P}^C$  as  $C \rightarrow -\infty$ .

## 3 Default probabilities under stress

Let  $h$  be a positive, Lebesgue-measurable function on  $(0, \infty)$ . We write  $h \in RV_\alpha$  if  $h$  is regularly varying with index  $\alpha \in \mathbb{R}$ , i.e.,

$$\lim_{x \rightarrow \infty} \frac{h(tx)}{h(x)} = t^\alpha, \quad t > 0,$$

and  $h \in RV_{-\infty}$  if  $h$  is rapidly varying with index  $-\infty$ , i.e.,

$$\lim_{x \rightarrow \infty} \frac{h(tx)}{h(x)} = \begin{cases} 0, & t > 1, \\ \infty, & 0 < t < 1. \end{cases}$$

For details on regularly varying functions, we refer to Bingham *et al.* (1987).

Let  $Z = GAU$  denote an elliptical random vector as in Equation (1). We assume that all variables are standardised so that  $\Sigma = AA^T$  is the correlation matrix of  $(Z_0, \dots, Z_d)^T$ . The correlation of  $Z_i$  and  $Z_j$  is denoted by  $\rho_{ij}$ ,  $i, j = 0, 1, \dots, d$ . We assume that the correlations with respect to the risk factor are positive, i.e.,  $\rho_{0i} > 0$ . The case  $\rho_{0i} \leq 0$  can be treated analogously.

In the following, denote by  $A_i$  the  $i$ -th row of  $A$  and let  $F_U$  denote the uniform distribution on  $\mathcal{S}_{d+1}$ .

It is well-known that  $\mathbf{P}(G > \cdot) \in RV_{-\alpha}$  implies  $\mathbf{P}(Z_i > \cdot) \in RV_{-\alpha}$ ,  $i = 0, \dots, d$ , see e.g. Theorem 7.35 of McNeil *et al.* (2005). For many distributions of interest in  $RV_{-\infty}$ , such as the normal distribution and the generalised hyperbolic distributions, we know that the mixing variable is in  $RV_{-\infty}$ , see e.g. Section 7.3 of McNeil *et al.* (2005).

**Theorem 1.** (i) If  $\mathbf{P}(G > \cdot) \in RV_{-\alpha}$ , then

$$\begin{aligned} & \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1, \dots, Z_d \leq D_d) \\ &= \int_{u \in \mathcal{S}_{d+1}, A_{0 \cdot} u > 0, \dots, A_{d \cdot} u > 0} (A_{0 \cdot} u)^\alpha dF_U(u) \left( \int_{u \in \mathcal{S}_{d+1}, A_{0 \cdot} u > 0} (A_{0 \cdot} u)^\alpha dF_U(u) \right)^{-1}. \end{aligned}$$

(ii) If  $\mathbf{P}(G > \cdot) \in RV_{-\infty}$ , then

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1, \dots, Z_d \leq D_d) = 1.$$

*Proof.* We first give a proof for the special case  $D_i = 0$  for  $i = 1, \dots, d$ .

Since the elliptical random vector is symmetric and continuous we can write

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq 0, \dots, Z_d \leq 0) = \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_0 > C, Z_1 > 0, \dots, Z_d > 0)}{\mathbf{P}(Z_0 > C)}. \quad (4)$$

For the numerator

$$\begin{aligned}\mathbf{P}(Z_0 > C, Z_1 > 0, \dots, Z_d > 0) &= \mathbf{P}\left(G > \frac{C}{A_0 \cdot U}, A_0 \cdot U > 0, \dots, A_d \cdot U > 0\right) \\ &= \int_{u \in \mathcal{S}_{d+1}, A_i \cdot u > 0, i=0, \dots, d} \mathbf{P}\left(G > \frac{C}{A_0 \cdot u}\right) F_U(\mathrm{d}u). \quad (5)\end{aligned}$$

For (i), it follows from  $\mathbf{P}(G > \cdot) \in RV_{-\alpha}$  that

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}(G > C/(A_0 \cdot u))}{\mathbf{P}(G > C)} = (A_0 \cdot u)^\alpha, \quad \text{for } A_0 \cdot u > 0.$$

Potter's bounds (de Haan and Ferreira, 2006, Proposition B.1.9) state that for arbitrary  $\varepsilon > 0$  and  $\delta > 0$  there exists  $C_0$  such that for all  $C \geq C_0$ ,  $C/(A_0 \cdot u) \geq C_0$ ,

$$\frac{\mathbf{P}(G > C/(A_0 \cdot u))}{\mathbf{P}(G > C)} < (1 + \varepsilon)(A_0 \cdot u)^\alpha \max\left((A_0 \cdot u)^\delta, (A_0 \cdot u)^{-\delta}\right),$$

and since the right-hand side is integrable, we obtain by Dominated Convergence that

$$\lim_{C \rightarrow \infty} \int_{u \in \mathcal{S}_{d+1}, A_i \cdot u > 0, i=0, \dots, d} \frac{\mathbf{P}(G > C/(A_0 \cdot u))}{\mathbf{P}(G > C)} F_U(\mathrm{d}u) = \int_{u \in \mathcal{S}_{d+1}, A_i \cdot u > 0, i=0, \dots, d} (A_0 \cdot u)^\alpha F_U(\mathrm{d}u).$$

Applying the same method to the denominator of Equation (4) completes the proof of (i).

For (ii), it suffices to consider the case  $d = 1$ , i.e.,  $\lim_{C \rightarrow \infty} \mathbf{P}(Z_1 > 0 | Z_0 > C) = 1$ , since the general case follows from

$$\mathbf{P}(Z_1 > 0, \dots, Z_d > 0 | Z_0 > C) \geq 1 - \sum_{i=1}^d (1 - \mathbf{P}(Z_i > 0 | Z_0 > C)).$$

Equality (5) implies

$$\begin{aligned}\mathbf{P}(Z_1 > 0 | Z_0 > C) &= \\ &= \int_{u \in \mathcal{S}_2, A_i \cdot u > 0, i=0,1} \mathbf{P}\left(G > \frac{C}{A_0 \cdot u}\right) F_U(\mathrm{d}u) \left( \int_{u \in \mathcal{S}_2, A_0 \cdot u > 0} \mathbf{P}\left(G > \frac{C}{A_0 \cdot u}\right) F_U(\mathrm{d}u) \right)^{-1}.\end{aligned}$$

Write  $u \in S_2$  in polar coordinates as  $u = (\cos \theta, \sin \theta)$ ,  $\theta \in [-\pi, \pi]$ , and let  $A$  be the Cholesky decomposition of the correlation matrix, i.e.,  $A_0 = (1, 0)^T$ ,  $A_1 = (\rho, \sqrt{1-\rho^2})$  with  $\rho := \rho_{01}$  the correlation of  $Z_0, Z_1$ . Hence,  $A_0 \cdot u = \cos \theta > 0$  if  $\theta \in (-\pi/2, \pi/2)$  and  $A_1 \cdot u = \rho \cos \theta + \sqrt{1-\rho^2} \sin \theta = \sin(\theta + \arcsin \rho) > 0$  if  $\theta \in (-\arcsin \rho, \pi - \arcsin \rho)$ . It follows that

$$\begin{aligned} \int_{u \in \mathcal{S}_2, A_i \cdot u > 0, i=0,1} \mathbf{P}\left(G > \frac{C}{A_0 \cdot u}\right) F_U(du) &= \int_{-\arcsin \rho}^{\pi/2} \mathbf{P}\left(G > \frac{C}{\cos \theta}\right) d\theta \\ &= \int_{-\pi/2}^{\pi/2} \mathbf{P}\left(G > \frac{C}{\cos \theta}\right) d\theta - \int_{\arcsin \rho}^{\pi/2} \mathbf{P}\left(G > \frac{C}{\cos \theta}\right) d\theta \end{aligned}$$

and

$$\int_{u \in \mathcal{S}_2, A_0 \cdot u > 0} \mathbf{P}\left(G > \frac{C}{A_0 \cdot u}\right) F_U(du) = \int_{-\pi/2}^{\pi/2} \mathbf{P}\left(G > \frac{C}{\cos \theta}\right) d\theta.$$

Since  $\rho > 0$ , we have  $\cos \theta < \cos(\arcsin \rho) = \sqrt{1-\rho^2}$  for  $\theta \in (\arcsin \rho, \pi/2)$ . Hence, by definition of rapidly varying functions and by Dominated Convergence,

$$\lim_{C \rightarrow \infty} \int_{\arcsin \rho}^{\pi/2} \frac{\mathbf{P}\left(G > \frac{C}{\sqrt{1-\rho^2}} \frac{\sqrt{1-\rho^2}}{\cos \theta}\right)}{\mathbf{P}\left(G > \frac{C}{\sqrt{1-\rho^2}}\right)} d\theta = 0.$$

On the other hand, for  $\theta \in (-\arcsin \rho, \arcsin \rho)$ ,

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}\left(G > \frac{C}{\sqrt{1-\rho^2}} \frac{\sqrt{1-\rho^2}}{\cos \theta}\right)}{\mathbf{P}\left(G > \frac{C}{\sqrt{1-\rho^2}}\right)} = \infty,$$

so that, putting everything together, we obtain

$$\lim_{C \rightarrow \infty} \mathbf{P}(Z_1 > 0 | Z_0 > C) = 1.$$

It remains to show that

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq 0, \dots, Z_d \leq 0) = \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1, \dots, Z_d \leq D_d), \quad (6)$$

for arbitrary  $D_1, \dots, D_d$ . Let  $i \in \{1, \dots, d\}$  and  $a > 0$ . Note that for  $C < -|D_i|/a$ ,

$$\mathbf{P}^C(Z_i - aZ_0 \leq 0) \leq \mathbf{P}^C(Z_i \leq aC) < \mathbf{P}^C(Z_i \leq D_i) < \mathbf{P}^C(Z_i \leq -aC) \leq \mathbf{P}^C(Z_i + aZ_0 \leq 0).$$

Hence,

$$\begin{aligned} \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 - aZ_0 \leq 0, Z_2 \leq 0, \dots, Z_d \leq 0) &\leq \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1, Z_2 \leq 0, \dots, Z_d \leq 0) \\ &\leq \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 + aZ_0 \leq 0, Z_2 \leq 0, \dots, Z_d \leq 0). \end{aligned}$$

Since Equation (5) is continuous in  $A_1$ , it follows that  $\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 + aZ_0 \leq 0, Z_2 \leq 0, \dots, Z_d \leq 0)$  is a continuous function in  $a \in \mathbb{R}$ , hence

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1, Z_2 \leq 0, \dots, Z_d \leq 0) = \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq 0, Z_2 \leq 0, \dots, Z_d \leq 0).$$

and therefore (6) is obtained by reiterating this argument.  $\square$

### Remarks

- (i) Default thresholds  $D_1, \dots, D_d$  determine the unconditional default probabilities  $\mathbf{P}(Z_i \leq D_i)$ . Note, however, that  $\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1, \dots, Z_d \leq D_d)$  does not depend on the  $D_i$ , i.e., in the limit, stressed default probabilities do not depend on the unconditional default probabilities, but only on the dependence structure of the  $Z_i$ .
- (ii) Theorem 1 implies in particular that the limiting default probability under stress is strictly smaller than 1 in the heavy-tailed case, provided that the variables are not perfectly correlated. This result can be attributed to the special structure of elliptical distributions, where a stress event may be caused by a large mixing variable and a uniform random vector on the sphere  $\mathcal{S}_{d+1}$  with components close to zero, whose signs may well differ, thus overall leading to potentially very large positive or negative realisations of the asset returns. In the light-tailed case, the tail behaviour of the mixing variable is too moderate to produce extreme overall behaviour of opposite signs.

(iii) Even without further making the limiting distribution of part (i) concrete, it can be numerically determined efficiently using Monte Carlo simulation. An efficient method to simulate uniform random variates on the unit sphere  $\mathcal{S}_{d+1}$ , is to draw  $(d+1)$ -dimensional independent normally distributed random variables  $Y = (Y_0, \dots, Y_d)^T$ , and transform them according to  $Y / \|Y\|$ , which produces the desired random variates, see e.g. Corollary 3.23 of McNeil *et al.* (2005) or Section 3.4.1.E. of Knuth (1998). Simulation has proven to be significantly faster than the numerical calculation of integrals in Proposition 3 below.

In the following two propositions, we express the integral in Theorem 1(i) in terms of beta functions: the incomplete beta function  $B(z; a, b)$  is defined by

$$B(z; a, b) := \int_0^z u^{a-1} (1-u)^{b-1} du = 2 \int_0^{\arcsin(\sqrt{z})} (\sin \theta)^{2a-1} (\cos \theta)^{2b-1} d\theta, \quad (7)$$

where the last equation follows from substituting  $u = (\sin \theta)^2$ . The regularized incomplete beta function is defined as

$$I_x(a, b) = \frac{B(x; a, b)}{B(a, b)}, \quad (8)$$

where  $B(a, b) := B(1, a, b)$ . Note that there exists the following relationship between an incomplete beta function and the distribution function  $t_v$  of the Student- $t$  distribution with parameter  $v$ :

$$t_v(x) = \begin{cases} \frac{1}{2} I_{v/(x^2+v)}\left(\frac{v}{2}, \frac{1}{2}\right), & x \leq 0, \\ \frac{1}{2} \left[ 1 + I_{x^2/(x^2+v)}\left(\frac{1}{2}, \frac{v}{2}\right) \right], & x > 0. \end{cases} \quad (9)$$

Proposition 2 covers the case  $d = 1$ , which corresponds to stressed default probabilities, whereas Proposition 3 deals with stressed bivariate default probabilities.

**Proposition 2.** *Let  $\mathbf{P}(G > \cdot) \in RV_{-\alpha}$ . Then,*

$$\lim_{C \rightarrow -\infty} \mathbf{P}(Z_1 \leq D_1 | Z_0 \leq C) = \frac{1}{2} + \frac{1}{2} I_{\rho^2}\left(\frac{1}{2}, \frac{\alpha+1}{2}\right) = t_{\alpha+1}\left(\frac{\sqrt{\alpha+1}\rho}{\sqrt{1-\rho^2}}\right) \in [1/2, 1), \quad (10)$$

where  $\rho := \rho_{01}$  denotes the correlation of  $Z_0$  and  $Z_1$ .

*Proof.* By Theorem 1(i),

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1) = \int_{u \in \mathcal{S}_2, A_0.u > 0, A_1.u > 0} (A_0.u)^\alpha dF_U(u) \left( \int_{u \in \mathcal{S}_2, A_0.u > 0} (A_0.u)^\alpha dF_U(u) \right)^{-1}.$$

Write  $u \in \mathcal{S}_2$  in polar coordinates as  $u = (\cos \theta, \sin \theta)$ ,  $\theta \in [-\pi, \pi]$ . As in the proof of Theorem 1(ii) we obtain

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1) = \int_{-\arcsin \rho}^{\pi/2} (\cos \theta)^\alpha d\theta \left( \int_{-\pi/2}^{\pi/2} (\cos \theta)^\alpha d\theta \right)^{-1}.$$

Using Equalities (7) and (8) yields

$$\begin{aligned} \int_{-\pi/2}^{\pi/2} (\cos \theta)^\alpha d\theta &= B\left(\frac{1}{2}, \frac{\alpha+1}{2}\right), \\ \int_{-\arcsin \rho}^{\pi/2} (\cos \theta)^\alpha d\theta &= \int_0^{\arcsin \rho} (\cos \theta)^\alpha d\theta + \frac{1}{2} B\left(\frac{1}{2}, \frac{\alpha+1}{2}\right) \\ &= \frac{1}{2} B\left(\rho^2; \frac{1}{2}, \frac{\alpha+1}{2}\right) + \frac{1}{2} B\left(\frac{1}{2}, \frac{\alpha+1}{2}\right) \end{aligned}$$

and therefore

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq D_1) = \frac{1}{2} + \frac{1}{2} I_{\rho^2} \left( \frac{1}{2}, \frac{\alpha+1}{2} \right).$$

The claim follows by observing that this expression corresponds to the respective Student  $t$ -distribution function, see Equation (9). □

We shall also assume in the following proposition that  $\rho_{12} \geq \rho_{01}\rho_{02}$ , which expresses that the specific components of  $Z_1$  and  $Z_2$  are correlated in a non-negative way. The sole reason for this assumption is to avoid awkward case differentiations, and it can easily be lifted.

**Proposition 3.** Let  $\mathbf{P}(G > \cdot) \in RV_{-\alpha}$ . Then,

$$\begin{aligned} \lim_{C \rightarrow -\infty} \mathbf{P}(Z_1 \leq D_1, Z_2 \leq D_2 | Z_0 \leq C) &= \frac{1}{2} t_{\alpha+1} \left( \frac{\sqrt{(\alpha+1)t}}{\sqrt{1-t^2}} \right) \\ &+ \int_{-\arcsin t}^{\pi/2} \left[ \frac{1}{2} - t_{\alpha+2} \left( - \left| \frac{\sqrt{\alpha+2}}{q_3(\varphi)} \right| \right) \right] (\cos \varphi)^\alpha d\varphi \left( B \left( \frac{1}{2}, \frac{\alpha+1}{2} \right) \right)^{-1} \\ &+ \int_{-\arcsin \rho_{01}}^{-\arcsin t} t_{\alpha+2} \left( - \left| \frac{\sqrt{\alpha+2}}{q_3(\varphi)} \right| \right) (\cos \varphi)^\alpha d\varphi \left( B \left( \frac{1}{2}, \frac{\alpha+1}{2} \right) \right)^{-1}, \end{aligned} \quad (11)$$

$$\text{where } q_3(\varphi) = \frac{\sqrt{1 - \rho_{02}^2 - q_1^2}}{\rho_{02} \cos \varphi + q_1 \sin \varphi} \text{ and } q_1 = \frac{\rho_{12} - \rho_{01}\rho_{02}}{\sqrt{1 - \rho_{01}^2}} \text{ and } t := \rho_{01} \wedge \frac{\rho_{02}}{\sqrt{q_1^2 + \rho_{02}^2}}.$$

*Proof.* Write  $u \in \mathcal{S}_3$  in polar coordinates as  $u = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)^T$ , with  $\theta \in [0, \pi]$ ,  $\varphi \in [-\pi, \pi]$ . Let  $A$  be given by  $A_0 = (1, 0, 0)^T$ ,  $A_1 = (\rho_{01}, \sqrt{1 - \rho_{01}^2}, 0)^T$  and  $A_2 = (\rho_{02}, q_1, q_2)^T$  with  $q_1 = \frac{\rho_{12} - \rho_{01}\rho_{02}}{\sqrt{1 - \rho_{01}^2}}$  and  $q_2 = \sqrt{1 - \rho_{02}^2 - q_1^2}$ . We have  $A_0 \cdot u = \sin \theta \cos \varphi > 0$  if  $\varphi \in (-\pi/2, \pi/2)$  and  $\theta \in (0, \pi)$ . For  $A_1 \cdot u = \rho_{01} \sin \theta \cos \varphi + \sqrt{1 - \rho_{01}^2} \sin \theta \sin \varphi = \rho_{01} \sin \theta \sin(\varphi + \arcsin \rho_{01}) > 0$  we obtain  $\varphi \in (-\arcsin \rho_{01}, \pi - \arcsin \rho_{01})$  on  $\theta \in (0, \pi)$ . Finally, for  $A_2 \cdot u = \sin \theta (\rho_{02} \cos \varphi + q_1 \sin \varphi) + q_2 \cos \theta > 0$  we need to distinguish four cases: First, if  $q_4(\varphi) := \rho_{02} \cos \varphi + q_1 \sin \varphi > 0$  and  $\theta \in (0, \pi/2)$ , then  $A_2 \cdot u > 0$  if  $\tan \theta > -q_2/q_4$ , which is fulfilled for all  $\theta \in (0, \pi)$ . Second, if  $q_4 > 0$  and  $\theta \in (\pi/2, \pi)$ , then  $A_2 \cdot u > 0$  if  $\tan \theta < -q_2/q_4$ , which implies  $\theta < \pi - \arctan(q_2/q_4) = \pi - \arcsin \left( \frac{q_3}{\sqrt{1 + q_3^2}} \right)$ , where  $q_3(\theta) := q_2/q_4$ . Third, for  $q_4 < 0$  and  $\theta \in (0, \pi/2)$ , we have  $A_2 \cdot u > 0$  if  $\tan \theta < -q_2/q_4 = q_2/|q_4|$ , so that  $\theta < \arctan(q_2/|q_4|) = \arcsin \left( \frac{|q_3|}{\sqrt{1 + q_3^2}} \right)$ . Fourth, if  $q_4 < 0$  and  $\theta \in (\pi/2, \pi)$ , then  $A_2 \cdot u > 0$  if  $\tan \theta > q_2/|q_4|$ , but this is not fulfilled for any  $\theta \in (\pi/2, \pi)$ , since  $\tan \theta < 0$ . Finally, we have  $q_4 = \rho_{02} \cos \varphi + q_1 \sin \varphi > 0$  for  $\varphi \in (-\pi/2, \pi/2)$  if  $\tan \varphi > -\rho_{02}/q_1$ , resp.



$\varphi > \arctan(-\rho_{02}/q_1) = \arcsin(-\rho_{02}/\sqrt{q_1^2 + \rho_{02}^2})$ . Putting everything together, we obtain

$$\begin{aligned} \lim_{C \rightarrow -\infty} \mathbf{P}(Z_1 \leq D_1, Z_2 \leq D_2 | Z_0 \leq C) &= \left( \int_{-\pi/2}^{\pi/2} \int_0^\pi (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi \right)^{-1} \\ &\left[ \int_t^{\pi/2} \int_0^{\pi/2} (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi + \int_t^{\pi/2} \int_{\pi/2}^{\pi - \arcsin(q_3/\sqrt{1+q_3^2})} (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi \right. \\ &\quad \left. + \int_{-\arcsin \rho_{01}}^t \int_0^{\arcsin(q_3/\sqrt{1+q_3^2})} (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi \right], \end{aligned}$$

with  $t := -\arcsin(\min(\rho_{01}, \rho_{02}/\sqrt{q_1^2 + \rho_{02}^2}))$ .

First,

$$\int_{-\pi/2}^{\pi/2} \int_0^\pi (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi = B\left(\frac{\alpha+2}{2}, \frac{1}{2}\right) B\left(\frac{1}{2}, \frac{\alpha+1}{2}\right).$$

Second,

$$\begin{aligned} &\int_t^{\pi/2} \int_0^{\pi/2} (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi \\ &= \frac{1}{4} B\left(\frac{\alpha+2}{2}, \frac{1}{2}\right) \left[ B\left(\frac{1}{2}, \frac{\alpha+1}{2}\right) + B\left(\rho_{01}^2 \wedge \frac{\rho_{02}^2}{q_1^2 + \rho_{02}^2}; \frac{1}{2}, \frac{\alpha+1}{2}\right) \right] \end{aligned}$$

and

$$\begin{aligned} &\int_t^{\pi/2} \int_{\pi/2}^{\pi - \arcsin(q_3(\varphi)/\sqrt{1+q_3(\varphi)^2})} (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi \\ &= \int_t^{\pi/2} \frac{1}{2} \left[ B\left(\frac{\alpha+2}{2}, \frac{1}{2}\right) - B\left(\frac{q_3(\varphi)^2}{1+q_3(\varphi)^2}, \frac{\alpha+2}{2}, \frac{1}{2}\right) \right] (\cos \varphi)^\alpha d\varphi \end{aligned}$$

and

$$\begin{aligned} &\int_{-\arcsin \rho_{01}}^t \int_0^{\arcsin(q_3(\varphi)/\sqrt{1+q_3(\varphi)^2})} (\sin \theta)^{\alpha+1} (\cos \varphi)^\alpha d\theta d\varphi \\ &= \int_{-\arcsin \rho_{01}}^t \frac{1}{2} B\left(\frac{q_3(\varphi)^2}{1+q_3(\varphi)^2}, \frac{\alpha+2}{2}, \frac{1}{2}\right) (\cos \varphi)^\alpha d\varphi. \end{aligned}$$




Figure 1: Asymptotic univariate PD's (left) and bivariate PD's (right) as a function of the tail index. Correlations are  $\rho_{01} = \rho_{02} = \rho$  and  $\rho_{12} = \rho^2$ . The choice of  $\rho_{12}$  implies that the asset returns are correlated only via the risk factor.

Putting everything together yields

$$\begin{aligned} \lim_{C \rightarrow -\infty} \mathbf{P}(Z_1 \leq D_1, Z_2 \leq D_2 | Z_0 \leq C) &= \frac{1}{4} \left[ 1 + I_{\rho_{12}^2 \wedge \frac{\rho_{02}^2}{q_1^2 + \rho_{02}^2}} \left( \frac{1}{2}, \frac{\alpha+1}{2} \right) \right] \\ &\quad + \frac{1}{2} \int_t^{\pi/2} \left[ 1 - I_{\frac{q_3(\varphi)^2}{1+q_3(\varphi)^2}} \left( \frac{\alpha+2}{2}, \frac{1}{2} \right) \right] (\cos \varphi)^\alpha d\varphi \left( B \left( \frac{1}{2}, \frac{\alpha+1}{2} \right) \right)^{-1} \\ &\quad + \frac{1}{2} \int_{-\arcsin \rho_{01}}^t I_{\frac{q_3(\varphi)^2}{1+q_3(\varphi)^2}} \left( \frac{\alpha+2}{2}, \frac{1}{2} \right) (\cos \varphi)^\alpha d\varphi \left( B \left( \frac{1}{2}, \frac{\alpha+1}{2} \right) \right)^{-1}, \end{aligned}$$

and replacing the incomplete beta functions by the Student- $t$  distributions, cf. Equation (9), yields the claim.  $\square$

Figure 1 shows examples of asymptotic univariate and bivariate PD's for varying tail index  $\alpha$ . This demonstrates how PD's depend on both the tail index and the correlations.

## 4 Default correlations under stress

In the case where  $\mathbf{P}(G > \cdot) \in RV_{-\alpha}$ , default correlations can be explicitly calculated using the results from Propositions 2 and 3. For the case where  $G$  is rapidly varying, we have the following result.

**Proposition 4.** Let  $\mathbf{P}(G > \cdot) \in RV_{-\infty}$ . Then,

$$\lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq D_1\}}, \mathbf{1}_{\{Z_2 \leq D_2\}}) = 0,$$

where  $\text{Corr}^C$  denotes the correlation under  $\mathbf{P}^C$ .

*Proof.* It is easily seen that for any probability measure  $\mathbf{P}$ ,

$$\text{Corr}(\mathbf{1}_{\{Z_1 \leq D_1\}}, \mathbf{1}_{\{Z_2 \leq D_2\}}) = \frac{\mathbf{P}(Z_1 > D_1, Z_2 > D_2) - (1 - \mathbb{E}(\mathbf{1}_{\{Z_1 \leq D_1\}}))(1 - \mathbb{E}(\mathbf{1}_{\{Z_2 \leq D_2\}}))}{\sqrt{\mathbb{E}(\mathbf{1}_{\{Z_1 \leq D_1\}})(1 - \mathbb{E}(\mathbf{1}_{\{Z_1 \leq D_1\}}))} \sqrt{\mathbb{E}(\mathbf{1}_{\{Z_2 \leq D_2\}})(1 - \mathbb{E}(\mathbf{1}_{\{Z_2 \leq D_2\}}))}}. \quad (12)$$

Using Theorem 1, this simplifies in the case of  $\mathbf{P}(G > \cdot) \in RV_{-\infty}$  to

$$\begin{aligned} \lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq D_1\}}, \mathbf{1}_{\{Z_2 \leq D_2\}}) &= \lim_{C \rightarrow -\infty} \frac{\mathbf{P}(Z_1 > D_1, Z_2 > D_2 | Z_0 \leq C)}{\sqrt{\mathbf{P}(Z_1 > D_1 | Z_0 \leq C)} \sqrt{\mathbf{P}(Z_2 > D_2 | Z_0 \leq C)}} \\ &= \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_1 \leq D_1, Z_2 \leq D_2, Z_0 > C)}{\sqrt{\mathbf{P}(Z_1 \leq D_1, Z_0 > C)} \sqrt{\mathbf{P}(Z_2 \leq D_2, Z_0 > C)}}, \end{aligned} \quad (13)$$

where the second equality follows from the symmetry of the  $Z_i$ .

Suppose first that  $\rho_{01} \neq \rho_{02}$ ; wlog. let  $\rho_{02} > \rho_{01}$ . As before, we first prove the claim for  $D_1 = D_2 = 0$ .

We write (13) in the form

$$\lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq 0\}}, \mathbf{1}_{\{Z_2 \leq 0\}}) = \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_1 \leq 0, Z_2 \leq 0, Z_0 > C)}{\mathbf{P}(Z_2 \leq 0, Z_0 > C)} \sqrt{\frac{\mathbf{P}(Z_2 \leq 0, Z_0 > C)}{\mathbf{P}(Z_1 \leq 0, Z_0 > C)}}. \quad (14)$$

The first term is bounded by 1. For  $C > 0$ , write the second term in the form

$$\begin{aligned} \mathbf{P}(Z_i \leq 0, Z_0 > C) &= \mathbf{P}\left(G > \frac{C}{A_0 U}, A_0 U > 0, A_i U < 0\right) \\ &= \int_{u \in \mathcal{S}_2, A_0 u > 0, A_i u < 0} \mathbf{P}\left(G > \frac{C}{A_0 u}\right) F_U(du) \\ &= \int_{-\pi/2}^{-\arcsin \rho_{0i}} \mathbf{P}\left(G > \frac{C}{\cos \theta}\right) d\theta. \end{aligned}$$

Since  $\rho_{02} > \rho_{01}$ , we have  $-\arcsin \rho_{02} < -\arcsin \rho_{01}$ , and

$$\sup_{\theta \in (-\pi/2, -\arcsin \rho_{02}]} \cos \theta = \cos(-\arcsin \rho_{02}) = \sqrt{1 - \rho_{02}^2}.$$

Hence, using that  $G$  is rapidly varying and via Dominated Convergence,

$$\lim_{C \rightarrow \infty} \int_{-\pi/2}^{-\arcsin \rho_{02}} \frac{\mathbf{P}(G > C / \cos \theta)}{\mathbf{P}(G > C / \sqrt{1 - \rho_{02}^2})} d\theta = 0,$$

whereas

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}(G > C / \cos \theta)}{\mathbf{P}(G > C / \sqrt{1 - \rho_{02}^2})} = \infty, \quad \text{for } \theta \in (-\arcsin \rho_{02}, -\arcsin \rho_{01}],$$

which implies that

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_2 \leq 0, Z_0 > C)}{\mathbf{P}(Z_1 \leq 0, Z_0 > C)} = 0 \tag{15}$$

and therefore Equation (14) is 0.

To complete the proof for  $\rho_{02} > \rho_{01}$  it remains to show that

$$\lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq D_1\}}, \mathbf{1}_{\{Z_2 \leq D_2\}}) = \lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq 0\}}, \mathbf{1}_{\{Z_2 \leq 0\}}) = 0$$

for arbitrary  $D_1, D_2 \in \mathbb{R}$ . For  $a > 0$  and  $C > |D_2|/a$ ,

$$\mathbf{P}(Z_2 + aZ_0 \leq 0, Z_0 > C) \leq \mathbf{P}(Z_2 \leq D_2, Z_0 > C) \leq \mathbf{P}(Z_2 - aZ_0 \leq 0, Z_0 > C).$$

For sufficiently small  $a$  we obtain from (15)

$$\begin{aligned} 0 &= \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_2 + aZ_0 \leq 0, Z_0 > C)}{\mathbf{P}(Z_1 \leq 0, Z_0 > C)} \leq \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_2 \leq D_2, Z_0 > C)}{\mathbf{P}(Z_1 \leq 0, Z_0 > C)} \\ &\leq \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_2 - aZ_0 \leq 0, Z_0 > C)}{\mathbf{P}(Z_1 \leq 0, Z_0 > C)} = 0 \end{aligned}$$

and therefore

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_2 \leq D_2, Z_0 > C)}{\mathbf{P}(Z_1 \leq 0, Z_0 > C)} = 0.$$

Applying the same argument to the numerator yields

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_2 \leq D_2, Z_0 > C)}{\mathbf{P}(Z_1 \leq D_1, Z_0 > C)} = 0.$$

Obviously,  $\frac{\mathbf{P}(Z_1 \leq D_1, Z_2 \leq D_2 | Z_0 > C)}{\mathbf{P}(Z_2 \leq D_2 | Z_0 > C)}$  is bounded by 1. Hence, by Equality (14),

$$\lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq D_1\}}, \mathbf{1}_{\{Z_2 \leq D_2\}}) = 0.$$

It remains to show that the claim also holds for  $\rho_{01} = \rho_{02}$ . Wlog. assume that  $D_1 \leq D_2$ . We now provide a proof for the case  $D_2 \leq 0$ . The case  $D_2 > 0$  is shown analogously.

It follows from (13) that

$$\lim_{C \rightarrow -\infty} \text{Corr}^C(\mathbf{1}_{\{Z_1 \leq D_1\}}, \mathbf{1}_{\{Z_2 \leq D_2\}}) \leq \lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_1 \leq 0, Z_2 \leq 0, Z_0 > C)}{\mathbf{P}(Z_1 \leq D_1, Z_0 > C)}.$$

The numerator can be written in the form

$$\mathbf{P}(Z_1 \leq 0, Z_2 \leq 0, Z_0 > C) = \int_{A_0.u > 0, A_1.u < 0, i=1,2} \mathbf{P}\left(G > \frac{C}{A_0.u}\right) F_U(du).$$

Express  $u \in \mathcal{S}_3$  in polar coordinates by  $u = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)^T$  with  $\theta \in [0, \pi]$ ,  $\varphi \in [-\pi, \pi]$ . Observe that the conditions  $A_0.u = \sin \theta \cos \varphi > 0$  and  $A_1.u = \rho_{01} \sin \theta \sin(\varphi + \arcsin \rho_{01}) < 0$  imply  $\theta \in (0, \pi)$  and  $\varphi \in (-\pi/2, -\arcsin \rho_{01})$ . Hence,

$$\begin{aligned} x &:= \sup_{A_0.u > 0, A_1.u < 0} A_0.u \\ &= \sup_{\theta \in (0, \pi), \varphi \in (-\pi/2, -\arcsin \rho_{01})} \sin \theta \cos \varphi \\ &= \sin(\pi/2) \cos(-\arcsin \rho_{01}) \\ &= \cos(-\arcsin \rho_{01}) = \sqrt{1 - \rho_{01}^2}. \end{aligned}$$

At  $\theta = \pi/2$ ,  $\varphi = -\arcsin \rho_{01}$ , the condition  $A_2.u < 0$  is not satisfied: since  $A_2.u = \sin \theta (\rho_{02} \cos \varphi + q_1 \sin \varphi) + q_2 \cos \theta$ , where  $q_1$  and  $q_2$  are as in Proposition 3, resp. the proof of Proposition 3 it follows from  $\rho_{02} = \rho_{01}$  that

$$A_2.u = \rho_{01} \underbrace{\cos(-\arcsin \rho_{01})}_{=\sqrt{1-\rho_{01}^2}} - \frac{\rho_{12} - \rho_{01}^2}{\sqrt{1-\rho_{01}^2}} \rho_{01} = \frac{\rho_{01}}{\sqrt{1-\rho_{01}^2}} (1 - \rho_{12}) \geq 0.$$

This implies that

$$y := \sup_{A_0.u > 0, A_1.u < 0, A_2.u < 0} A_0.u < x.$$

Hence, by the property that  $G$  is rapidly varying and via Dominated Convergence,

$$\lim_{C \rightarrow \infty} \int_{u \in \mathcal{S}_3, A_0.u > 0, A_1.u < 0, A_2.u < 0} \frac{\mathbf{P}(G > C/(A_1.u))}{\mathbf{P}(G > C/y)} F_U(du) = 0. \quad (16)$$

For the denominator we have

$$\begin{aligned} \mathbf{P}(Z_1 \leq D_1, Z_0 > C) &= \int_{u \in \mathcal{S}_3, A_0.u > 0, A_1.u < 0} \mathbf{P}\left(G > \max\left(\frac{|D_1|}{|A_1.u|}, \frac{C}{A_0.u}\right)\right) F_U(du) \\ &= \int_0^\pi \int_{-\pi/2}^{-\arcsin \rho_{01}} \mathbf{P}\left(G > \max\left(f(\theta, \varphi), \frac{C}{\sin \theta \cos \varphi}\right)\right) \sin \theta d\varphi d\theta, \end{aligned} \quad (17)$$

where

$$f(\theta, \varphi) := \frac{|D_1|}{|\rho_{01} \sin \theta \sin(\varphi + \arcsin \rho_{01})|}.$$

Obviously, the maximum in the integrand of (17) is given by  $f(\theta, \varphi)$  if

$$\frac{\sin \theta \cos \varphi}{|\rho_{01} \sin \theta \sin(\varphi + \arcsin \rho_{01})|} = \frac{\cos \varphi}{|\rho_{01} \sin(\varphi + \arcsin \rho_{01})|} \geq \frac{C}{|D_1|}.$$

Hence,  $f(\theta, \varphi)$  is the maximum in a neighbourhood of  $\varphi = -\arcsin \rho_{01}$ . On the other hand, for every  $\varphi \in (-\pi/2, -\arcsin \rho_{01})$  the maximum is given by  $C/(A_0.u)$  for  $C$  sufficiently large.

Hence, we choose  $C$  large enough so that

$$\max \left( \frac{|D_1|}{|\rho_{01} \sin \theta \sin(\varphi + \arcsin \rho_{01})|}, \frac{C}{A_{0,u}} \right) = \frac{C}{A_{0,u}} \quad \text{and} \quad A_{0,u} = \sin \theta \cos \varphi > y$$

for  $(\theta, \varphi) \in M$ , where  $M \subset (0, \pi) \times (-\pi/2, -\arcsin \rho_{01})$  is a set of positive Lebesgue measure.

Hence, by the property that  $G$  is rapidly varying and via Dominated Convergence, we obtain from (17) that

$$\lim_{C \rightarrow \infty} \frac{\mathbf{P}(Z_1 \leq D_1, Z_0 > C)}{\mathbf{P}(G > C/y)} \geq \lim_{C \rightarrow \infty} \int_M \frac{\mathbf{P}(G > C/(\sin \theta \cos \varphi))}{\mathbf{P}(G > C/y)} \sin \theta \, d\varphi \, d\theta = \infty.$$

Together with (16), this proves the claim.  $\square$

That default correlations under stress converge to zero in light-tailed models can be explained as follows: In regression analysis, correlation – expressed as  $R^2$  – measures the degree of the linear relationship between two random variables. In the case of default correlations, there are only four possible scenarios: both variables are zero, both variables are one, and exactly one variable is one and the other is zero (and vice versa). In the light-tailed case, since, asymptotically, default is a sure event, only the event that both variables take value one remains. However, for any large stress level, the probability that both variables are zero vanishes most quickly and probability mass is pushed into the remaining three cases. No line will succeed in adequately describing the relationship of those variables and in particular will not capture the variance in the centralized variables.

## 5 Implications for credit portfolio modelling

The main results of this paper are formulae for asymptotic stressed default probabilities in credit portfolio models with elliptically distributed risk factors and asset variables  $Z \stackrel{\mathcal{L}}{=} GAU$ .

We have shown that for any  $D_i \in \mathbb{R}$

$$\lim_{C \rightarrow -\infty} \mathbf{P}(Z_i \leq D_i | Z_0 \leq C) = 1,$$

if  $\mathbf{P}(G > \cdot)$  is rapidly varying and, if  $\mathbf{P}(G > \cdot)$  is regularly varying,

$$\lim_{C \rightarrow -\infty} \mathbf{P}(Z_i \leq D_i | Z_0 \leq C) = t_{\alpha+1} \left( \frac{\sqrt{\alpha+1} \rho}{\sqrt{1-\rho^2}} \right) \in [1/2, 1).$$

This behaviour of limiting default probabilities is fundamentally different to tail dependence, which is a popular measure in finance to assess the ability of a bivariate distribution to generate joint extreme events: For two random variables  $Y_1$  and  $Y_2$  with distribution functions  $F_1$  and  $F_2$ , the coefficient of (lower) tail dependence of  $Y_1$  and  $Y_2$  is

$$\lambda_l(Y_1, Y_2) := \lim_{q \rightarrow 0^+} \mathbf{P}(Y_2 \leq F_2^{\leftarrow}(q) | Y_1 \leq F_1^{\leftarrow}(q)), \quad (18)$$

where  $F_i^{\leftarrow}$  denotes the inverse of the df  $F_i$ . The tail dependence coefficient depends only on the copula rather than the bivariate distribution function, see e.g. Joe (1997); Nelsen (1999); McNeil *et al.* (2005). For heavy-tailed elliptical distributions, i.e., the mixing variable  $G$  is in  $RV_{-\alpha}$ , the tail dependence is given by

$$\lambda_l(Y_1, Y_2) = \frac{\int_{\pi/2 - \arcsin \rho}^{\pi/2} (\cos \theta)^\alpha d\theta}{\int_0^{\pi/2} (\cos \theta)^\alpha d\theta} = 2t_{\alpha+1} \left( -\sqrt{\frac{(\alpha+1)(1-\rho)}{1+\rho}} \right),$$

where  $\rho := \text{Corr}(Y_1, Y_2)$ , see Hult and Lindskog (2002); Schmidt (2002); McNeil *et al.* (2005). Hence,  $\lambda_l(Y_1, Y_2) > 0$  for  $\rho > -1$ . In contrast, the tail dependence is zero for a normal distribution, which is the most frequently used distribution in structural credit portfolio models, e.g. normal distributions are used in Moody's KMV model (Crosbie and Bohn (2002)). Due to zero tail dependence, normally distributed models are usually considered less sensitive to extreme stress than heavy-tailed models. The results in this paper show that this is not necessarily the case: in the limit, the impact of stress on default probabilities is higher in light-tailed models than in heavy-tailed models.

To analyze the precise difference between tail dependence and asymptotic stressed PD's,






Figure 2: Tail dependence coefficient and asymptotic PD under stress as a function of the tail index  $\alpha$ . Left: correlation parameter 0.4; right: correlation parameter 0.7.

we write the tail dependence of  $Z_0$  and  $Z_1$  in the form

$$\lambda_t(Z_0, Z_1) = \lim_{C \rightarrow -\infty} \mathbf{P}(Z_1 \leq C | Z_0 \leq C), \quad (19)$$

which is equivalent to definition (18) since  $Z_0$  and  $Z_1$  are identically distributed. Hence, for calculating tail dependence the conditional probability  $\mathbf{P}^C(Z_1 \leq C)$  has to be evaluated whereas the stressed default probability  $\mathbf{P}^C(Z_1 \leq D)$  is evaluated at a constant  $D$ . For light-tailed elliptical distributions, the variable  $Z_1$  only attains limit values in the range

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(C \leq Z_1 \leq D) = 1$$

for any  $D \in \mathbb{R}$ , whereas extreme events outside this range have positive probability in the heavy-tailed case:

$$\lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 \leq C) > 0, \quad \lim_{C \rightarrow -\infty} \mathbf{P}^C(Z_1 > D) > 0.$$

The resulting difference between tails dependence and asymptotic stressed PD's as a function of the tail index  $\alpha$  is shown in Figure 2.

Turning now to stressed default correlations, we observe a behaviour similar to tail dependence: we have shown that stressed default correlations converge to 0 in the light-tailed case and to a positive number in the heavy-tailed case. Hence, in light-tailed models extreme stress scenarios tend to heavily increase the expected loss whereas tail risk measures, which are driven




Figure 3: PD's under stress as a function of the stress probability  $p := F(C)$ , with  $F$  the distribution function of the respective model and  $C$  the stress level. Models considered are the normal distribution and the  $t$ -distribution with parameter 3. Correlations are 0.4. Left: unconditional PD is 0.1 (i.e.,  $D = F^{(-1)}(0.1)$ ); right: unconditional PD is 0.01.

by the dependence of default events, are less affected.

It is important to note, however, that the asymptotic behaviour analysed in this paper is not necessarily representative for typical stress scenarios in credit risk management. To gain further insight and provide a heuristic answer, we consider PD's under stress for various stress levels and compare them in light- and heavy-tailed models. Figure 3 shows PD's under stress for both normally distributed and  $t$ -distributed ( $\nu = 3$ ) models as a function of the stress probability  $p := F(C)$ , where  $F$  is the distribution function of the respective model and  $C$  is the stress level. The correlation is chosen to be 0.4. Despite converging to a value smaller than 1, PD's under stress in the  $t$ -distributed model dominate the normally distributed case unless the stress probability is very small: If the unconditional PD is 10%, then for stress probabilities greater than  $10^{-6}$ , the PD under stress in the  $t$ -distributed model is greater than the respective PD in the normal model. If the unconditional PD is 1%, then the threshold lies beyond  $10^{-13}$ .

Hence, aside from providing useful information for stress testing, our results indicate that gauging the suitability of a distribution family for credit portfolio modelling solely on asymptotic behaviour may be misleading.

## References

- N. Bingham and R. Kiesel. Semi-parametric modelling in finance: theoretical foundations. *Quantitative Finance*, 2(4):241–250, 2002.
- N. Bingham, C. Goldie, and J. Teugels. *Regular Variation*. Cambridge University Press, Cambridge, MA, 1987.
- BIS. Principles for sound stress testing practices and supervision. Basel Committee on Banking Supervision, Bank for International Settlements, January 2009.
- G. Bonti, M. Kalkbrener, C. Lotz, and G. Stahl. Credit risk concentrations under stress. *Journal of Credit Risk*, 2(3):115–136, 2006.
- M. Brunnermeier, A. Crockett, C. Goodhart, A. Persaud, and H. Shin. The fundamental principles of financial regulation. *Geneva Reports on the World Economy*, 11, 2009.
- S. Cambanis, S. Huang, and G. Simons. On the theory of elliptically contoured distributions. *Journal of Multivariate Analysis*, 11(3):368–385, 1981.
- B. Crosbie and J. Bohn. Modelling default risk. Working paper, KMV Corporation, 2002. Available from [www.kmv.com](http://www.kmv.com).
- L. de Haan and A. Ferreira. *Extreme Value Theory: An Introduction*. Springer, New York, 2006.
- K. Duellmann and M. Erdelmeier. Crash testing German banks. *International Journal of Central Banking*, 5(3):139–175, 2009.
- P. Embrechts, D. D. Lambrigger, and M. V. Wüthrich. Multivariate extremes and the aggregation of dependent risks: examples and counter-examples. *Extremes*, 12(2):107–127, 2009.
- K.-T. Fang, S. Kotz, and K.-W. Ng. *Symmetric Multivariate and Related Distributions*. Chapman and Hall, London, 1990.

- H. Hult and F. Lindskog. Multivariate extremes, aggregation and dependence in elliptical distributions. *Advances in Applied Probability*, 34(3):587–608, 2002.
- H. Joe. *Multivariate Models and Dependence Concepts*. Chapman & Hall/CRC, 1997.
- M. Kalkbrener and N. Packham. Correlation under stress in normal variance mixture models. *Mathematical Finance*, forthcoming, 2013.
- C. Klüppelberg, G. Kuhn, and L. Peng. Semi-parametric models for the multivariate tail dependence function – the asymptotically dependent case. *Scandinavian Journal of Statistics*, 35(4):701–718, 2008.
- G. Knuth. The Art of Computer Programming, vol. 2: Seminumerical Algorithms. *Reading, Massachusetts*, 1998.
- J. Larosière et al. Report. The High-Level Group on Financial Supervision in the EU, February 2009.
- G. Mainik and P. Embrechts. Diversification in heavy-tailed portfolios: properties and pitfalls. *Annals of Actuarial Science*, 7(01):26–45, 2013.
- A. McNeil, R. Frey, and P. Embrechts. *Quantitative Risk Management*. Princeton University Press, Princeton, NJ, 2005.
- R. C. Merton. On the pricing of corporate debt: The risk structure of interest rates. *The Journal of Finance*, 29(2):449–470, May 1974.
- R. B. Nelsen. *An Introduction to Copulas*. Springer, 1999.
- R. Schmidt. Tail dependence for elliptically contoured distributions. *Mathematical Methods of Operations Research*, 55(2):301–327, 2002.
- A. Turner. The Turner Review — A Regulatory Response to the Global Banking Crisis. *FSA*, March, 2009.

No.	Author/Title	
210.	Buschmann, Christian / Heidorn, Thomas The Liquidity Reserve Funding and Management Strategies	2014
209.	Heidorn, Thomas / Mokinski, Frieder / Rühl, Christoph / Schmaltz, Christian The Impact of Fundamental and Financial Traders on the Term Structure of Oil	2014
208.	Krasnosselski, Nikolai / Cremers, Heinz / Sanddorf-Köhle, Walter Messung des Marktrisikos mit generalisierter autoregressiver bedingter heteroskedastischer Modellierung der Volatilität	2014
207.	Winkler, Adalbert The Lender of Last Resort in Court	2014
206.	Winkler, Adalbert Der lender of last resort vor Gericht	2013
205.	Lang, Michael The Early Warnings of Balance-of-Payments problems: Kaminsky and Reinhart Revisited	2013
204.	Odermann, Alexander / Cremers, Heinz Komponenten und Determinanten des Credit Spreads – Empirische Untersuchung während Phasen von Marktstress	2013
203.	Vogel, Heinz-Dieter / Bannier, Christina E. / Heidorn, Thomas Functions and characteristics of corporate and sovereign CDS	2013
202.	Loechel, Horst / Packham, Natalie / Walisch, Fabian Determinants of the onshore and offshore Chinese Government yield curves	2013
201.	Kostka, Genia China's evolving green planning system: Are targets the answer?	2013
200.	Moormann, Jürgen / Grau, Corinna Exploring the interrelation between process management and organiza-tional culture: A critical review	2013
199.	Libman, Alexander / Obydenkova, Anastassia Communism or Communists? Soviet Legacies and Corruption in Transition Economies	2013
198.	Völker, Florian / Cremers, Heinz / Panzer, Christof Integration des Marktliquiditätsrisikos in das Risikoanalysekonzept des Value at Risk	2012
197.	Inklaar, Robert / Koetter, Michael / Noth, Felix Who's afraid of big bad banks? – Bank competition, SME and industry growth	2012
196.	Philipp Boeing / Elisabeth Mueller / Philipp Sandner What Makes Chinese Firms Productive? Learning from Indigenous and Foreign Sources of Knowledge	2012
195.	Krones, Julia / Cremers, Heinz Eine Analyse des Credit Spreads und seiner Komponenten als Grundlage für Hedge Strategien mit Kreditderivaten	2012
194.	Herrmann-Pillath, Carsten Performativity of Economic Systems: Approach and Implications for Taxonomy	2012
193.	Boldyrev, Ivan A. / Herrmann-Pillath, Carsten Moral Sentiments, Institutions, and Civil Society: Exploiting Family Resemblances between Smith and Hegel to Resolve Some Conceptual Issues in Sen's Recent Contributions to the Theory of Justice	2012
192.	Mehmke, Fabian / Cremers, Heinz / Packham, Natalie Validierung von Konzepten zur Messung des Marktrisikos - insbesondere des Value at Risk und des Expected Shortfall	2012
191.	Tinschert, Jonas / Cremers, Heinz Fixed Income Strategies for Trading and for Asset Management	2012

Printed edition: € 25.00 + € 2.50 shipping

Download working paper:

<http://www.frankfurt-school.de/content/de/research/workingpapers.html>

**Order address / contact**

Frankfurt School of Finance & Management

Sonnemannstr. 9 – 11 ■ 60314 Frankfurt/M. ■ Germany

Phone: +49 (0) 69 154 008 – 830 ■ Fax: +49 (0) 69 154 008 –4830

eMail: [r.ewel@fs.de](mailto:r.ewel@fs.de)

Further information about Frankfurt School of Finance & Management  
may be obtained at: <http://www.fs.de>