

Giacomini, Enzo; Härdle, Wolfgang Karl

Working Paper

Value-at-risk calculations with time varying copulae

SFB 649 Discussion Paper, No. 2005-004

Provided in Cooperation with:

Collaborative Research Center 649: Economic Risk, Humboldt University Berlin

Suggested Citation: Giacomini, Enzo; Härdle, Wolfgang Karl (2005) : Value-at-risk calculations with time varying copulae, SFB 649 Discussion Paper, No. 2005-004, Humboldt University of Berlin, Collaborative Research Center 649 - Economic Risk, Berlin

This Version is available at:

<https://hdl.handle.net/10419/25023>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

Value-at-Risk Calculations with Time Varying Copulae

Enzo Giacomini*
Wolfgang Härdle*



* CASE - Center for Applied Statistics and Economics,
Humboldt-Universität zu Berlin, Germany

This research was supported by the Deutsche
Forschungsgemeinschaft through the SFB 649 "Economic Risk".

<http://sfb649.wiwi.hu-berlin.de>
ISSN 1860-5664

SFB 649, Humboldt-Universität zu Berlin
Spandauer Straße 1, D-10178 Berlin



Value-at-Risk Calculations with Time Varying Copulae

Enzo Giacomini, Wolfgang Härdle
Center for Applied Statistics and Economics
Institut für Statistik und Ökonometrie
Humboldt-Universität zu Berlin
Spandauer Str. 1, D-10178 Berlin
Germany
E-mail: giacomini@wiwi.hu-berlin.de

Value-at-Risk (VaR) of a portfolio is determined by the multivariate distribution of the risk factors increments. This distribution can be modelled through copulae, where the copulae parameters are not necessarily constant over time. For an exchange rate portfolio, copulae with time varying parameters are estimated and the VaR simulated accordingly. Backtesting underlines the improved performance of time varying copulae.

Value-at-Risk and Copulae

At time t a linear portfolio composed of d positions $w = (w_1, \dots, w_d)^\top$ on assets with prices $S_t = (S_{1,t}, \dots, S_{d,t})^\top$ and log prices $Z_t = \ln S_t$, has value

$$(1) \quad V_t = \sum_{j=1}^d w_j e^{Z_{j,t}}$$

The *profit and loss* (P&L) function is defined as $L_{t+1} = (V_{t+1} - V_t)$. Defining $X_{t+1} = (Z_{t+1} - Z_t)$ as the time increment in the risk factors from period t to $t + 1$, the P&L can be expressed as:

$$(2) \quad L_{t+1} = \sum_{j=1}^d w_j S_{j,t} (e^{X_{j,t+1}} - 1)$$

The *Value-at-Risk* (VaR) is calculated as the α -quantile from F_L , the distribution of L :

$$(3) \quad VaR = F_L^{-1}(\alpha)$$

The 1-dimensional distribution F_L depends on the d -dimensional distribution F_X . Using *copulae*, the marginal distributions F_{X_j} from each univariate increment can be *separately* modelled from their dependence structure and then *coupled* together to form the multivariate distribution F_X .

In the following, the dependence parameter $\hat{\theta}$ and the joint distribution $\hat{F}_X$ from a sample $\{X_t\}_{t=1}^T$ of log returns from exchange rate positions are estimated with copulae. A Monte Carlo simulation based on $\hat{F}_X$ generates different P&L samples. The quantiles at different levels from these simulation samples are then used as estimators for the Value-at-Risk of differem portfolio.

Computing Value-at-Risk with Copulae

A copula is a d -dimensional distribution function $C : [0, 1]^d \rightarrow [0, 1]$ with uniform marginals on the interval $[0, 1]$. As in Nelsen (1998), multivariate distributions can be modelled via:

Theorem 1 (Sklar's theorem) *Let F be a d -dimensional distribution function with marginals $F_{X_1}, \dots, F_{X_d}$. Then there exists a copula C with*

$$(4) \quad F(x_1, \dots, x_d) = C\{F_{X_1}(x_1), \dots, F_{X_d}(x_d)\}$$

for every $x_1, \dots, x_d \in \overline{\mathbb{R}}$. If $F_{X_1}, \dots, F_{X_d}$ are continuous, then C is unique. On the other hand, if C is a copula and $F_{X_1}, \dots, F_{X_d}$ are distribution functions, then the function F defined in (4) is a joint distribution function with marginals $F_{X_1}, \dots, F_{X_d}$.

The estimation of the Value-at-Risk, based on an *i.i.d.* sample $\{X_t\}_{t=1}^T$ is implemented in the following procedure:

1. specification of marginal distributions $F_{X_j}(x_j)$
2. specification of copula $C(u_1, \dots, u_d; \theta)$
3. fitting the copula C to obtain $\hat{\theta}$
4. generation of Monte Carlo data $X_{T+1} \sim C(u_1, \dots, u_d; \hat{\theta})$
5. generation of a sample of portfolio losses $L_{T+1}(X_{T+1})$
6. estimation of $\widehat{VaR}_{T+1}$, the empirical quantile at level α from $L_{T+1}(X_{T+1})$.

For copulae belonging to a parametric family $C = \{C_\theta, \theta \in \Theta\}$ and univariate marginals $F_{X_j}(x_j; \delta_j)$, the density of X is given by:

$$f(x_1, \dots, x_d; \delta_1, \dots, \delta_d, \theta) = c\{F_{X_1}(x_1; \delta_1), \dots, F_{X_d}(x_d; \delta_d); \theta\} \prod_{j=1}^d f_j(x_j; \delta_j)$$

where

$$c(u_1, \dots, u_d) = \frac{\partial^d C(u_1, \dots, u_d)}{\partial u_1 \dots \partial u_d}$$

In the IFM (*inference for margins*) method, the log-likelihood function for each of the marginal distributions

$$(5) \quad \ell_j(\delta_j) = \sum_{t=1}^T \ln f_j(x_{j,t}; \delta_j), j = 1, \dots, d$$

is maximized to obtain estimates $(\hat{\delta}_1, \dots, \hat{\delta}_d)^\top$. The function

$$(6) \quad \ell(\theta, \hat{\delta}_1, \dots, \hat{\delta}_d) = \sum_{t=1}^T [\ln c\{F_{X_1}(x_{1,t}; \hat{\delta}_1), \dots, F_{X_d}(x_{d,t}; \hat{\delta}_d); \theta\}]$$

is then maximized over θ to get the dependence parameter estimate $\hat{\theta}$. The estimates $\hat{\theta}_{IFM} = (\hat{\delta}_1, \dots, \hat{\delta}_d, \hat{\theta})^\top$ solve

$$(\partial \ell_1 / \partial \delta_1, \dots, \partial \ell_d / \partial \delta_d, \partial \ell / \partial \theta) = 0$$

Backtesting

This procedure is applied to a daily exchange rate portfolio (DEM/USD and GBP/USD from 01.12.1979 to 01.04.1994) with $T = 250$. The univariate risk factor increments (log returns) are assumed to be Gaussian distributed with parameters estimated from the data. The selected copulae belong to the bivariate one-parametric Gumbel family:

$$(7) \quad C(u, v) = \exp(-[(\ln u)^\theta + (\ln v)^\theta]^{\theta^{-1}}), 1 \leq \theta \leq \infty$$

To evaluate the performance of the copula in the VaR calculations, different portfolio compositions are used to generate P&L samples. The quantiles from the samples at four levels $\alpha_1 = 0.05$, $\alpha_2 = 0.01$, $\alpha_3 = 0.005$ and $\alpha_4 = 0.001$ are used as estimators for VaR.

The estimated VaR is compared with the realization of the P&L function, an *exceedance* occurring for each P&L value smaller than the estimated VaR. The ratio of the number of exceedances to the number of observations gives the empirical level $\hat{\alpha}$.




Figure 1: Value-at-Risk at levels $\alpha_1 = 0.05$ (yellow), $\alpha_2 = 0.01$ (green), $\alpha_3 = 0.005$ (red), and $\alpha_4 = 0.001$ (blue), P&L (black), estimated at each time from a Monte Carlo sample of 10.000 P&L values generated with Gumbel copula, $w = (2, 1)^\top$.




Figure 2: Value-at-Risk at level $\alpha = 0.05$ (yellow), P&L (black) and exceedances (red), $\hat{\alpha} = 0.0573$, $w = (2, 1)^\top$. P&L samples generated with Gumbel copula.

Table 1: Gumbel copula, empirical levels $\hat{\alpha}$ for different portfolios.

Portfolio $w^\top$	level $\alpha(\times 10^2)$			
	5	1	0.5	0.1
	empirical level $\hat{\alpha}(\times 10^2)$			
(1, 1)	6.05	2.45	1.75	0.83
(1, 2)	6.34	2.74	1.75	1.00
(2, 1)	5.73	2.24	1.58	0.69
(2, 3)	6.22	2.56	1.75	0.92
(3, 2)	5.99	2.30	1.55	0.74
(-1, 2)	1.64	0.37	0.20	0.11
(1, -2)	2.01	0.51	0.43	0.11
(-2, 1)	4.44	1.49	0.95	0.40
(2, -1)	4.09	1.35	1.09	0.49

REFERENCES

- Durrelman, V., Nikeghbali, A. and Roncalli, T. (2000). Which Copula is the Right One ?, *Groupe de Recherche Opérationnelle Crédit Lyonnais*.
- Embrechts, P., McNeil, A. and Straumann, D. (1999). *Correlation and Dependence in Risk Management: Properties and Pitfalls Correlation, Risk Management: Value at Risk and Beyond*, Cambridge University Press, Cambridge.
- Franke, J., Härdle, W. and Hafner, C. (2004). *Statistics of Financial Markets*, Springer-Verlag, Heidelberg.
- Giacomini, E. and Härdle, W. (2005). Nonparametric Risk Management with Adaptive Copulae, *Discussion Paper, SFB 649*.
- Härdle, W., Herwartz, H. and Spokoiny, V. (2003). Time Inhomogeneous Multiple Volatility Modeling, *Journal of Financial Econometrics*, 2003 1 (1): 55-95.
- Härdle, W., Kleinow, T. and Stahl, G. (2002). *Applied Quantitative Finance*, Springer-Verlag, Heidelberg.
- Joe, H. (1997). *Multivariate Models and Dependence Concepts*, Chapman & Hall, London.
- Nelsen, R. (1998). *An Introduction to Copulas*, Springer-Verlag, New York.

ACKNOWLEDGEMENT

Financial support of *Deutsche Forschungsgemeinschaft* via *SFB 649 "Ökonomisches Risiko"*, Humboldt-Universität zu Berlin, is gratefully acknowledged.

RÉSUMÉ

La Value at Risk (VaR) d'un portefeuille est déterminée par la distribution multivariée des incréments des facteurs de risques. Cette distribution peut être modélisée par des copules dont les paramètres ne sont pas nécessairement constants par rapport au temps. Pour un portefeuille de taux de change, des copules dépendant du temps sont estimés et la VaR est ainsi simulée. Le backtesting confirme l'amélioration apportée par les copules dépendant du temps.

SFB 649 Discussion Paper Series

For a complete list of Discussion Papers published by the SFB 649, please visit <http://sfb649.wiwi.hu-berlin.de>.

- 001 "Nonparametric Risk Management with Generalized Hyperbolic Distributions" by Ying Chen, Wolfgang Härdle and Seok-Oh Jeong, January 2005.
- 002 "Selecting Comparables for the Valuation of the European Firms" by Ingolf Dittmann and Christian Weiner, February 2005.
- 003 "Competitive Risk Sharing Contracts with One-sided Commitment" by Dirk Krueger and Harald Uhlig, February 2005.
- 004 "Value-at-Risk Calculations with Time Varying Copulae" by Enzo Giacomini and Wolfgang Härdle, February 2005.

SFB 649, Spandauer Straße 1, D-10178 Berlin
<http://sfb649.wiwi.hu-berlin.de>

This research was supported by the Deutsche
Forschungsgemeinschaft through the SFB 649 "Economic Risk".

