ECONSTOR
Make Your Publications Visible.

A Service of
ZBW
Leibniz-Informationszentrum Wirtschaft
Leibniz Information Centre for Economics

Klein, Ingo

Working Paper

Van Zwet ordering for Fechner asymmetry

IWQW Discussion Papers, No. 08/2011

Provided in Cooperation with:
Friedrich-Alexander University Erlangen-Nuremberg, Institute for Economics

Suggested Citation: Klein, Ingo (2011) : Van Zwet ordering for Fechner asymmetry, IWQW Discussion Papers, No. 08/2011, Friedrich-Alexander-Universität Erlangen-Nürnberg, Institut für Wirtschaftspolitik und Quantitative Wirtschaftsforschung (IWQW), Nürnberg

This Version is available at:
https://hdl.handle.net/10419/48667

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

WWW.ECONSTOR.EU

Mitglied der Leibniz-Gemeinschaft

IWQW

Institut für Wirtschaftspolitik und Quantitative Wirtschaftsforschung

Diskussionspapier
Discussion Papers

No. 08/2011

Van Zwet Ordering for Fechner Asymmetry

Ingo Klein
University of Erlangen-Nuremberg

ISSN 1867-6707

Van Zwet Ordering for Fechner Asymmetry

Ingo Klein[1]

Department of Statistics and Econometrics
University of Erlangen-Nuremberg, Germany

Abstract

There are several procedures to construct a skewed distribution. One of these procedures splits the value of a parameter of scale for the two halfs of a symmetric distribution. Fechner proposed this procedure in his famous book "Kollektivmaßlehre (1897), p. 295ff.". A similar proposal comes from Fernández et al. (1995). We consider the very general approach from Arellano-Valle et al. (2005) of splitting a scale parameter and show that this technique of generating skewed distributions incorporates a well-defined parameter of skewness. It is well-defined in the sense that the parameter of skewness is compatible with the ordering $\preceq_2$ of van Zwet (1964) which is the strongest ordering in the hierarchy of orderings discussed by Oja (1981). For this family of skewed distributions it will be shown that the measure proposed by Arnold & Groeneveld (1995) is a measure of skewness in the sense of Oja (1981). In the special case considered by Fechner (1897) this measure and the skewness parameter coincide.

Keywords: Skewness; skewness to the right; skewness ordering, measure of skewness

1 Introduction

Starting with a symmetric density f Arellano-Valle et al. (2005) introduce an asymmetric distribution

$$f(x;\gamma) = \frac{2}{a(\gamma)+b(\gamma)}\left(f\left(\frac{x}{a(\gamma)}\right)I(x<0) + f\left(\frac{x}{b(\gamma)}\right)I(x\geq 0)\right) \tag{1}$$

where $a(.)$, $b(.)$ are known positive functions with domain Γ. In the following we call $a(.)$ and $b(.)$ skewness functions. This is a general kind of generating skewness by splitting a scale parameter for the negative and positive half of a distribution that has several well-known special cases. In the following we call this family the AGQ family of skewed distributions.

Setting $a(\gamma) = \gamma$ and $b(\gamma) = 1/\gamma$ for $\Gamma = \mathbb{R}^+$ we get an asymmetric density discussed by Fernández et al. (1995) and Theodossiou (1998). Another choice is $a(\gamma) = 1+\gamma$ and $b(\gamma) = 1-\gamma$ for $\gamma \in \Gamma = (-1,1)$. If f is the normal density we get an asymmetric density already considered by Fechner (1897), p. 295ff. in his famous book "Kollektivmaßlehre".

[1]Correspondence Author: Ingo Klein, Department of Statistics and Econometrics, University of Erlangen Nuremberg, D-90403 Nuremberg, Lange Gasse 20, E-Mail: ingo.klein@wiso.uni-erlangen.de

Arellano-Valle et al. (2005) discuss the statistical properties of (1) for different choices of $a(.)$, $b(.)$ and $f(.)$. Furthermore, they discuss inferential aspects like parameter estimation by the method of moments and by maximum likelihood. For (1) Cassart et al. (2008) construct optimal tests on skewness.

Arellano-Valle et al. (2005, p. 429) or Cassart et al. (2008, p. 2500) call γ an "asymmetry" or a "skewness" parameter. But, due to the work of van Zwet (1964), Oja (1981), Balanda & McGillivray (1990) and others there exist conditions a parameter has to satisfy to be a skewness parameter. Especially, a skewness parameter must hold a specified ordering of skewness. This still has to be proven for the parameter γ in (1).

Klein & Fischer (2006) show for the special choice $a(\gamma) = \gamma$ and $b(\gamma) = 1/\gamma$ (see Fernández et al. (1995)) that γ is a skewness parameter. In detail, they show that $f(x;\gamma)$ is skewed to the right for $\gamma < 1$ and that under some conditions of regularity concerning the symmetric density f γ holds the strong ordering of skewness of van Zwet (1964). This ordering concerns the convexity of

$$\Lambda(x;\gamma_1,\gamma_2)) = F^{-1}(F(.;\gamma_1);\gamma_2) \;\; x \in supp(F(.;\gamma_1)) \tag{2}$$

with the cumulative distribution function $F(.;\gamma)$ and the quantile function $F^{-1}(.;\gamma)$ corresponding to (1).

In this paper we will generalize the results of Klein & Fischer (2006). We show that $\Lambda(.;\gamma_1,\gamma_2)$ is convex or concave for positive, monotone increasing functions $a(.)$ and positive, monotone decreasing functions $b(.)$ with existing derivative $a'(.)$ and $b'(.)$. The conditions of regularity on f are the same as in Klein & Fischer (2006).

The paper is organized in the following way. In section 2 we derive the distribution function, the quantile function, the score function and a function ϕ that is important to show the convexity of (2). Section 3 contains the central result that the AGQ family is a family of skewed distributions in the sense of van Zwet. In section 4 we show for the AGQ family that the skewness measure proposed by Arnold & Groeneveld (1995) is a skewness measure in the sense of Oja. In section 6 we discuss some examples for the skewness functions $a(.)$ and $b(.)$.

2 Some concepts for the measurement of skewness

Oja (1981) p. 7 introduces a location-scale-skewness family of distributions as a family of distributions such that each pair of distributions is skewness comparable. This means that for each pair of distributions holds the van Zwet ordering of skewness.

Definition 1 *Let $\mathcal{F}$ be a family of cumulative distribution functions and $F, G \in \mathcal{F}$ and G^{-1} the quantile function of G.*

1. *F and G will be called skewness comparable if $G^{-1}(F(x))$ is either convex or concave on the support of F.*

2. *F is not more skewed to the right than G (shortly: $F \preceq_2 G$) if $G^{-1}(F(x))$ is convex on the support of F.*

3. *$\mathcal{F}$ is a locations-scale-skewness family if each pair $(F, G) \in \mathcal{F}$ is skewness comparable.*

Transformations of location and scale do not matter the fact that $G^{-1}(F(x))$ is convex or concave. This was proven by Klein & Fischer (2006). We summarize the result in the following lemma.

Lemma 1 *Let F, G be cumulative distribution functions with quantile function G^{-1}. Then*

$$G^{-1}(F(x)) \text{ convex (concave) on } supp(F) \Longrightarrow c + dG^{-1}\left(F\left(\tfrac{x-a}{b}\right)\right),\ b, d > 0,$$
$$a, c \in \mathbb{R} \text{ convex (concave) on } supp(F).$$

This lemma allows to discuss only distributions F and G with unimodal densities and common modus 0.

Whether $G^{-1}(F(.))$ is convex or concave can be checked by the sign of its second derivative. The following lemma proven by Klein & Fischer (2006), p. 1167 or stated by Arnold & Groeneveld (1995), p. 35 gives a sufficient and necessary condition for the convexity (concavity) of $G^{-1}(F(.))$.

Lemma 2 *Let F, G be continuous, cumulative distribution functions with densities f and g. f and g shall be differentiable on $\mathbb{R}$. Define $\phi_f(x) = -f'(x)/f^2(x)$, $\phi_g(x) = -g'(x)/g(x)^2$, $x \in \mathbb{R}$. F^{-1} and G^{-1} are the quantile functions corresponding to F and G. Then $G^{-1}(F(x))$ is convex (concave) on $\mathbb{R}$ iff*

$$\phi_f(F^{-1}(u)) - \phi_g(G^{-1}(u)) \leq (\geq) 0 \text{ for all } u, v \in (0, 1). \tag{3}$$

Skewness shall be measured by a functional that maps a set of distributions in the real numbers and satisfies some requirements that are plausible for the concept of skewness. Oja (1981) gives the following definition for a measure of skewness. If $F \in \mathcal{F}$ belongs to the random variable X, $a \times F + b$ denotes the distribution function of the transformed random variable $aX + b$, $a, b \in \mathbb{R}$.

Definition 2 *Let $\mathcal{F}$ be a family of distributions. $T : \mathcal{F} \to \mathbb{R}$ is a measure of skewness in $\mathcal{F}$ if*

1. *$T(a \times F + b) = sgn(a)T(F)$ for all $a, b \in \mathbb{R}$, $F \in \mathcal{F}$.*
2. *$T(F) \leq T(G)$ if $F, G \in \mathcal{F}$ and $F \preceq_2 G$.*

As a consequence of this definition for a measure of skewness holds

$$T((-1) \times F) = -T(F) \tag{4}$$

This means that reflection of the distribution changes the sign of the measure of skewness.

From the large class of possible measure of skewness we will discuss the proposal of Arnold & Groeneveld (1995) in more detail. If Y is a random variable with uniquely defined modus y_M they propose

$$AG = P(Y < y_M) - P(Y \geq y_M) = 1 - 2P(Y < y_M)$$

as a measure of skewness. AG takes values in $[-1, 1]$. As Ferreira & Steel (2006) p. 823 pronounce this measure "is fairly intuitive for unimodal distributions with negative (positive) values for left (right) skewed distributions and 0 for symmetric distributions".

To show that the AGQ family is a location-scale-skewness family in the sense of Oja the functions $G^{-1}(F(x))$ and ϕ_f have to be computed for this special family.

3 Some functions for the AGQ family

Let F denote the cumulative distribution function of a random variable X and assume that F is continuous on $\mathbb{R}$ and has a density f which itself is differentiable on $\mathbb{R} \setminus \{0\}$. Further, we assume that X is symmetrically distributed. Without restriction of generality we assume that the median of X is 0. Otherwise, we consider $Y = X - \text{median}(X)$. F^{-1} denotes the quantile function of X. We also assume that the support of F is $\mathbb{R}$. The results of the paper can be easily transferred to the case of a random variable X with compact support.

Let X_γ be the random variable with density (1). Then it is easy to verify that X_γ has the following cumulative distribution function

$$\begin{aligned} F(x;\gamma) = & \frac{2a(\gamma)}{a(\gamma)+b(\gamma)} F\left(\frac{x}{a(\gamma)}\right) I(x<0) \\ & + \frac{a(\gamma)-b(\gamma)}{a(\gamma)+b(\gamma)} + \frac{2b(\gamma)}{a(\gamma)+b(\gamma)} F\left(\frac{x}{b(\gamma)}\right) I(x \geq 0). \end{aligned} \tag{5}$$

Inverting the cumulative distribution function leads to the quantile function of X_γ:

$$\begin{aligned} F^{-1}(u;\gamma) = & a(\gamma) F^{-1}\left(\frac{a(\gamma)+b(\gamma)}{2a(\gamma)} u\right) I(A) \\ & + b(\gamma) F^{-1}\left(\frac{a(\gamma)+b(\gamma)}{2b(\gamma)}\left(u - \frac{a(\gamma)-b(\gamma)}{a(\gamma)+b(\gamma)}\right)\right) I(\bar{A}) \end{aligned} \tag{6}$$

with $A = \{u \in [0,1] | u \leq a(\gamma)/(a(\gamma)+b(\gamma)\}$.

With this quantile function we derive the median of X_γ as

$$F^{-1}(0.5;\gamma) = \begin{cases} a(\gamma) F^{-1}\left(\frac{a(\gamma)+b(\gamma)}{a(\gamma)}\right) \leq 0 & \text{for } a(\gamma) > b(\gamma) \\ b(\gamma) F^{-1}\left(\frac{3b(\gamma)-a(\gamma)}{4b(\gamma)}\right) \geq 0 & \text{for } a(\gamma) < b(\gamma) \end{cases} \tag{7}$$

With the help of the cumulative and the inverse distribution functions we get (2). Γ denotes the domain of $a(.)$ and $b(.)$. To restrict the number of cases we have to discuss, we assume that $a(\gamma)$ is monotone increasing with existing derivative $a'(\gamma) > 0$ and $b(\gamma)$ is monotone decreasing with existing derivative $b'(\gamma) < 0$ for $\gamma \in \Gamma$. Then, $a(\gamma)/b(\gamma)$ is monotone increasing and $b(\gamma)/a(\gamma)$ monotone decreasing on Γ.

Let $\gamma_1, \gamma_2 \in \Gamma$ with $\gamma_2 < \gamma_1$ be fixed. Then we get after some tedious calculations

$$\Lambda(x;\gamma_1,\gamma_2) = \begin{cases} a(\gamma_2) F^{-1}\left(\frac{1+b(\gamma_2)/a(\gamma_2)}{1+b(\gamma_1)/a(\gamma_1)} F\left(\frac{x}{a(\gamma_1)}\right)\right) \\ \quad \text{for } x \leq a(\gamma_1) F^{-1}\left(\frac{1}{2}\frac{1+b(\gamma_1)/a(\gamma_1)}{1+b(\gamma_2)/a(\gamma_2)}\right) \\ b(\gamma_2) F^{-1}\left(\frac{1+a(\gamma_2)/b(\gamma_2)}{1+b(\gamma_1)/a(\gamma_1)} F\left(\frac{x}{a(\gamma_1)}\right) - \frac{a(\gamma_2)-b(\gamma_2)}{2b(\gamma_2)}\right) \\ \quad \text{for } a(\gamma_1) F^{-1}\left(\frac{1}{2}\frac{1+b(\gamma_1)/a(\gamma_1)}{1+b(\gamma_2)/a(\gamma_2)}\right) < x \leq 0 \\ b(\gamma_2) F^{-1}\left(\frac{1+a(\gamma_2)/b(\gamma_2)}{1+a(\gamma_1)/b(\gamma_1)} \frac{a(\gamma_1)-b(\gamma_1)}{2b(\gamma_1)} - \frac{a(\gamma_2)-a(\gamma_2)}{2b(\gamma_2)} + \frac{1+a(\gamma_2)/b(\gamma_2)}{1+a(\gamma_1)/b(\gamma_1)} F\left(\frac{x}{b(\gamma_1)}\right)\right) \\ \quad \text{for } x > 0 \end{cases} \tag{8}$$

Whether $\Lambda(.;\gamma_1,\gamma_2)$ is convex or concave can be checked with the following ϕ-function: Let $\phi(x) = -f'(x)/f^2(x)$ and $\phi(x;\gamma) = -f'(x;\gamma)/f^2(x;\gamma)$ for $x \in \mathbb{R}$. Then it is easy to show that

$$\begin{aligned}\phi(x;\gamma) = &\frac{1}{2}\left(1+\frac{b(\gamma)}{a(\gamma)}\right)\phi\left(\frac{x}{a(\gamma)}\right)I(x<0) \\ &+\frac{1}{2}\left(1+\frac{a(\gamma)}{b(\gamma)}\right)\phi\left(\frac{x}{b(\gamma)}\right)I(x\geq 0).\end{aligned} \tag{9}$$

Let $I \subseteq supp(F(.;\gamma_1)$ such that $f'(x;\lambda_1)$ and $f'(F^{-1}(F(x;\lambda_1);\lambda_2);\lambda_2)$ exists for $x \in I$ then $\Lambda(.;\gamma_1,\gamma_2)$ is convex (concave) on I iff

$$\phi(x;\gamma_1) - \phi\left(F^{-1}\left(F(x;\gamma_1);\gamma_2\right);\gamma_2\right) \leq (\geq) 0$$

for $x \in I$. Obviously, the interval I may not contain $x = 0$ or $x = F^{-1}(a(\gamma_2)/(a(\gamma_2) + b(\gamma_2));\gamma_1)$.

An important property is the monotonicity of ϕ to show under which conditions $\Lambda(.;\gamma_1,\gamma_2)$ is convex or concave. Klein & Fischer (2006) discuss some examples for distributions such that $\phi(.)$ is monotone increasing. To these distributions belong among others the Gaussian and the t distribution, the Laplace distribution and the generalized secant hyperbolic distribution of Vaughan (2002). A counterexample is the generalized t distribution of McDonald & Newey (1988).

4 AGQ family as a location-scale-skewness family

Let $\mathcal{F}$ be the AGQ family with densities (1). Then we have to show that $F(.;\gamma_1), F(.;\gamma_2))$ are skewness comparable. This means that $F^{-1}(F(x;\gamma_1);\gamma_2)$ is either convex or concave on $\mathbb{R}$. We will prove this result under some assumptions. The first set of assumptions concerns the symmetric density f. Especially, $\phi(x) = -f'(x)/f(x)^2$ has to be monotone increasing with derivative $\phi'(x) > 0$. The second set of assumptions concerns the skewness functions $a(.)$ and $b(.)$. $a(.)$ and $b(.)$ shall be differentiable on Γ with $a'(\gamma) > 0$ and $b'(\gamma) < 0$ for $\gamma \in \Gamma$. This means that $a(\gamma)/b(\gamma)$ is monotone increasing on the common domain Γ.

Notice that the ϕ-function of $F(.;\gamma)$ is only defined for $x \neq 0$ because $f(.;\gamma)$ is continuous, but not differentiable at $x = 0$. This demands a special treatment at $x = 0$.

Two lemmata will prepare the main result that the AGQ family is a location-scale-skewness family in the sense of Oja.

Lemma 3 *Let F be a continuous distribution function with unimodal and symmetric density function f that is continuous on $\mathbb{R}$ and differentiable for $\{\mathbb{R}\setminus 0\}$ such that $\phi'(x) > 0$ for $x \neq 0$. Denote $\mathcal{F} = \{F(.;\gamma)|\gamma \in \Gamma\}$ the AGQ family of distributions with positive skewness functions $a(.)$ and $b(.)$. Furthermore, we assume that $a(.)$ an d b(.) are differentiable with $a'(\gamma) > 0$ and $b'(\gamma) < 0$ for $\gamma \in \Gamma$. Then*

$$\frac{\partial\phi(F^{-1}(u;\gamma);\gamma)}{\partial\gamma} = \frac{\partial\phi(x;\gamma)}{\partial\gamma}|_{x=F^{-1}(u;\gamma)}\frac{\partial F^{-1}(u;\gamma)}{\partial\gamma} < 0$$

for $u < a(\gamma)/a(\gamma) + b(\gamma))$ *or* $u > a(\gamma)/a(\gamma) + b(\gamma))$ *and* $u \in (0, 1)$.

Proof:

- Discussion of $\partial\phi(x;\gamma)/\partial\gamma$ for $x \neq 0$:

  Let $x < 0$:

$$\begin{aligned}\frac{\partial\phi(x;\gamma)}{\partial\gamma} &= 1/2\frac{\partial b(\gamma)/a(\gamma)}{\partial\gamma}\phi\left(\frac{x}{a(\gamma)}\right)\\ &\quad +1/2(1+b(\gamma)/a(\gamma))\phi'\left(\frac{x}{a(\gamma)}\right)\left(-\frac{x}{a(\gamma)^2}\right)a'(\gamma) > 0\end{aligned}$$

  because $a'(\gamma) > 0$, $b'(\gamma) < 0$, $\phi(x) < 0$ for $x < 0$ and $\phi'(x) > 0$ for $x \in \mathbb{R}$.

  Let $x > 0$. Then we get

$$\begin{aligned}\frac{\partial\phi(x;\gamma)}{\partial\gamma} &= 1/2\frac{\partial a(\gamma)/b(\gamma)}{\partial\gamma}\phi\left(\frac{x}{b(\gamma)}\right)\\ &\quad +1/2(1+a(\gamma)/b(\gamma))\phi'\left(\frac{x}{b(\gamma)}\right)\left(-\frac{x}{b(\gamma)^2}\right)b'(\gamma) > 0\end{aligned}$$

  due to $\phi(x) > 0$, $x > 0$.

- Discussion of $\partial F^{-1}(.;\gamma)/\partial\gamma$ for $u \neq a(\gamma)/(a(\gamma) + b(\gamma)$:

  Moreover, fur $u < a(\gamma)/(a(\gamma) + b(\gamma))$

$$\begin{aligned}\frac{\partial F^{-1}(u;\gamma)}{\partial\gamma} &= a'(\gamma)F^{-1}\left(1/2\left(1+b(\gamma)/a(\gamma)\right)u\right)\\ &\quad +1/2a(\gamma)\frac{1}{f\left(F^{-1}\left(1/2(1+b(\gamma)/a(\gamma))\,u\right)\right.}\frac{\partial b(\gamma)/a(\gamma)}{\partial\gamma} < 0\end{aligned}$$

  because $F^{-1}\left(1/2\left(1+b(\gamma)/a(\gamma)\right)u\right) < 0$ for $u < a(\gamma)/(a(\gamma)+b(\gamma))$ and $\frac{\partial b(\gamma)/a(\gamma)}{\partial\gamma} < 0$.

  Fur $u > a(\gamma)/(a(\gamma) + b(\gamma))$ we get

$$\begin{aligned}\frac{\partial F^{-1}(u;\gamma)}{\partial\gamma} &= b'(\gamma)F^{-1}\left(1/2\left(1+a(\gamma)/b(\gamma)\right)u + 1 - a(\gamma)/b(\gamma)\right)\\ &\quad +1/2b(\gamma)\frac{1}{f\left(F^{-1}\left(1/2(1+b(\gamma)/a(\gamma))\,u + 1 - a(\gamma)/b(\gamma)\right)\right.}\\ &\quad \cdot\frac{\partial a(\gamma)/b(\gamma)}{\partial\gamma}(u-1) < 0\end{aligned}$$

  because $F^{-1}\left(1/2\left(1+a(\gamma)/b(\gamma)\right)u + 1 - a(\gamma)/b(\gamma)\right) > 0$ for $u > a(\gamma)/(a(\gamma) + b(\gamma))$, $\frac{\partial b(\gamma)/a(\gamma)}{\partial\gamma} > 0$ and $u - 1 < 0$.

- Combining these results we get

$$\frac{\partial \phi(F^{-1}(u;\gamma);\gamma)}{\partial \gamma} = \frac{\partial \phi(x;\gamma)}{\partial \gamma}|_{x=F^{-1}(u;\gamma)} \frac{\partial F^{-1}(u;\gamma)}{\partial \gamma} < 0$$

for $u < a(\gamma)/a(\gamma) + b(\gamma))$ or $u > a(\gamma)/(a(\gamma) + b(\gamma)$ and $u \in (0,1)$. □

Lemma 4 *Let F be a continuous distribution function with unimodal and symmetric density function f that is positive and continuous on $\mathbb{R}$. Let $F(.;\gamma_1)$, $F(.;\gamma_2)$ be elements of the AGQ family of distributions. Then*

$$\Lambda'(x;\gamma_1,\gamma_2) = \frac{\partial F^{-1}(F(x;\gamma_1);\gamma_2)}{\partial x} > 0 \ \textit{for}\ x \in \mathbb{R}$$

and continuous on $\mathbb{R}$.

Proof: It holds

$$\Lambda'(x;\gamma_1,\gamma_2) = \frac{f(x;\gamma_1)}{f(F^{-1}(F(x;\gamma_1);\gamma_2);\gamma_2)} \quad x \in \mathbb{R}. \tag{10}$$

$f(.;\gamma) > 0$ and $F(.;\gamma)$ are continuous functions on $\mathbb{R}$. $F^{-1}(u;\gamma)$ is continuous on $(0,1)$. Therefore, $\Lambda'(.;\gamma_1,\gamma_2) > 0$ and continuous for $x \in \mathbb{R}$. □

Theorem 1 *Let F be a continuous distribution function with unimodal and symmetric density function f that is positive and continuous on $\mathbb{R}$ and differentiable for $\{\mathbb{R}\backslash 0\}$ such that $\phi'(x) > 0$ for $x \neq 0$. Denote $\mathcal{F} = \{F(.;\gamma)|\gamma \in G\}$ the AGQ family of distributions skewness functions $a(.)$ and $b(.)$. Furthermore, we assume that $a(.)$ and b(.) are differentiable with $a'(\gamma) > 0$ and $b'(\gamma) < 0$ for $\gamma \in \Gamma$. Then $\mathcal{F}$ is a location-scale-skewness family.*

Proof: We have to show that all members of $\mathcal{F}$ are skewness comparable. This means that $\Lambda(x;\gamma_1,\gamma_2)$ is either convex or concave on $\mathbb{R}$ for all $\gamma_1,\gamma_2 \in \Gamma$.

With lemma 3 we know that $\phi(F^{-1}(u;\gamma)$ is monotone decreasing in γ for $u \neq a(\gamma)/(a(\gamma)+b(\gamma)$.

For $\gamma_2 < \gamma_1$ it is

$$\frac{a(\gamma_2)}{a(\gamma_2)+b(\gamma_2)} < \frac{a(\gamma_1)}{a(\gamma_1)+b(\gamma_1)}.$$

The interval $(0,1)$ can be divided into three subsets:

$$I_1 = \left(0, \frac{a(\gamma_2)}{a(\gamma_2)+b(\gamma_2)}\right), \quad I_2 = \left(\frac{a(\gamma_2)}{a(\gamma_2)+b(\gamma_2)}, \frac{a(\gamma_1)}{a(\gamma_1)+b(\gamma_1)}\right), \quad I_3 = \left(\frac{a(\gamma_1)}{a(\gamma_1)+b(\gamma_1)}\right).$$

$\phi(F^{-1}(u;\gamma))$ is monotone decreasing in γ for $u \in I_1 \cup I_2 \cup I_3)$. This means that

$$\phi(F^{-1}(u;\gamma_1),\gamma_1) - \phi(F^{-1}(u;\gamma_2);\gamma_2) \ \text{ for } u \in I_1 \cup I_2 \cup I_3.$$

Therefore, $\Lambda(.;\gamma_1,\gamma_2)$ is convex on the intervals

$$J_1 = \left(-\infty, F^{-1}\left(\frac{a(\gamma_2)}{(a(\gamma_2)+b(\gamma_2)}\right)\right), \quad J_2 = \left(F^{-1}\left(\frac{a(\gamma_2)}{(a(\gamma_2)+b(\gamma_2)}, 0\right)\right), \quad J_3 = (0,\infty)$$

corresponding to I_1, I_2 and I_3. With lemma 4 $\Lambda'(.;\gamma_1,\gamma_2)$ is continuous on $\mathbb{R}$. Therefore, we can generalize the convexity of $\Lambda(.;\gamma_1,\gamma_2)$ from the three subsets J_1, J_2, J_3 to the whole real line.

A similar discussion for $\gamma_2 > \gamma_1$ leads to the conclusion that $\Lambda(.;\gamma_1,\gamma_2)$ is concave on $\mathbb{R}$ for this case. $\square$.

Notice that $\phi(F^{-1}(u;\gamma))$ is strictly decreasing. Therefore, $\Lambda(.;\gamma_1,\gamma_2)$ is strictly increasing and either strictly convex or strictly concave on $\mathbb{R}$.

5 Skewness measure of Arnold & Groeneveld for the AGQ family

Arnold & Groeneveld (1995) propose

$$AG = P(Y < y_M) - P(Y > y_M) = 1 - 2P(Y < y_M).$$

as a measure of skewness for a random variable Y with unique modus y_M. For a symmetric density $f(.)$ with modus 0 $f(,;\gamma)$ also has modus 0 with $F(0;\gamma) = a(\gamma)/(a(\gamma)+b(\gamma)$. Therefore, we get for the skewness measure of Arnold & Groeneveld

$$AG(F(.;\gamma)) = 1 - 2\frac{a(\gamma)}{a(\gamma)+b(\gamma)} = \frac{b(\gamma)-a(\gamma)}{a(\gamma)+b(\gamma)}.$$

Notice that AG does not depend on the underlying symmetric density $f(.)$. Under the assumption that $a(.)$ ($(b(.)$) is monotone increasing (decreasing) AG is monotone decreasing in γ. This means

$$\gamma_2 < \gamma_1 \iff AG(\gamma_1) < AG(\gamma_2).$$

Now, we want to show for the AGQ family that AG is a measure of skewness in the sense of Oja.

Corollary 5.1 *Let F be a continuous distribution function with unimodal and symmetric density function f that is continuous on $\mathbb{R}$ and differentiable for $\{\mathbb{R}\setminus 0\}$ such that $\phi'(x) > 0$ for $x \neq 0$. Denote $\mathcal{F} = \{F(.;\gamma)|\gamma \in G\}$ the AGQ family of distributions skewness functions $a(.)$ and $b(.)$. Further, we assume that $a(.)$ and b(.) are differentiable with $a'(\gamma) > 0$ and $b'(\gamma) < 0$ for $\gamma \in \Gamma$. Then AG is measure of skewness in the sense of Oja.*

Proof:

1. Let X_γ be distributed with $F(x;\gamma)$, $\gamma \in \Gamma$. X_γ has modus 0 such that $(X_\gamma - b)/a$ has modus $-b/a$.

   Consider $a \times F(.;\gamma) + b$ for $a, b \in \mathbb{R}$. Then

$$AG(a \times F(.;\gamma) + b) = 1 - 2P\left(\frac{X_\gamma - b}{a} < -\frac{b}{a}\right).$$

We get

$$AG(a \times F(.;\gamma) + b) = \begin{cases} 1 - 2P(X_\gamma < 0) = AG(F(.;\gamma) & \text{for } a > 0 \\ 1 - 2P(X_\gamma > 0) \\ = 1 - 2(1 - P(X_\gamma < 0)) \\ = -1 + 2P(X_\gamma < 0) = -AG(F(.;\gamma) & \text{for } a < 0 \end{cases}$$

Hence, $AG(a \times F(.;\gamma) + b) = sgn(a)AG(F(.;\gamma)$.

2. It remains to show that if $F(.;\gamma_1) \preceq_2 F(.;\gamma_2)$ then it holds $AG(\gamma_1) \leq AG(\gamma_2)$.

   From lemma 3 and the proof of theorem 1 we know that $\phi(F^{-1}(.;\gamma)$ is a strictly decreasing function in γ for $u \in I_1 \cup I_2 \cup I_3$ if $\Lambda(.;\gamma_1,\gamma_2)$ is strictly convex on $\mathbb{R}$. Therefore, it holds for $\gamma_1, \gamma_2 \in \Gamma$:

$$\phi(F^{-1}(.;\gamma_1)) - \phi(F^{-1}(.;\gamma_2) \Longrightarrow \gamma_2 < \gamma_1.$$

   $AG(.)$ is strictly decreasing on Γ. This leads to

$$\Lambda(.;\gamma_1,\gamma_2) \text{ is srictly convex on } \mathbb{R} \Longrightarrow AG(F(.;\gamma_1)) < AG(F(.;\gamma_2)).$$

   for $\gamma_1, \gamma_2 \in \Gamma$. $\square$

6 Special cases

6.1 Symmetry for $\gamma = 0$ and invariance w.r.t to reflection

For monotone increasing (decreasing) functions $a(.)$ ($b(.)$) the density (1) is symmetric around 0 if $a(\gamma) = b(\gamma)$.

1. For $a(\gamma) = \gamma$ and $b(\gamma) = 1/\gamma$ we get symmetry for $\gamma = 1$.
2. Setting $a(\gamma) = 1 + \gamma$ and $b(\gamma) = 1 - \gamma$ gives symmetry for $\gamma = 0$.

After the reparametrization $a(\gamma) = e^\gamma$ and $b(\gamma) = e^{-\gamma}$ in the first case we get also symmetry for $\gamma = 0$. Therefore, we restrict us on functions $a(.)$ and $b(.)$ such that $a(\gamma) = b(\gamma)$ for $\gamma = 0$. This means that 0 is in the domain Γ.

Let X_γ the random variable corresponding to the density (1). $-X_\gamma$ denotes the reflected random variable with reflected properties of skewness. If X_γ is skewed to the right, $-X_\gamma$ should be skewed to the left in the same manner. It is plausible to require that the skewness of $-X_\gamma$ should be the same as the skewness of $X_{-\gamma}$. This leads to the condition

$$f(x;-\gamma) = f(-x;\gamma) \;\; x \in \mathbb{R}$$

for $\gamma \in \Gamma$. For $x = 0$ we get

$$f(0;-\gamma) = \frac{2}{a(-\gamma) + b(-\gamma)} f(0) = \frac{2}{a(\gamma) + b(\gamma)} f(0) = f(0;\gamma).$$

This condition is satisfied if

$$b(\gamma) = a(-\gamma) \;\; \gamma \in \Gamma. \tag{11}$$

The splitting of scale originally proposed by Fechner gives an example for this special case. Another example is $a(\gamma) = e^{\gamma}$ and $b(\gamma) = e^{-\gamma}$ for $\gamma \in \mathbb{R}$.

Under (11) Arnold & Groeneveld's measure of skewness AG takes the form

$$AG(\gamma) = \frac{a(-\gamma) - a(\gamma)}{a(\gamma) + a(-\gamma)}.$$

6.2 Asymmetric distributions due to Fechner

We follow Cassart et al. (2008) and discuss a slightly more general version of Fechner's normal density with different scale parameters for positive and negative arguments. Set

$$a(\gamma) = 1 + a\gamma \;\text{ and }\; b(\gamma) = 1 - b\gamma, \;\; \gamma \in (-1, 1) \tag{12}$$

with $a, b > 0$. Obviously, $a(\gamma)$ is monotone increasing and $b(\gamma)$ is monotone decreasing for $\gamma > 0$. This proofs the following corollary.

Corollary 6.1 *Let F be a continuous distribution function with unimodal and symmetric density function f that is continuous on $\mathbb{R}$ and differentiable for $\{\mathbb{R} \setminus 0\}$ such that $\phi'(x) > 0$ for $x \neq 0$. Consider the special AGQ family of skewed densities*

$$f(x;\gamma) = \frac{2}{2 + (a-b)\gamma}\left(f\left(\frac{x}{1+a\gamma}\right) I(x \leq 0) + f\left(\frac{1}{1-b\gamma}\right) I(x > 0)\right) \;\; x \in \mathbb{R}$$

with $a, b > 0$ and $\gamma \in (-1, 1)$. Then, this family is location-scale-skewness family in the sense of Oja.

Under the setting (12) we get for the Arnold & Groeneveld's measure of skewness

$$AG(F(.;\gamma)) = -\frac{(a+b)\gamma}{2 + (a-b)\gamma}.$$

In the Fechner case with $a = b = 1$ we get $AG(F.;\gamma) = -\gamma$ for $\gamma \in [-1, 1]$. This means that Arnold & Groeneveld's skewness measure and the negative of the skewness parameter γ are identical.

6.3 Asymmetric distribution due to Fernández et al.

We consider the proposal of Fernández et al. for skewed distributions with the modified choice

$$a(\gamma) = e^{\gamma} \;\text{ and }\; b(\gamma) = e^{-\gamma} \;\; \gamma \in \mathbb{R}. \tag{13}$$

$a(.)$ is monotone increasing and $b(.)$ monotone decreasing on $\mathbb{R}$ and $a(\gamma) > b(\gamma)$ for $\gamma > 0$. This proves the following corollary which was already proven by Klein & Fischer (2006).

Corollary 6.2 *Let F be a continuous distribution function with unimodal and symmetric density function f that is continuous on $\mathbb{R}$ and differentiable for $\{\mathbb{R} \setminus 0\}$ such that $\phi'(x) > 0$ for $x \neq 0$. Consider the special AGQ family of skewed densities*

$$f(x;\gamma) = \frac{2}{e^{\gamma} + e^{-\gamma}} \left(f\left(xe^{-\gamma}\right) I(x \leq 0) + f\left(xe^{\gamma}\right) I(x > 0) \right) \quad x \in \mathbb{R}$$

with $\gamma \in \mathbb{R}$. Then, this family is location-scale-skewness family in the sense of Oja.

The calculation of Arnold & Groeneveld's AG is straightforward.

7 Summary

There are several procedures to construct a skewed distribution. One of these procedures splits the value of a parameter of scale for the two sides of a symmetric distribution. We show that the most general form of this technique of generating skewed distributions proposed by Arellano-Valle et al. (2005) incorporates a well-defined parameter of skewness. It is well-defined in the sense that the generated distributions are skewed to the right if the parameter of skewness takes values less than 1. As second property we show that the parameter of skewness is compatible with the ordering $\preceq_2$ of van Zwet (1964) which is the strongest ordering in the hierarchy of orderings discussed by Oja (1981). In this sense the generated skewed distributions can be ordered by the parameter of skewness. We show how the measure of skewness of Arnold & Groeneveld depends on the skewness parameter.

References

[1] Arellano-Valle, R.B., H.W. Gomez, F.A. Quintana: *Statistical Inference for a General Class of Asymmetric Distributions*. Journal of Statistical Planning and Inference, **128**:427-443, 2005.

[2] Arnold, B.C. and R.A. Groeneveld: *Measuring Skewness with Respect to the Mode*. The American Statistician, **49**(1):34-38, 1995.

[3] Balanda, K. P. and H. L. MacGillivray: *Kurtosis and Spread*. The Canadian Journal of Statistics, **18**(1):17-30, 1990.

[4] Cassart, D., M. Hallin, D. Paindaveine: *Optimal Detection of Fechner-Asymmetry*. Journal of Statistical Planning and Inference, **138**:2499-2525, 2008.

[5] Fechner, G.T.: *Kollektivmaßlehre*. Engelmann, Leipzig, 1897.

[6] Fernández, C., J. Osiewalski and M. F. J. Steel: *Modelling and Inference with ν-Spherical Distributions*. Journal of American Statistical Association, **90**:1331-1340, 1995.

[7] Klein, I., M. Fischer: *Skewness and Splitting the Scale Parameter*. Communications in Statistics, Theory and Methods, **35**(7):1159-1171, 2006.

[8] McDonald, J. B. and W. K. Newey: *Partially Adaptive Estimation of Regression Models via the Generalized-t-Distribution*. Econometric Theory, **1**(4):428-457, 1988.

[9] Oja, H.: *On Location, Scale, Skewness and Kurtosis of Univariate Distributions*. Scandinavian Journal of Statistics, **18**:154-168, 1981.

[10] Theodossiou, P.: *Financial Data and the Skewed Generalized t distribution*. Management Science, **44**(12/1):1650-1661, 1998.

[11] Van Zwet, W. R.: *Convex Transformations of Random Variables*. Mathematical Centre Tracts No. 7. Mathematical Centre, Amsterdam, 1964.

[12] Vaughan, D. C.: *The Generalized Hyperbolic Secant distribution and its Application*. Communications in Statistics – Theory and Methods, **31**(2):219-238, 2002.

Diskussionspapiere 2011
Discussion Papers 2011

01/2011 **Klein, Ingo, Fischer, Matthias, Pleier, Thomas:** Weighted Power Mean Copulas: Theory and Application

02/2011 **Kiss, David:** The Impact of Peer Ability and Heterogeneity on Student Achievement: Evidence from a Natural Experiment

03/2011 **Zibrowius, Michael:** Convergence or divergence? Immigrant wage assimilation patterns in Germany

04/2011 **Klein, Ingo, Christa, Florian:** Families of Copulas closed under the Construction of Generalized Linear Means

05/2011 **Schnitzlein, Daniel**: How important is the family? Evidence from sibling correlations in permanent earnings in the US, Germany and Denmark

06/2011 **Schnitzlein, Daniel**: How important is cultural background for the level of intergenerational mobility?

07/2011 **Steffen Mueller**: Teacher Experience and the Class Size Effect - Experimental Evidence

Diskussionspapiere 2010
Discussion Papers 2010

01/2010 **Mosthaf, Alexander, Schnabel, Claus and Stephani, Jens:** Low-wage careers: Are there dead-end firms and dead-end jobs?

02/2010 **Schlüter, Stephan and Matt Davison:** Pricing an European Gas Storage Facility using a Continuous-Time Spot Price Model with GARCH Diffusion

03/2010 **Fischer, Matthias, Gao, Yang and Herrmann, Klaus:** Volatility Models with Innovations from New Maximum Entropy Densities at Work

04/2010 **Schlüter, Stephan, Deuschle, Carola:** Using Wavelets for Time Series Forecasting – Does it Pay Off?

05/2010 **Feicht, Robert, Stummer, Wolfgang:** Complete closed-form solution to a stochastic growth model and corresponding speed of economic recovery.

06/2010 **Hirsch, Boris, Schnabel, Claus:** Women Move Differently: Job Separations and Gender.

07/2010 **Gartner, Hermann, Schank, Thorsten, Schnabel, Claus:** Wage cyclicality under different regimes of industrial relations.

08/2010 **Tinkl, Fabian**: A note on Hadamard differentiability and differentiability in quadratic mean.

09/2010 **Tinkl, Fabian**: Asymptotic theory for M estimators for martingale differences with applications to GARCH models.

Diskussionspapiere 2009
Discussion Papers 2009

01/2009 **Addison, John T. and Claus Schnabel:** Worker Directors: A German Product that Didn't Export?

02/2009 **Uhde, André and Ulrich Heimeshoff:** Consolidation in banking and financial stability in Europe: Empirical evidence

03/2009 **Gu, Yiquan and Tobias Wenzel:** Product Variety, Price Elasticity of Demand and Fixed Cost in Spatial Models

04/2009 **Schlüter, Stephan:** A Two-Factor Model for Electricity Prices with Dynamic Volatility

05/2009 **Schlüter, Stephan and Fischer, Matthias**: A Tail Quantile Approximation Formula for the Student t and the Symmetric Generalized Hyperbolic Distribution

06/2009 **Ardelean, Vlad:** The impacts of outliers on different estimators for GARCH processes: an empirical study

07/2009 **Herrmann, Klaus:** Non-Extensitivity versus Informative Moments for Financial Models - A Unifying Framework and Empirical Results

08/2009 **Herr, Annika:** Product differentiation and welfare in a mixed duopoly with regulated prices: The case of a public and a private hospital

09/2009 **Dewenter**, Ralf, **Haucap**, Justus and **Wenzel**, Tobias: Indirect Network Effects with Two Salop Circles: The Example of the Music Industry

10/2009 **Stuehmeier**, Torben and **Wenzel**, Tobias: Getting Beer During Commercials: Adverse Effects of Ad-Avoidance

11/2009 **Klein**, Ingo, **Köck**, Christian and **Tinkl**, Fabian: Spatial-serial dependency in multivariate GARCH models and dynamic copulas: A simulation study

12/2009 **Schlüter**, Stephan: Constructing a Quasilinear Moving Average Using the Scaling Function

13/2009 **Blien,** Uwe, **Dauth,** Wolfgang**, Schank,** Thorsten and **Schnabel,** Claus**:** The institutional context of an "empirical law": The wage curve under different regimes of collective bargaining

14/2009 **Mosthaf**, Alexander, **Schank**, Thorsten and **Schnabel**, Claus: Low-wage employment versus unemployment: Which one provides better prospects for women?

Diskussionspapiere 2008
Discussion Papers 2008

01/2008 **Grimm, Veronika and Gregor Zoettl**: Strategic Capacity Choice under Uncertainty: The Impact of Market Structure on Investment and Welfare

02/2008 **Grimm, Veronika and Gregor Zoettl**: Production under Uncertainty: A Characterization of Welfare Enhancing and Optimal Price Caps

03/2008 **Engelmann, Dirk and Veronika Grimm**: Mechanisms for Efficient Voting with Private Information about Preferences

04/2008 **Schnabel, Claus and Joachim Wagner**: The Aging of the Unions in West Germany, 1980-2006

05/2008 **Wenzel, Tobias**: On the Incentives to Form Strategic Coalitions in ATM Markets

06/2008 **Herrmann, Klaus**: Models for Time-varying Moments Using Maximum Entropy Applied to a Generalized Measure of Volatility

07/2008 **Klein, Ingo and Michael Grottke:** On J.M. Keynes' "The Principal Averages and the Laws of Error which Lead to Them" - Refinement and Generalisation