

Jaschke, Stefan R.

Working Paper

A note on stochastic volatility, GARCH models, and hyperbolic distributions

SFB 373 Discussion Paper, No. 1998,23

Provided in Cooperation with:

Collaborative Research Center 373: Quantification and Simulation of Economic Processes,
Humboldt University Berlin

Suggested Citation: Jaschke, Stefan R. (1997) : A note on stochastic volatility, GARCH models, and hyperbolic distributions, SFB 373 Discussion Paper, No. 1998,23, Humboldt University of Berlin, Interdisciplinary Research Project 373: Quantification and Simulation of Economic Processes, Berlin, <https://nbn-resolving.de/urn:nbn:de:kobv:11-10056700>

This Version is available at:

<https://hdl.handle.net/10419/61282>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

A Note on Stochastic Volatility, GARCH models, and Hyperbolic Distributions

Stefan R. Jaschke, *

December 13, 1997

We establish a relation between stochastic volatility models and the class of generalized hyperbolic distributions. These distributions have been found to fit exceptionally well to the empirical distribution of stock returns. We review the background of hyperbolic distributions and prove stationary distributions of certain GARCH-type models to be generalized hyperbolic.

1 Introduction

The aim of this paper is to gather some results about the empirical distribution of stock returns and some attempts to match the “heavy tails” of these distributions in stochastic processes models, and to give an introductory overview of the appearance of generalized hyperbolic distributions in this context.

There are some well-known empirical facts about log returns on stocks:

- (i) The empirical distribution is leptokurtic compared to the normal distribution.
- (ii) Although there is no significant serial correlation in stock returns, there is serial correlation in squared log returns.

*Humboldt-Universität Berlin, Institut für Mathematik. The main part of the study of hyperbolic distributions and the research into stochastic volatility models was done when I was a guest at Aarhus University in 1994. As I continued to work in a different field, the work never made it into a paper. During a discussion three years later I was encouraged to make a small note about the subject. I like to thank Uwe Küchler, Michael Sørensen, and Rolf Tschernig for helpful comments on this paper. The note was finalized at the Sonderforschungsbereich 373 at Humboldt University Berlin. Financial support by the Deutsche Forschungsgemeinschaft is gratefully acknowledged.

Empirical and theoretical investigations of (i) have a long history. Mandelbrot and Fama proposed *Pareto-stable distributions* to explain the excess kurtosis in stock returns ((Fama; 1965), (Mandelbrot and Taylor; 1967)). Mittnik and Rachev (1993) give an overview and comparison of alternative distributions in modeling stock returns. Evidence against stable Paretian distributions had been accumulated, and Mittnik and Rachev suggest the double Weibull-distribution for stock returns, whose density is

$$f(x; \alpha, \lambda) = \frac{1}{2} \lambda |x|^{\alpha-1} \exp(-\lambda x^\alpha) \quad (\alpha > 0, \lambda > 0).$$

One of their arguments in favor of the Weibull distribution is that tails decrease exponentially. (They estimate α to be close to 1.)

Hyperbolic distributions, which also have exponentially decreasing tails, were independently suggested as distributions of German stock returns by Eberlein and Keller (1994) and Küchler et al. (1994). (The logarithm of the density of a hyperbolic distribution is a hyperbola.) Hyperbolic distributions seem to fit exceptionally well to the return in German stocks represented in the Stock index DAX. Barndorff-Nielsen fitted generalized hyperbolic distributions to Danish stock returns (Barndorff-Nielsen; 1994).

Let $R_k = \log S_k - \log S_{k-1}$ denote the (one-period) log return, given a stock price process $(S_t)_{t \in [0, \infty)}$. Suppose that (R_k) is stationary. We think that the important question now is what kind of stochastic process (R_k) is, rather than looking at its marginal distributions only. Küchler et al. suggest

$$\begin{aligned} dX_t &= m(X_t)dt + \sigma dW_t \\ R_n &= X_n \\ m(x) &= -\frac{\sigma^2}{2} \left(\alpha \frac{x - \mu}{\sqrt{\delta^2 + (x - \mu)^2}} - \beta \right) \end{aligned}$$

$m(x)$ is chosen in such a way that the stationary distribution of X is hyperbolic. This approach is not satisfying for several reasons. First, a nontrivial $m(x)$ induces an autocorrelated series (R_k) which is not what we observe. Second, in the continuous time version of the model, the stock price would be

$$S_t = \exp\left\{\int_0^t X_s ds + \mu t\right\},$$

which is absolutely continuous and of bounded variation. So, there would be arbitrage in the continuous-time option pricing theory (Harrison and Pliska; 1981), if the locally riskless security $S_t^0 = \exp\{\int_0^t r_s ds\}$ wasn't identical to S_t .




Figure 1: Autocorrelation in weekly log returns in the DAX 1974 – 1992

In this sense, the model proposed in (Küchler et al.; 1994) is not suitable for the valuation of derivative securities.

Barndorff-Nielsen (1994) suggests modeling $\log S_t$ with processes whose increments are independent and generalized hyperbolically distributed. Stock returns are, however, dependent, although there is no significant linear dependence (serial correlation). The fact, that there is significant serial correlation in squared residuals of many economic time series led to the development of stochastic volatility models, one important instance being the GARCH models.

Figure (1) shows the autocorrelation and partial autocorrelation of the weekly (log) return in the DAX and its square, respectively. The dotted line shows the level of significance as given by the S-Plus function **acf**. All lags except the third show insignificant autocorrelation. Judging from both the autocorrelation function and the partial autocorrelation function, weekly DAX returns very much look like white noise. Looking at squared returns, however, significant autocorrelation up to for weeks can be seen. There also seems to be some autocorrelation of squared returns at 13 weeks (a quarter of a year).

In section 2, we review some facts about generalized hyperbolic distributions and normal variance-mean mixtures. We show in section 3 that certain GARCH-type models produce stationary marginal distributions that

are generalized hyperbolic.

2 Generalized Hyperbolic Distributions as Normal Variance-Mean Mixture Distributions

Hyperbolic distributions were introduced by Barndorff-Nielsen (1977) in the context of the distribution of the size of sand particles taken from the Danish coast. Plots of the log histogram of their log size seemed to have linear tails and could be very well fitted by hyperbolae. This naturally led to the definition of the hyperbolic distribution, whose log density is a hyperbola:

$$\log h(x; \alpha, \beta, \mu, \delta) = c(-\alpha\sqrt{\delta^2 + (x - \mu)^2} + \beta(x - \mu)). \quad (1)$$

(c is a norming constant, μ is a location parameter, δ a scale parameter, and $(\alpha + \beta)x$ and $(-\alpha + \beta)x$ are the asymptotes of the hyperbola.)

Empirical distributions that apparently have exponentially decreasing tails abound. The examples given in Barndorff-Nielsen (1977) include (log) diamond sizes from mining areas in South Africa, (log) personal income in Australia, and measurements of the velocity of light by Michelson.

There is a natural multivariate extension of (1) where the density is a hyperboloid. It turns out that the marginal and conditional distributions of multivariate hyperbolic distributions are not necessarily hyperbolic. Instead, hyperbolic distributions can be embedded into the class of *generalized hyperbolic distributions*, which is invariant under margining, conditioning, and affine transformations. Many important distributions are in this class or a limiting case of it. Namely, the Gaussian distribution, Student's t-distribution, the Laplace-distribution, the gamma distribution, and the reciprocal gamma distribution. (See Barndorff-Nielsen and Blaesild (1981)[p.20].)

Definition (Barndorff-Nielsen et al.; 1982)

A random variable $X \in \mathbf{R}^n$ is said to be distributed according to a normal variance-mean mixture with location μ , drift β , structure matrix Δ , and mixing distribution F , if there is a random variable u with a distribution F on $[0, \infty)$ and the conditional distribution of X under u is normal:

$$P^{X|u} = N_n(\mu + u\beta, u\Delta).$$

$\mu, \beta \in \mathbf{R}^n$, Δ is symmetric and positive definite, $\det(\Delta) = 1$. We denote this distribution by $NVMM(\mu, \beta, \Delta, F)$. (The condition $|\Delta| = 1$ makes the choice of parameters unique and excludes the case $\Delta = 0$, where any distribution F

on $[0, \infty)$ could be written as $NVMM(0, 1, 0, F)$. For $n = 1$, the parameter $\Delta = 1$ is irrelevant and we write $NVMM(\mu, \beta, F)$.

If ϕ is the characteristic function of F , the characteristic function ψ of $NVMM(\mu, \beta, \Delta, F)$ is given by

$$\psi(t) = e^{i\langle t, \mu \rangle} \phi(\langle \beta, t \rangle + \frac{i}{2} \langle t, \Delta t \rangle), \quad (2)$$

as is easily seen. A simple consequence of (2) is that the sum of i.i.d. NVMM variables with the same β and Δ is again a NVMM distribution:

$$NVMM(\mu, \beta, \Delta, F)^{*n} = NVMM(n\mu, \beta, \Delta, F^{*n}).$$

Furthermore, this class of mixtures is closed under margining, conditioning and affine transformations. Details are given in (Barndorff-Nielsen et al.; 1982, p.147).

Generalized hyperbolic distributions can now be defined as normal variance-mean mixtures where the mixing distribution is a *generalized inverse Gaussian distribution* $N^-(\lambda, \chi, \psi)$ that is defined by having the following density:

$$f(x) = \frac{(\psi/\chi)^{\lambda/2}}{2K_\lambda(\sqrt{\psi\chi})} x^{\lambda-1} \exp\left\{-\frac{1}{2}(\chi x^{-1} + \psi x)\right\}. \quad (x > 0)$$

(K_λ is a Bessel function. The parameter domain is $\lambda \in \mathbf{R}$, $\chi > 0$, $\psi > 0$. Additionally, $\chi = 0$ is allowed for $\lambda > 0$ and $\psi = 0$ is allowed for $\lambda < 0$.) Let

$$\begin{aligned} H(\lambda, \alpha, \beta, \mu, \delta, \Delta) &= NVMM(\mu, \beta\Delta, \Delta, N^-(\lambda, \delta^2, \kappa^2)), \\ \kappa^2 &= \alpha^2 - \langle \beta, \Delta\beta \rangle \end{aligned}$$

denote the generalized hyperbolic distribution. The density and characteristic function of the generalized hyperbolic distribution are given in Barndorff-Nielsen et al. (1982)[p.148]. For $\lambda = (n+1)/2$, the density simplifies to

$$\frac{(\kappa/\delta)^\lambda}{(2\pi)^{\lambda-1} 2\alpha K_\lambda(\delta\kappa)} e^{-\alpha\sqrt{\delta^2 + \langle x-\mu, \Delta^{-1}(x-\mu) \rangle} + \langle \beta, x-\mu \rangle},$$

whose logarithm is a hyperboloid in x .

Maximum likelihood estimation for hyperbolic distributions with parameters $H(1, \alpha, \beta, \mu, \delta)$ is discussed in Barndorff-Nielsen and Blaesild (1981), where it is noted that the computation of the MLE is "from a theoretical as well as from a computational point of view rather unpleasant", [p.29]. Namely, the log-likelihood function may have saddle points.

Back to the feature that gave rise to the definition of the hyperbolic distribution. It is shown in Barndorff-Nielsen and Blaesild (1981)[pp.34] that for the density h of the univariate generalized hyperbolic distribution

$$h(x; \lambda, \alpha, \beta, \mu = 0, \delta) \sim |x|^{\lambda-1} e^{-(\pm\alpha-\beta)x} \quad \text{as } x \rightarrow \pm\infty$$

for $\lambda > 0$. It is dubious whether one can see the effect of the power $|x|^{\lambda-1}$ in the log histogram. That is, generalized hyperbolic distributions very much “look like” hyperbolic distributions in the tails. That so much is known about generalized hyperbolic distributions makes it worthwhile investigating whether *normal variance-mean mixture distributions* implied by certain stochastic models are in fact *generalized hyperbolic*.

3 Normal Variance-Mean Mixture Distributions Generated by Stochastic Processes

If W is a one-dimensional Wiener-process and τ is an independent random time with distribution F , then

$$\mu + \beta\tau + W_\tau \sim \text{NVMM}(\mu, \beta, F).$$

Moreover, if $\{\tau_a\}$ is an increasing stochastic process which is independent of W , then $X_a := \mu + \beta\tau_a + W_{\tau_a}$ is a stochastic process whose increments are mixed normal. For financial markets, $a \rightarrow \tau_a$ can be thought of as a random time change mapping calendar time to “internal market time” or “operational time”. The generalized hyperbolic distribution with $\lambda = -\frac{1}{2}$ is now naturally obtained by letting τ_a be a certain first hitting time of an independent Wiener-process B :

$$\tau_a := \inf_{t \geq 0} \{\kappa t + B_t \geq a\delta\}.$$

τ_a is a Levy-process, i.e. it has independent increments. The increments are inverse Gaussian distributed: $\tau_b - \tau_a \sim N^-(\frac{1}{2}, ((b-a)\delta)^2, \kappa^2)$. Now, $X_a = W_{\tau_a}$ is a Levy-process with generalized hyperbolic increments. X was termed *Gaussian-inverse Gaussian process* and its Levy decomposition was given by Barndorff-Nielsen (1994). X is a pure jump process, and the Levy measure is not integrable at 0. That means, X has infinitely many (small) jumps in any time interval. X can be thought of as a limit of compound Poisson processes.

Normal variance-mean mixtures also appear in stochastic volatility models where log returns are – conditionally on some stochastic volatility σ – multivariate Gaussian:

$$R_k = \mu + \sigma_k^2 \beta + \sigma_k z_k.$$

σ is a stochastic process and z_k is Gaussian white noise (z_k i.i.d. $N_n(0, \Delta)$; $\beta, \mu \in \mathbf{R}^n$). If σ^2 is a stationary process with marginal distribution F and independent of (z_k) , then R is a stationary process with marginal distribution NVMM(μ, β, Δ, F).

One simple example of a stochastic model for $r_t := \sigma_t^2$ is

$$dr_t = (\omega - \theta r_t)dt + cr_t dW_t, \quad (3)$$

where W_t is an independent Wiener-process. In the framework of Nelson (1990), (3) is a weak limit of the volatility process in the GARCH(1,1)-model

$$r_{(k+1)h} = \omega_h + r_{kh}(\beta_h + c_h B_{kh}^2)$$

for $h \rightarrow 0$, and $B_{kh} \sim \text{i.i.d.} N(0, h)$, $\omega_h = \omega h$, $\beta_h = 1 - c\sqrt{h/2} - \theta h$, $c_h = c/\sqrt{2h}$. Nelson (1990) shows that the stationary distribution of (3) is reciprocal gamma if $1 + 2\theta/c^2 > 0$ and $\omega > 0$. That is, if $1/r_0 \sim \Gamma(1 + 2\theta/c^2, 2\omega/c^2)$, then r is (strictly) stationary.¹ Now, the stationary distribution of R is generalized hyperbolic $H(-(1 + 2\theta/c^2), \beta, \beta, \mu, 2\sqrt{\omega}/c)$. (Note that this one has tails $\sim |x|^{\lambda-1}$ for $x \rightarrow \infty$ and $\sim e^{2\beta x}$ for $x \rightarrow -\infty$ which is not quite what we observe.)

Yet another simple volatility model is

$$dr_t = (\omega - \theta r_t)dt + c\sqrt{r_t}dW_t.$$

Its stationary distribution is a gamma distribution: $\Gamma(2\omega/c^2, 2\theta/c^2)$. Hence, the stationary distribution of R is generalized hyperbolic with parameters $H(2\omega/c^2, \sqrt{4\theta/c^2 + \beta^2}, \beta, \mu, 0)$.

4 Conclusion

- (i) For stock returns, Gaussian tails ($\sim e^{-x^2}$) are too light while Pareto-stable tails ($\sim |x|^\alpha$) seem to be too heavy. The tails of generalized hyperbolic distributions ($\sim |x|^\alpha e^{-\lambda x}$) – which Barndorff-Nielsen calls *semi-heavy* – seem to fit just right to the distribution of (log) stock returns.

¹Let $\Gamma(\lambda, b)$ denote the gamma distribution with density $(b^\lambda/\Gamma(\lambda))x^{\lambda-1}e^{-bx}$, $x > 0$. Then $\Gamma(\lambda, b) = N^-(\lambda, 0, 2b)$ and, if $X \sim \Gamma(\lambda, b)$, then $X^{-1} \sim N^-(-\lambda, 2b, 0)$.

- (ii) Many stochastic volatility models produce – almost by definition – log returns whose stationary distribution is a *normal variance-mean mixture*. As the subclass of generalized hyperbolic distributions has several nice properties and is well-studied, it is worthwhile to investigate whether new (or old) stochastic volatility models produce log returns that are in fact generalized hyperbolic.

References

- Barndorff-Nielsen, O. (1977). Exponentially decreasing distributions for the logarithm of particle size, *Proc. R. Soc. London A* **353**: 401–419.
- Barndorff-Nielsen, O. (1994). Gaussian-inverse Gaussian processes and the modelling of stock returns, presented at the 2. Workshop on Stochastics and Finance 1994 in Berlin.
- Barndorff-Nielsen, O. and Blaesild, P. (1981). Hyperbolic distributions and ramifications: Contributions to theory and application, in C. T. et al. (ed.), *Statistical Distributions in Scientific Work*, Vol. 4, D. Reidel Publishing Company, pp. 19–44.
- Barndorff-Nielsen, O., Kent, J. and Sørensen, M. (1982). Normal variance-mean mixtures and z distributions, *International Statistical Review* **50**: 145–159.
- Eberlein, E. and Keller, U. (1994). Hyperbolic distributions in finance, Research report, Institut für Mathematische Stochastik, Universität Freiburg. presented at the 2. Workshop on Stochastics and Finance 1994 in Berlin.
- Fama, E. (1965). The behavior of stock market prices, *Journal of Business* **38**: 34–105.
- Harrison, J. and Pliska, S. (1981). Martingales and stochastic integrals in the theory of continuous trading, *Stochastic Processes and Applications* **11**: 215–260.
- Küchler, U., Neumann, K., Sørensen, M. and Streller, A. (1994). Stock returns and hyperbolic distributions, Discussion Paper 23, Sonderforschungsbereich 373, Humboldt-Universität zu Berlin. presented at the 2. Workshop on Stochastics and Finance 1994 in Berlin.
- Mahieu, R. J. (1995). *Financial Market Volatility*, Universitaire Pers Maastricht.

- Mandelbrot, B. and Taylor, H. (1967). On the distribution of stock price differences, *Operations Research* **15**: 1057+.
- Mittnik, S. and Rachev, S. (1993). Modeling asset returns with alternative stable distributions, *Econometric Reviews* **12**(3): 261–330.
- Nelson, D. B. (1990). ARCH models as diffusion approximations, *Journal of Econometrics* **45**: 7–38.