

A Service of



Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre

Werwatz, Axel; Müller, Christian

Working Paper
Simultaneous-equations models

SFB 373 Discussion Paper, No. 2000,55

## **Provided in Cooperation with:**

Collaborative Research Center 373: Quantification and Simulation of Economic Processes, Humboldt University Berlin

Suggested Citation: Werwatz, Axel; Müller, Christian (2000): Simultaneous-equations models, SFB 373 Discussion Paper, No. 2000,55, Humboldt University of Berlin, Interdisciplinary Research Project 373: Quantification and Simulation of Economic Processes, Berlin, https://nbn-resolving.de/urn:nbn:de:kobv:11-10047813

This Version is available at: https://hdl.handle.net/10419/62253

#### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

#### Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



# Simultaneuos-Equations Models

Axel Werwatz and Christian Müller

## 1 Introduction

A simultaneuos-equations model (SEQM) consists of several interdependent equations. Typically, these equations are not standard regression equations with an **endogenous variable** on the left-hand side and one or several **exogenous regressors** on the right-hand side that are independent of the error term. Rather, endogenous variables may also appear on the right hand side of the equations that comprise the SEQM.

But SEQMs are not merely a collection of equations with endogenous regressors. They are truly systems of equations in the sense that there are cross-equation relationships between the variables.

The well-known macroeconometric model of Klein (1950) is a good example to illustrate these points. The Klein's model consists of six equations, three **statistical equations** and three **identities**. The three statistical equations look like standard regression equations:

$$C_{t} = \alpha_{0} + \alpha_{1}P_{t} + \alpha_{2}P_{t-1} + \alpha_{3}(W_{t}^{p} + W_{t}^{g}) + \epsilon_{1t}$$

$$I_{t} = \beta_{0} + \beta_{1}P_{t} + \beta_{2}P_{t-1} + \beta_{3}K_{t-1} + \epsilon_{2t}$$

$$W_{t}^{p} = \gamma_{0} + \gamma_{1}Y_{t} + \gamma_{2}Y_{t-1} + \gamma_{3}A_{t} + \epsilon_{3t}$$

$$(1)$$

Here, the  $\alpha$ s,  $\beta$ s and  $\gamma$ s are unknown regression coefficients,  $\epsilon_{1t}$ ,  $\epsilon_{2t}$  and  $\epsilon_{3t}$  are unobservable error terms and all capital letters denote observable variables, whose meaning will be described below as necessary.

Klein's model is completed by the following three identities:

$$\begin{aligned}
 Y_t &= C_t &+ I_t &+ G_t \\
 P_t &= Y_t &+ T_t &- W_t^p \\
 K_t &= K_{t-1} &+ I_t 
 \end{aligned}
 \tag{2}$$

These identities neither include unknown coefficients nor error terms. They hold "by definition". Nonetheless, they are an integral part of the model.

The first equation of (2), for instance, says that total spending  $Y_t$  in an economy in year t is the sum of private consumption spending  $C_t$ , investment spending  $I_t$  and government spending  $G_t$ . This is an accounting relationship. Similarly, the second equation of (2) states that we obtain private profits  $P_t$  if we subtract from total spending  $Y_t$  indirect taxes  $T_t$  and the total wage bill of private enterprises  $W_t^p$ . These identities introduce interdependencies between the variables of the statistical equations (1).

Note, for instance, that  $C_t$  depends on  $W_t^p$  via the first equation of (1) and that  $W_t^p$  depends on  $Y_t$  via the third equation of (1). But  $Y_t$  depends on  $C_t$  though the first identity in (2) which implies that  $W_t^p$  depends on  $Y_t$ . In this way, the first and third equation of (1) are interdependent or simultaneous. This relatedness is not a result of some relationship between the error terms  $\epsilon_{1t}$  and  $\epsilon_{3t}$  of these equations. Rather, it is a result of the equations in (1) and (2) truly being a system of equations with various cross-equation relationships between the observable variables of the system. This simulateneity has important consequences if we want to consistently estimate the unknown coefficients in (1).

## 2 Estimating Simultaneous-Equations Models

### 2.1 Identification

The equations of Klein's SEQM defined by (1) and (2) are motivated by economic theory. They are an attempt to write down in a parsimonious way the "structure" of the economy from an aggregate, macro perspective. The parameters of these equations are thus the proper targets of estimation: after all, they supposedly tell us "how the economy works". In the following sections, we will briefly discuss two approaches of estimating the parameters of a structural SEQM such as Klein's model.

A necessary condition for consistent estimation is identification. Identification in SEQMs is a serious issue and warrants an extended treatment that can't be provided in the format of this chapter. But any good econometrics textbook features an extensive discussion of this matter. See, for instance, Greene (1998) or Goldberger (1991).

Intuitively speaking, identification is difficult because of the interrelatedness of the variables, which makes it hard to disentangle their various relationships. A necessary and easily verified condition for identification in SEQMs is that the number of exogenous variables excluded from any equation must be as least as large as the number of endogenous variables included in that equation ("order condition").

In Klein's model of equations (1) and (2), the endogenous variables are  $C_t$ ,  $I_t$ ,  $W_t^p$ ,  $Y_t$ ,  $K_t$  and  $P_t$  (note that the latter three are endogenous despite not being a left-hand side variable of any statistical equation). All other variables are exogenous or predetermined. You may verify that in the first equation of (1) there are two endogenous ( $C_t$  and  $P_t$ ) and six excluded exogenous variables ( $K_{t-1}$ ,  $Y_{t-1}$ ,  $A_t$ ,  $G_t$ ,  $T_t$  and  $W_t^g$ ). Hence, the order equation is satisfied for this equation. Greene (1998) shows that the condition is satisfied for the entire system (as well as other, sufficient conditions for identification).

### 2.2 Some Notation

Let us write down the M statistical equations (i.e., the equations that have unknown coefficients and error terms) of the structural form (i.e., the equations as they are suggested by economics) of a SEQM:

$$\begin{array}{lclclclcl} \mathbf{y}_1 & = & \mathbf{Y}_1^T \ \beta_1 & + & \mathbf{X}_1^T \ \gamma_1 & + & \varepsilon_1 \\ \vdots & & & \vdots & & & \\ \mathbf{y}_M & = & \mathbf{Y}_M^T \ \beta_M & + & \mathbf{X}_M^T \ \gamma_M & + & \varepsilon_M. \end{array}$$

Focusing on the first equation,  $y_1$  is the left-hand side variable,  $Y_1$  is the vector of endogenous right-hand side variables,  $x_1$  is the vector of exogenous or predetrmined right-hand side variables and  $\beta_1$  and  $\gamma_1$  are the objects of interest: the unknown structural parameters of the model.

We can illustrate the notation by using, say, the M-th equation of Klein's model introduced above: M=3 (the coefficients of the identities in (2) are known and need not be estimated),  $y_M=W_t^p$ ,  $Y=Y_t$ ,  $\beta_M=\gamma_1$ ,  $X_M^T=\begin{bmatrix}1 & Y_{t-1} & A_t\end{bmatrix}$ ,  $\gamma_M=\begin{bmatrix}\gamma_0 & \gamma_2 & \gamma_3\end{bmatrix}^T$  and  $\varepsilon_M=\varepsilon_{3t}$ .

Suppose that we have T observations of all the variables of the model. Then we can write down the SEQM in terms of the data matrices and vectors as

follows:

$$\begin{array}{rclcrcl} y_1 & = & Y_1\beta_1 & + & X_1\gamma_1 & + \epsilon_1 \\ \vdots & & & \vdots & & \vdots \\ y_M & = & Y_M\beta_M + & X_M\gamma_M & + \epsilon_M. \end{array}$$

Using again the M-th equation to illustrate the notation,  $y_M$  and  $\epsilon_M$  are  $T \times 1$  vectors whereas  $Y_M$  and  $X_M$  are matrices with n rows that have as many columns as  $Y_M^T$  and  $X_M^T$ , respectively.

Defining  $Z_m = (Y_1, X_1)$  and  $\delta_m = (\beta_M, \gamma_M)$  for m = 1, ..., M, we can write the system even more compactly as

$$y_1 = Z_1 \delta_1 + \epsilon_1$$

$$\vdots \qquad \vdots$$

$$y_M = Z_M \delta_M + \epsilon_M.$$
(3)

Estimating  $\delta_1, \ldots, \delta_M$  by applying ordinary least squares to each equation is not a consistent estimation procedure as some of the right-hand side variables are endogenous and therefore correlated with the error terms.

For instance, in the second equation of (1),  $P_t$  is correlated with the error term. This can be seen by substituting the first identity of (2) for  $X_t$  into the second identity of (2) and observing that  $P_t$  is a linear function of  $I_t$ .

## 2.3 Two-Stage Least Squares

Endogenous regressors can be treated by instrumenting them and this is precisely what the method of two-stage least squares does.

1. In the first stage, the endogenous regressor is regressed by ordinary least squares on all exogenous regressors and the predicted values of this regression are obtained. Denoting the matrix of observations of **all** exogenous or predetermined variables of the SEQM as X we can write the matrix of predicted values of the endogenous regressors of the m-th equation as

$$\hat{Y}_m = X\{(X^T X)^{-1} X^T Y_m\} \tag{4}$$

2. In the second stage, the two-stage least squares estimator for equation m is obtained by regressing  $y_m$  on the fitted values  $\hat{Y}_m$  and the matrix of exogenous or predetermined regressors of equation  $m, X_m$ .

Using the notation of (3), Greene (1998) shows that the two-stage least squares estimator of the parameters of equation m can be written as

$$\hat{\delta}_{m,2SLS} = \{ (Z_m^T X)(X^T X)^{-1} (X^T Z_m) \}^{-1} Z_m X (X^T X)^{-1} X^T y_m.$$

$$= \{ \widehat{Z}_m^T \widehat{Z}_m \}^{-1} \widehat{Z}_m^T y_m$$
(5)

This procedure, applied to all equations  $m=1,\ldots,M$ , is consistent but not efficient. This is because it does not exploit the fact that the equations form a system or interdependent equations but rather estimates the equations one-by-one. In the following section, we will describe the three-stage least squares estimation procedure which builds on but also improves upon two-stage least squares.

## 2.4 Three-Stage Least Squares

Let us rewrite (3) such that the system nature of the SEQM is even more evident:

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_M \end{pmatrix} = \begin{pmatrix} Z_1 & 0 & \cdots & 0 \\ 0 & Z_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & Z_M \end{pmatrix} + \begin{pmatrix} \delta_1 \\ \delta_2 \\ \vdots \\ \delta_M \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_M \end{pmatrix}$$
(6)

(where y, Z and  $\epsilon$  all have  $T \times M$  rows) or simply as

$$y = Z\delta + \epsilon \tag{7}$$

Using this "system" notation, we can write the two-stage least squares estimator (5) for all M equations very compactly as

$$\widehat{\delta}_{2SLS} = \{\widehat{Z}^T \widehat{Z}\}^{-1} \widehat{Z}^T y,\tag{8}$$

where

$$\widehat{Z} = \begin{pmatrix}
X(X^{T}X)^{-1}X^{T}Z_{1} & 0 & \cdots & 0 \\
0 & X(X^{T}X)^{-1}X^{T}Z_{2} & \cdots & 0 \\
\vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & X(X^{T}X)^{-1}X^{T}Z_{M}
\end{pmatrix}$$

$$= \begin{pmatrix}
\widehat{Z}_{1} & 0 & \cdots & 0 \\
0 & \widehat{Z}_{2} & \cdots & 0 \\
\vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & \widehat{Z}_{M}
\end{pmatrix}$$
(9)

Written in this notation, the two-stage least-squares estimator looks like a "system" estimator but it is really equation-by-equation instrumental variables regression.

As remarked earlier, this estimator is consistent but it is not efficient. A more efficient estimator is suggested by looking at the covariance matrix of the long residual vector  $\epsilon$  of (7).

$$E[\epsilon \epsilon^{T}] = \bar{\Sigma} = \begin{pmatrix} \sigma_{11}I & \sigma_{12}I & \cdots & \sigma_{1M}I \\ \sigma_{21}I & \sigma_{22}I & \cdots & \sigma_{2M}I \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{M1}I & \sigma_{M2}I & \cdots & \sigma_{MM}I \end{pmatrix} = \Sigma \otimes I, \tag{10}$$

where I is an  $T \times T$  identity matrix,  $\otimes$  denotes the Kronecker product and  $\Sigma$  is the covariance matrix of the error terms  $\epsilon_1, \epsilon_2, \ldots, \epsilon_M$ . This non-spherical covariance matrix suggestes to improve upon (8) by using generalized least squares. This is precisely the idea behind the three-stage least squares estimator.

Suppose we knew the covariance matrix  $\Sigma$ . Then a generalized least squares, instrumental variables estimator could be computed as

$$\delta_{GLS} = \{\widehat{Z}^T(\Sigma^{-1} \otimes I)\widehat{Z}\}^{-1}\widehat{Z}^T(\Sigma^{-1} \otimes I)y. \tag{11}$$

In practice,  $\Sigma$  is unknown and has to be estimated. A consistent estimator of element (i, j) of  $\Sigma$  is based on the sum of squares of the residuals of the *i*-th and *j*-th equation after estimating each equation by two-stage least squares:

$$\hat{\sigma}_{ij} = \frac{(y_i - Z_i \hat{\delta}_i)^T (y_j - Z_j \hat{\delta}_j)}{T}, \tag{12}$$

where  $\hat{\delta}_i$  and  $\hat{\delta}_j$  are the two-stage least squares estimator for equations i and j as defined in (5). Estimating each element of  $\Sigma$  in this way defines the estimator  $\hat{\Sigma}$  of  $\Sigma$  and allows to compute the feasible generalized lest squares estimator:

$$\widehat{\delta}_{3SLS} = \{\widehat{Z}^T(\widehat{\Sigma}^{-1} \otimes I)\widehat{Z}\}^{-1}\widehat{Z}^T(\widehat{\Sigma}^{-1} \otimes I)y. \tag{13}$$

An estimator of the asymptotic covariance matrix of (13) is given by

$$\widehat{\operatorname{Asy. Var}}(\widehat{\delta}_{3SLS} = \{\widehat{Z}^T(\widehat{\Sigma}^{-1} \otimes I)\widehat{Z}\}^{-1}$$
(14)

Summing up, the three-stage least squares estimator of  $\delta$  in (7) is obtained by carrying out the following three-steps:

- 1. The first stage is identical to the two-stage procedure: instruments for the endogenous regressors are computed as the predicted values of an ordinary least squares regression of each endogenous regressor on all exogenous regressors.
- 2. In the second stage, the two-stage least squares estimator for each equation m is computed and the residuals are used according to (12) to obtain an estimate of  $\Sigma$ , the covariance matrix of the error terms of the SEQM.
- 3. In the third stage, the estimate of  $\Sigma$  is used to calculate the generalized least squares estimator defined in (13) and an estimate of its covariance matrix as described in (14)

## 2.5 Computation

```
{d3sls, cov3, d2sls} = seq(seqlist1, seqlist2)
    estimates a SEQM via two-stage and three-stage least squares
```

The quantlet seq is all you need to estimate a SEQM with XploRe. It belongs to the metrics quantlib which you have to load before you can use seq:

```
library("metrics")
```

seq requires two inputs and will return two vectors and a matrix along with an ANOVA-type table in the output window. The two inputs are lists. The first list (seqlist1) is a list of matrices that contain the data. The second list (seqlist2) is a list of string vectors that contain variable names.

#### seqlist1

If there are M statistical equations (i.e., equations with unknown coefficents and an error term) in the system then  $\mathtt{seqlist1}$  is a list of M+2 matrices. The first matrix consists of the cocatenated vectors of observations of the left-hand side variables. The next M matrices contain the observations of the right-hand side variables of the M statistical equations of the system. The last and (M+2)nd matrix contains the observations of all exogenous and predetermined variables of the system. That is, the M+2 matrices are:

```
\begin{array}{lll} 1. & & (y_1, y_2, \dots, y_M) \\ 2. & & Z_1 \\ 3. & & Z_2 \\ \vdots & & \vdots \\ (M+1). & Z_M \\ (M+2). & X \end{array}
```

#### seqlist2

is a list of M+2 string vectors. Each vector contains the names of the variables that form the corresponding matrix in seqlist1. For instance, the first string vector of seqlist2 contains the names of the left-hand side variables of the system whereas the last string vector contains the names of all exogenous and predetermined variables of the system.

The XploRe code for the Klein's model will serve as an example:

```
; reading in the data
data=read("klein.dat")

; getting rid of the missing values of the lagged variables
data=data[2:rows(data),]

; assigning the columns of "data" to the variables
C=data[,2]
P=data[,3]
Wp=data[,4]
I=data[,5]
K1=data[,6]
Y=data[,7]
Wg=data[,8]
```

```
G=data[,9]
T=data[,10]
W=data[,11]
P1=data[,12]
Y1=data[,13]
A=data[,14]
; preparing the matrices that form seqlist1
; first matrix: concatenated vectors
                                 of left-hand side variables
y=C~I~Wp
; a column of "1"s for constant term
ONE=matrix(rows(data),1)
; second through (M+1)st matrix:
                       matrices of right-hand side variables
z1=P~P1~W~ONE
z2=P~P1~K1~ONE
z3=Y~Y1~A~ONE
; (M+2)nd matrix: matrix of instruments
                      (exogenous and predetermined variables)
x=ONE~G~T~Wg~A~P1~K1~Y1
; forming a list of matrices
seqlist1=list(y,z1,z2,z3,x)
; now we will prepare the string vectors that will form
; seglist2 vector with names of left-hand side variables
yl="C"|"I"|"Wp"
; vectors with names of right-hand side variables
                                     of equations 1 though M
11="P"|"P1"|"W"|"ONE"
12="P"|"P1"|"K1"|"ONE"
13="E"|"E1"|"A"|"ONE"
; vector with names of instruments
xl="ONE"|"G"|"T"|"Wg"|"A"|"P1"|"K1"|"Y1"
; putting the string vectors into a list
seqlist2=list(y1,11,12,13,x1)
```

```
; finally, we call the seq-quantlet \{d1,s,d2\}=seq(seqlist1,seqlist2)
```

Q seq01.xpl

seq produces two kinds of output: ANOVA-style tables in the output window and coefficient estimates and their estimated standard errors as vectors or matrices.

There is a table of estimation results both for the two-stage (shown first) and three-stage least squares procedures (shown last). Both tables are designed to be easily readable and self-explanatory. We hope that you are going to agree after looking at the three-stage least squares table for the Klein's model:

[ 1,]	"=======			========"
			es estimates"	
•			======================================	=======""
	"EQ dep.		κ∠ · =========	"
•	" 1 C		0.980	11
•	" 2 I			II .
-	" 3 Wp		0.986	II .
•	-			======="
			Std.Err.	t"
[11,]	"			"
[12,]	"P	0.125	0.108	1.155"
[13,]	"P1	0.163	0.100	1.624"
[14,]	''W	0.790	0.038	20.826"
[15,]		16.441	1.305	12.603"
[16,]	"			"
[17,]	"P	-0.013	0.162	-0.081"
[18,]	"P1	0.756	0.153	4.942"
[19,]	"K1	-0.195	0.033	-5.990"
[20,]	"ONE	28.178	6.794	4.148"
[21,]	"			"
[22,]	"E	0.400	0.032	12.589"
[23,]	"E1	0.181	0.034	5.307"
[24,]	'' A	0.150	0.028	5.358"
[25,]		1.797	1.116	1.611"
•				=======""
	"INSTRUMENT		$\mathtt{Std}.\mathtt{Dev}.$	11
Γ28.]	"			"

[29,]	"ONE	1.000	0.000"
[30,]	"G	9.914	3.910"
[31,]	"T	6.805	2.032"
[32,]	"Wg	5.119	1.957"
[33,]	"A	0.000	6.205"
[34,]	"P1	16.376	4.028"
[35,]	"K1	200.495	9.919"
[36,]	"Y1	57.986	8.919"
[37,]	"=======	=========	=======================================

Besides showing the estimation results in the output window you can also access some of them as vectors or matrices. Specifically, the following quantities are returned by seq:

#### d3sls

the vector of three-stage least squares estimates of the elements of  $\delta$  in (7). It is computed according to (13).

#### cov3

the estimated covariance matrix of the three-stage least squares estimator, computed according to (14).

#### d2s1s

the vector of two-stage least squares estimates of the elements of  $\delta$  in (7). It is obtained by computing (5) for each equation and stacking-up the resulting vectors.

In the following section, we will use seq to estimate a money-demand system.

## 3 Application: Money Demand

Economists often refer to the money stock as one important determinant of the price level. Therefore, the evolution of the amount of money in the economy is also a focus of monetary policy analysis. A convenient tool for this is to look at the so called demand for money. The recent economic literature analyzes the long-run demand for money (denoted m) as a function of aggregated income (y,) short and long term interest  $(i^s, i^l)$  and inflation rates within a cointegration framework. If there is more than one cointegration relationship and/or the relationship of interest helps to explain more than just one variable, these relationships are estimated more efficiently within a system rather than as a single equation (see (Ericsson 1999) on that). The explanatory variables considered in the money demand equation might cointegrate not only to a money demand function but also to a stationary spread between long and short term interest rates and a stationary real interest rate. Therefore, in a study about European money demand, Müller and Hahn (2000) applied a system specification to determine whether or not there exists a stationary relationship between the money stock, aggregated income, 3-months interest rates, government bond yield and a measure of European inflation. All data are weighted sums of the series of each of the eleven countries except for the price measure which has been obtained as the ratio of nominal and real income. In case of the interest rates the weights are real income shares and in the cases of money and income the official EURO rates have been used.

Using a system approach suggests to consider a reduced form regression, where no endogenous variables may enter any of the equations on the right hand side. In contrast to that, the change in money stock is often considered to depend e.g. on the current change in inflation (Lütkepohl and Wolters 1998) and the same is true for relationship between the short and the long term interest rates. That's why the reduced form regression is used to identify the long-run relationships (cointegration relationships) while in a second step the model is re-written to yield a structural form, as described above. Thus, the structural SEQM is:

$$\Delta(m-p)_{t} = \alpha_{1,1} + \alpha_{1,2}\Delta(m-p)_{t-1} + \alpha_{1,3}\Delta y_{t-1} + \alpha_{1,4}\Delta i_{t-1}^{l} + \alpha_{1,5}\Delta i_{t-1}^{s} + \alpha_{1,6}\Delta^{2}p_{t-1} + \alpha_{1,7}\Delta^{2}p_{t} + \alpha_{1,8}ec1_{t-1} + \varepsilon_{1,t}$$
(15)

$$\Delta y_{t} = \alpha_{2,1} + \alpha_{2,2} \Delta (m-p)_{t-1} + \alpha_{2,3} \Delta y_{t-1} + \alpha_{2,4} \Delta i_{t-1}^{l} + \alpha_{2,5} \Delta i_{t-1}^{s} + \alpha_{2,6} \Delta^{2} p_{t-1} + \alpha_{1,8} e c 1_{t-1} + \alpha_{2,9} e c 2_{t-1} + \varepsilon_{2,t}$$
(16)

$$\Delta i_t^l = \alpha_{3,1} + \alpha_{3,2} \Delta (m-p)_{t-1} + \alpha_{3,3} \Delta y_{t-1} + \alpha_{3,4} \Delta i_{t-1}^l + \alpha_{3,5} \Delta i_{t-1}^s + \alpha_{3,6} \Delta^2 p_{t-1} + \varepsilon_{3,t}$$
(17)

$$\Delta i_t^s = \alpha_{4,1} + \alpha_{4,2} \Delta (m-p)_{t-1} + \alpha_{4,3} \Delta y_{t-1} + \alpha_{4,4} \Delta i_{t-1}^l + \alpha_{4,5} \Delta i_{t-1}^s + \alpha_{4,6} \Delta^2 p_{t-1} + \alpha_{4,10} \Delta i_t^l + \varepsilon_{4,t}$$
(18)

$$\Delta^{2} p_{t} = \alpha_{5,1} + \alpha_{5,2} \Delta (m-p)_{t-1} + \alpha_{5,3} \Delta y_{t-1} + \alpha_{5,4} \Delta i_{t-1}^{l} + \alpha_{5,5} \Delta i_{t-1}^{s} + \alpha_{5,6} \Delta^{2} p_{t-1} + \alpha_{5,9} ec2_{t-1} + \varepsilon_{1,t}$$

$$(19)$$

$$ec1_t = (m-p)_t - 1.574y_t + 3.405i_t^l - 2.061i_t^s$$
 (20)

$$ec2_t = i^l - 4\Delta p \tag{21}$$

Here,  $\Delta = 1 - L$  where L is the backshift operator (i.e.,  $\Delta y_{t-1} = (1 - L)y_{t-1} = y_{t-1} - y_{t-2}$ ). It can be easily verified that the order condition holds and all parameters are identified. In our case the endogenous variables are  $\Delta m - p, \Delta y_t, \Delta i_t^l, \Delta i_t^s$  and  $\Delta^2 p_t$  (all variables except the interest rates in logarithms). There are no purely exogenous variables but predetermined (lagged endogenous) variables only. The error correction terms are given by the identities (20) and (21). They are the results of the first step, the reduced form regression. Thus, we re-estimate the model including the short-run adjustment parameters by 3SLS under more general assumptions about the contemporaneous structure than in the reduced rank model.

We have used the following XploRe code to estimate the parameters of equations (15) to (19):

```
; reading in the data
z=read("eu.raw")
; getting rid of missing values due to lagged variables
z=z[4:rows(z),]
; assigning columns of z to variable names
dmp = z[,4]
```

```
dmp1
         = z[,5]
         = z[,7]
dу
         = z[,8]
dy1
         = z[,12]
d2p
         = z[,13]
d2p1
dil
         = z[,15]
dil1
         = z[,16]
         = z[,18]
dik
dik1
         = z[,19]
         = z[,21]
ec11
ec21
         = z[,23]
; creating the matrices for seqlist1
lhs=dmp~dy~dil~dik~d2p
one= matrix(rows(z),1)
z1=one~dmp1~dy1~dil1~dik1~d2p1~d2p~ec11
z2=one~dmp1~dy1~dil1~dik1~d2p1~ec11~ec21
z3=one~dmp1~dy1~dil1~dik1~d2p1
z4=one~dmp1~dy1~dil1~dik1~d2p1~dil
z5=one~dmp1~dy1~dil1~dik1~d2p1~ec21
x=one^dmp1^dy1^dil1^dik1^d2p1^ec11^ec21
; forming seqlist1 as a list of matrices
seqlist1=list(lhs,z1,z2,z3,z4,z5,x)
; creating list of string vectors
yl="dmp"|"dy"|"dzl"|"dzk"|"d2p"
zl1="one"|"dmp1"|"dy1"|"dil1"|"dik1"|"d2p1"|"d2p"|"ec11"
zl2="one"|"dmp1"|"dy1"|"dil1"|"dik1"|"d2p1"|"ec11"|"ec21"
zl3="one"|"dmp1"|"dy1"|"dil1"|"dik1"|"d2p1"
z14="one"|"dmp1"|"dy1"|"dil1"|"dik1"|"d2p1"|"dil"
z15="one"|"dmp1"|"dy1"|"dil1"|"dik1"|"d2p1"|"ec21"
xl="one"|"dmp1"|"dy1"|"dil1"|"dik1"|"d2p1"|"ec11"|"ec21"
; forming seqlist2 as a list of string vectors
seqlist2=list(y1,z11,z12,z13,z14,z15,x1)
; finally, calling seq to estimate the model
{d3sls,cov3,d2sls}=seq(seqlist1,seqlist2)
```

Q seq02.xpl

These lines of XploRe code yield the following table in the output window (table of two-stage least squares estimates has been omitted). It shows for each equation the left-hand side variable (top panel), three-stage least squares coefficient estimates, estimated standard errors and t-values. The bottom panel

of the table lists the predetermined and exogenous variables that were used as instruments in the estimation procedure, along with their sample means and standard deviations.

[ 1,] [ 2,]	"====== " 3stag	======== e Least-squares	s estimates"	======="
[ 3,] [ 4,] [ 5,]	" EQ dep	. var.	R2"	
[ 6,]	" 1 dmp		0.643 "	
[7,]	"2 dy		0.506 "	
[8,]	" 3 dzl		0.279 "	
[ 9,]	"4 dzk		0.079 "	
[10,]	" 5 d2p		0.299 "	
[11,]	"======	=========		======="
[12,]	"VARIABLE	Coef.Est.	Std.Err.	t"
[13,]	"			"
[14,]	"one	-0.050	0.074	-0.684"
[15,]	"dmp1	0.680	0.113	6.023"
[16,]	"dy1	0.124	0.114	1.089"
[17,]	"dil1	-0.126	0.153	-0.821"
[18,]	"dik1	0.029	0.149	0.195"
[19,]	"d2p1	-0.897	0.541	-1.658"
[20,]	"d2p	-1.421	1.078	-1.319"
[21,]	"ec11	-0.018	0.026	-0.669"
[22,]	"			"
[23,]	"one	0.184	0.054	3.403"
[24,]	"dmp1	0.270	0.087	3.082"
[25,]	"dy1	0.393	0.106	3.694"
[26,]	"dil1	0.016	0.129	0.124"
[27,]	"dik1	0.041	0.098	0.418"
[28,]	"d2p1	0.922	0.242	3.809"
[29,]	"ec11	0.061	0.019	3.262"
[30,]	"ec21	-0.159	0.054	-2.928"
[31,]	"			"
[32,]	"one	-0.002	0.001	-1.980"
[33,]	"dmp1	0.148	0.098	1.516"
[34,]	"dy1	0.192	0.121	1.590"
[35,]	"dil1	0.425	0.138	3.075"
[36,]	"dik1	-0.092	0.103	-0.894"

	"d2p1	0.136	0.261	0.522"
[38,]	"			"
[39,]	"one	-0.003	0.002	-1.698"
[40,]	"dmp1	0.120	0.174	0.686"
[41,]	"dy1	0.365	0.221	1.656"
[42,]	"dil1	0.400	0.374	1.069"
[43,]	"dik1	0.108	0.157	0.687"
[44,]	"d2p1	-0.022	0.371	-0.060"
[45,]	"dil	-0.270	0.762	-0.354"
[46,]	"			"
[47,]	"one	-0.004	0.002	-2.472"
[48,]	"dmp1	0.030	0.047	0.646"
[49,]	"dy1	0.009	0.057	0.154"
[50,]	"dil1	-0.077	0.068	-1.144"
[51,]	"dik1	0.062	0.048	1.296"
[52,]	"d2p1	-0.323	0.125	-2.574"
[53,]	"ec21	0.072	0.028	2.596"
[54,]	"========		=======================================	-======""
	"INSTRUMENTS	Mean	Std.Dev.	II .
[56,]	"			"
[57,]	"one	1.000	0.000"	
[58,]	"dmp1	0.000	0.005"	
[59,]	"dy1	0.006	0.005"	
[60,]	"dil1	-0.001	0.004"	
[61,]	"dik1	-0.002	0.006"	
[62,]	"d2p1	0.000	0.002"	
[63,]	"ec11	-2.813	0.027"	
[64,]	"ec21	0.052	0.010"	
[65,]	"========		==============	"

Interpreting the results, two groups of estimators are of particular interest. These are first the structural or contemporaneous explanatory variables' parameters ( $\alpha_{4,10}$  and  $\alpha_{1,7}$ ) and, second the parameters of the error correction terms ( $\alpha_{.,8}$  and  $\alpha_{.,9}$ ), the so called short-run adjustment parameters.

Within the first group, we obtain a negative relationship between real money growth and inflation growth, which is indicated by the coefficient of -1.42. Its sign does not come as a surprise since an increase in inflation will naturally depreciate the value of nominal money stock. The coefficient does not seem to be statistically significant however, which is indicated by the marginal probability

of 0.187. Similarly the changes in the long-term interest rate do not seem to have a significant impact on short rate movements of the same period.

The second group of interest provides some insight into the feasibility and effects of monetary policy as well as to some extent into some basic economic relationships. To start with, the error correction term which is given in (20) and labeled as the long-run money demand enters the money and income growth equations (15) and (16). This term indicates what effect money demand has on the respective variables in excess of the long-run equilibrium. In the first equation we assumed it to lead to a slow-down in money growth. This feature should be present if one expects money to be demanded in quite the same way as many other ordinary commodities. Thus, in such a case we would observe an inherent tendency to restore equilibrium. The estimation results suggests however that this adjustment does not take place. This is because although the corresponding coefficient yields the correct sign it has too large a standard error compared to its magnitude. When the true coefficient is zero then the we would also have to assume that there is no money demand in Europe altogether and the cointegrating relationship should better be rewritten in such a way that it is normalised on a variable in whose equation the error correction term enters significantly. This could be the income equation for example. Sticking to the interpretation of a long-run money demand equilibrium we notice that excess demand of real money will lead to higher income growth in the next period as indicated by  $\alpha_{2.8}$ . Of course, re-formulating the ec1 term does not affect the significance of the coefficient but it could change the sign and will change magnitude and economic interpretation, which will not be done here since we are investigating the hypothesis of the existence of a money demand.

The second error correction term has an interpretation as a real interest rate. When real interest rates are high, the respective coefficient in the income equation indicates that income growth will be less in the following period. This, too, is economically reasonable, because credits are more expensive in that case. Already in the first step, in the reduced form estimation, we found no evidence of an endogenous tendency for the long-term interest rate to adjust to deviations from the long-run real interest equilibrium level. Therefore the error correction term for real interest rates have not been included in eq. (19). Instead, as the corresponding coefficient of the current estimation  $(\alpha_{2,9})$  implies also, these deviations may help to predict future inflation.

Since we used the error correction terms obtained in the reduced form regression and applied the zero adjustment coefficient restrictions identified in this first step, there was not much more to learn about the effect of excess money

demand on prices, say. Therefore the additional insight from this 3SLS estimation is mainly the sensitivity of the short-run adjustment estimates when explicit structural assumptions enter the model. It has to be pointed out however, that no final conclusions can be drawn yet because as the t-statistic of the additional structural explanatory variables indicate, their inclusion might not have contributed much to explain the underlying data generating process. That's why it is not quite clear which of variables are really part of this process and which are not. In order to find out more about that some model selection procedures could be applied. A natural extension in that direction would be e.g. to systematically exclude unnecessary variables due to some criteria like t-values, F-statistics, Akaike or Schwartz criteria to obtain more efficient estimates of the remaining **true** model.

## References

- Ericsson, N. (1999). Empirical modelling of money demand, in: H. Lütkepohl and J. Wolters (eds.), *Money Demand in Europe*, Physika-Verlag, Heidelberg, pp. 29–49.
- Greene, W. H. (1998). *Econometric Analysis*, 3rd edn, Prentice Hall, New York.
- Goldberger, A. S. (1991). A Course in Econometrics, Harvard University Press, Cambridge.
- Klein, L. R. (1950). Economic Fluctuations in the United States, 1921–1941, Cowles Commission for Research in Economics, New Haven.
- Lütkepohl, H. and Wolters, J. (1998). A money demand system for German M3, Empirical Economics 23(3): 371–386.
- Müller, C. and Hahn, E., (2000). Money Demand in Europe: Evidence from the Past SFB 373 Discussion Paper, Humboldt University Berlin.